



Gastrointestinal parasites in Danish cats

Thamsborg, Stig Milan; Starostka, Trine Parsberg; Larsen, Birgit ; Enemark, Heidi

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Thamsborg, S. M., Starostka, T. P., Larsen, B., & Enemark, H. (2015). *Gastrointestinal parasites in Danish cats*. Abstract from 6th Conference of the Scandinavian-Baltic Society for Parasitology (CSBSP6), Uppsala, Sweden.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

GASTROINTESTINAL PARASITES IN DANISH CATS

Stig M. Thamsborg¹, Trine P. Starostka², Birgit Larsen², Heidi L. Enemark²

¹*Department of Veterinary Disease Biology, University of Copenhagen, Denmark.*

²*National Veterinary Institute, Technical University of Denmark, Denmark.*

smt@sund.ku.dk

Introduction

Gastrointestinal parasites, including protozoa, are known to cause diarrhea in animals and humans. The objective of this study was to investigate the prevalence and genetic variation of *Cryptosporidium* spp., *Giardia* spp. in cats from Zealand and to study possible risk factors associated with these infections.

Materials and Methods

Faecal samples and questionnaire data were collected from a total of 315 cats including 284 cats from 2 veterinary clinics and 2 shelters (study population), and 34 cats from 3 catteries (cases). Cysts and oocysts were quantified by immunofluorescence microscopy (IFA), and all samples positive by IFA underwent PCR amplification and sequencing of the 18S rDNA, *hsp70* and *gdh* genes.

Results

Within the study population (n=284) the prevalence of *Giardia* and *Cryptosporidium* was 7.0% (n=20) and 6.7% (n=19), respectively. Eight of the cats were infected with both parasites (2.8 %). The prevalence among cats (n=34) in catteries was 47.1 % (n=16) for *Giardia* and 35.3 % (n=12) for *Cryptosporidium*. Nine of the cats were infected with both parasites (26.5 %). Significant risk factors associated with *Cryptosporidium* spp. infection were: *Giardia* spp. infection (study population), age (cases) and “cause for the veterinary consultation” (cases). Risk factors significantly associated with *Giardia* spp. infection were: *Cryptosporidium* spp. infection (cases), outdoor access (study population) and “other animals in the household” (study population). A significant positive association between level of (oo-)cyst excretion and clinical illness/diarrhea was found. Catteries had significantly more *Giardia* and *Cryptosporidium* infected cats compared to veterinary clinics and shelters. Molecular analyses of *Giardia* isolates revealed predominantly *Giardia duodenalis* assemblage F, but *Giardia duodenalis* assemblage A was also detected, whereas all *Cryptosporidium* isolates were identified as *Cryptosporidium felis*.

Conclusion

Infections with *Cryptosporidium* and *Giardia* represent a real problem among cats and especially in catteries where it, despite treatment and hygiene measures, can be difficult to control these infections. *C. felis* and *G. duodenalis* Assemblage A are capable of infecting humans, and close contact with cats is a potential source of infection, although the risk is currently considered to be low.