



## Inhibition studies of natural resin acids to *Clostridium perfringens* and *Escherichia coli* O149

Roy, Krisna; Lyhs, Ulrike; Pedersen, Karl

*Publication date:*  
2015

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Roy, K., Lyhs, U., & Pedersen, K. (2015). *Inhibition studies of natural resin acids to Clostridium perfringens and Escherichia coli O149*. Abstract from 1st International Conference on Necrotic Enteritis in Poultry, Copenhagen, Denmark.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## **Inhibition studies of natural resin acids to *Clostridium perfringens* and *Escherichia coli* O149**

**Krisna Roy, Ulrike Lyhs and Karl Pedersen**

*National Veterinary Institute, Technical University of Denmark, Bülowsvej 27, 1870 Frederiksberg C, Copenhagen, Denmark*

### **Introduction**

As feed antibiotics are no longer allowed in Europe and some other countries, natural means to control the proliferation of pathogenic bacteria in the intestinal lumen of animals are needed. Natural resin acids are such natural products which have antimicrobial properties. In the present study, we aimed to evaluate the inhibitory activity of a resin-based product at different concentrations on intestinal bacterial pathogens.

### **Materials and methods**

The targeted product was Progres® (Suomen Rehu) containing 8% resin acids. *Clostridium perfringens* isolated from chickens, turkeys and pigs, respectively, and *Escherichia coli* O149 from pigs were tested. Growth of the pathogens was tested at 0.01%, 0.1% and 0.5% concentrations of the product. Inhibitory bioactivity of the product was examined via OD<sub>600</sub> measurements on growing cultures, by a 10-fold broth dilution method (DM), and by using an agar diffusion method (ADM). The OD method was followed only in one strain of *E. coli* O149. The DM was applied to three strains of each of the bacteria. Samples were taken after during incubation, where after 10-fold dilutions were made and plated onto blood agar plates. Counts were expressed as colony forming unit per ml (cfu/ml). The ADM was run on one strain of each of the bacteria, where zones of inhibition (mm) were measured. Subsequently, 10 *Cl. perfringens* strains, five from pigs, four from chickens and one from turkey, were tested with ADM against 0.5%, 1% and 5%.

### **Results**

OD measurements were difficult to interpret due to a considerable contribution of the test product to the turbidity. Therefore c.f.u. measurements were considered more accurate. In DM, no *Cl. perfringens* was found at any concentration of the product, indicating an efficient inhibition of *Cl. perfringens*. At 0.1% and 0.5% of the product, there was apparently lower cfu/ml of two strains of *E. coli* O149 compared to the corresponding controls, but *E. coli* was considerably less inhibited than *Cl. perfringens*. In ADM, zone of inhibition (ZI) was evolved around the product-concentration of 0.5% (ZI: 8 to 10 mm), 1% (8.5 to 12.0 mm), and 5% (9.0 to 19.5 mm) when performed on all ten strains of *Cl. perfringens*. No strain of *E. coli* O149 was inhibited by the product at any concentration in ADM, and not at 0.01% in DM.

### **Conclusion**

*Cl. perfringens* was inhibited even at low concentrations of the product containing resin acids, but there seemed to be some strain variation. *E. coli* O149 was only inhibited by high concentrations.