



## Electron-beam-induced dynamic charging of thin films

Beleggia, Marco; Migunov, V.; Dunin-Borkowski, R.E.

*Publication date:*  
2015

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*

Beleggia, M., Migunov, V., & Dunin-Borkowski, R. E. (2015). *Electron-beam-induced dynamic charging of thin films*. Poster session presented at 3rd Conference on Frontiers of Aberration Corrected Electron Microscopy, Kasteel Vaalsbroek, Netherlands.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# ELECTRON-BEAM-INDUCED DYNAMIC CHARGING OF THIN FILMS

Marco Beleggia<sup>§,†</sup>, Vadim Migunov<sup>‡</sup>, and Rafal E. Dunin-Borkowski<sup>‡</sup>

<sup>§</sup>Center for Electron Nanoscopy, Technical University of Denmark,  
DK-2800 Kgs. Lyngby, Denmark, mb@cen.dtu.dk

<sup>†</sup>Helmholtz-Zentrum Berlin für Materialien und Energie,  
Hahn-Meitner-Platz 1, D-14109 Berlin, Germany

<sup>‡</sup>Ernst Ruska-Centre for Microscopy and Spectroscopy with Electrons and  
Peter Grünberg Institute, Forschungszentrum Jülich, D-52425 Jülich, Germany

## Abstract

Primary electrons that are inelastically scattered generate electron-hole pairs in the illuminated area of a thin film, thereby enhancing its charge-carrier density and conductivity. The charge density gradient near the rim of the illumination patch triggers a drift-diffusion process that leads to a steady-state equilibrium with a local bias within a circular region that is larger than the beam diameter. The sign of the potential is dictated by the difference in mobility between electrons and holes and is generally positive. The magnitude of the local bias, which is also dependent on the mobilities, is determined primarily by the ratio between intrinsic and extrinsic carrier densities.

A proportion of the generated electrons that have sufficient energy may escape into vacuum, leaving holes behind and triggering larger-scale drift processes involving neutralizing currents that originate from electrical contacts to ground. This process also affects the steady-state equilibrium, both in the magnitude and in the extent of the local bias. The two effects cannot easily be kept separated.

In order to measure electron-beam-induced dynamical charging, we create a circular patch of illumination on a pristine region of a thin film by using the condenser system of the electron microscope. We then use the imaging lenses to record a time series of Fresnel defocus images of the illuminated area. As the currents start to flow and balance each other out, the out-of-focus appearance of the spot changes, reflecting the evolution of the local bias. We vary experimental parameters such as film composition, temperature, beam energy, beam current and beam current density and compile a comprehensive dataset that can be interpreted on the basis of the drift-diffusion equation. We are interested in characterizing the steady state and the previously observed asymmetry between charging and discharging characteristic times that goes by the name of the "Beriman effect", i.e. the charge footprint of the beam that appears to remain stable once the illumination is turned off.

Our study is relevant for understanding the operating principles of charge-based phase plates for electrons, such as the "hole-free" and "Volta" devices. More generally, the development of an understanding of electron-beam-induced charging processes and resulting local electric fields promises to be beneficial for a wide variety of *in situ* electron microscopy experiments involving gases, liquids, electrochemical processes (beam-assisted electrocatalysis), junctions, etc.