



## On the accuracy of HITEMP-2010 calculated emissivities of Water Vapor and Carbon Dioxide

Alberti, M.; Weber, R.; Mancini, M.; Fateev, A.; Clausen, S.

*Publication date:*  
2015

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*

Alberti, M., Weber, R., Mancini, M., Fateev, A., & Clausen, S. (2015). *On the accuracy of HITEMP-2010 calculated emissivities of Water Vapor and Carbon Dioxide*. Abstract from 12th International Conference on Energy for a Clean Environment, Lisboa, Portugal.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## On the accuracy of HITEMP-2010 calculated emissivities of Water Vapor and Carbon Dioxide\*

M. Alberti<sup>a†</sup>, R. Weber<sup>a</sup>, M. Mancini<sup>a</sup>, A. Fateev<sup>b</sup> and S. Clausen<sup>b</sup>

<sup>a</sup> *Institute for Energy Process Engineering and Fuel Technology, Clausthal University of Technology, Germany*

<sup>b</sup> *Department of Chemical and Biochemical Engineering, Technical University of Denmark, Denmark*

### Abstract

Nowadays, spectral Line-by-Line calculations using either HITRAN or HITEMP data bases are frequently used for calculating gas radiation properties like absorption coefficients or emissivities. Such calculations are computationally very expensive because of the vast number of spectral lines and, therefore, these calculations are usually taken as a benchmark to validate less complex models. In order to evaluate the accuracy of these simplified models, the accuracy of the Line-by-Line predictions using HITEMP data base must be known.

We present here some of our recent full spectrum measurements of spectral transmissivities which were done for H<sub>2</sub>O-CO<sub>2</sub>-N<sub>2</sub> mixtures for temperatures up to 1770K. Using these measured data it is possible to compare the Line-by-Line calculation using HITEMP-2010 on the basis of total (spectrally averaged) emissivity. At high pressures, also a proper lineshape treatment is essential for the accuracy of the Line-by-Line model. Spectral measurements that span the spectral range of at least one whole absorption band are hardly available in literature. Therefore, we compare the standard model with more advanced models which take the line-mixing effects into account. However, we show that also the standard model can be adjusted to yield reliable results.

*Keywords: Line-By-Line, HITEMP-2010*

---

\* See also M. Alberti, R. Weber, M. Mancini, A. Fateev, and S. Clausen, On the accuracy of HITEMP-2010 calculated emissivities of Water Vapor and Carbon Dioxide, 10<sup>th</sup> International Conference on Industrial Furnaces and Boilers (INFUB-10), Porto, 2015

†Corresponding author email: [alberti@ievb.tu-clausthal.de](mailto:alberti@ievb.tu-clausthal.de)