



Compositional Heterogeneity of the Upper mantle beneath the Siberian Craton: Reconciling Thermal, Seismic and Gravity Data

Artemieva, Irina; Herceg, Matija; Cherepanova, Yulia

Published in:
Goldschmidt2013: Conference Abstracts

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Artemieva, I., Herceg, M., & Cherepanova, Y. (2013). Compositional Heterogeneity of the Upper mantle beneath the Siberian Craton: Reconciling Thermal, Seismic and Gravity Data. In *Goldschmidt2013: Conference Abstracts* (pp. 620)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Please ensure that your abstract fits into one column on one page and complies with the *Instructions to Authors* available from the Abstract Submission web page.

Compositional heterogeneity of the upper mantle beneath the Siberian craton: reconciling thermal, seismic and gravity data

Irina Artemieva¹, Matija Herceg¹, Yulia Cherepanova¹, Hans Thybo¹

¹IGN, Univ. Copenhagen, Denmark, irina@geo.ku.dk

We present a new regional model for the upper mantle structure below the Siberian craton. The model includes (i) thermal model of the lithosphere structure constrained by surface heat flow data, (ii) seismic velocity heterogeneity constrained by global tomography models corrected for temperature variations in the upper mantle, (iii) and a new regional model for the density structure based on the GOCE satellite gravity data. We also present a new regional seismic crustal model SibCrust, which is used to calculate crustal correction to gravity field. Thermal model and seismic tomography models are used as independent constraints on lithosphere thickness, required to convert mantle residual density anomalies to density heterogeneity. New regional upper mantle model is compared with regional and world-wide petrological data on upper mantle velocities and densities constrained by mantle-derived xenoliths. The results indicate a significant compositional heterogeneity of the lithospheric mantle of the region. Seismic velocity and density anomalies cannot be explained by variations in $mg\#$ alone. Compared to the adjacent orogenic belts and the West Siberian basin, density structure of the cratonic mantle is in a close agreement with isopycnic condition, although significant regional variations are well correlated with basement topography.