



## MicroRNA regulation of TLRs in a post-influenza animal model

Brogaard, Louise; Heegaard, Peter M. H.; Larsen, Lars Erik; Dürrwald, R.; Schlegel, M.; Skovgaard, Kerstin

*Publication date:*  
2015

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Brogaard, L., Heegaard, P. M. H., Larsen, L. E., Dürrwald, R., Schlegel, M., & Skovgaard, K. (2015). *MicroRNA regulation of TLRs in a post-influenza animal model*. Abstract from TOLL2015 Targeting Innate Immunity, Marbella, Spain.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# MICRORNA REGULATION OF TLRs IN A POST-INFLUENZA ANIMAL MODEL

L. Brogaard<sup>1</sup>, P. M. H. Heegaard<sup>1</sup>, L. E. Larsen<sup>1</sup>, R. Dürrwald<sup>2</sup>, M. Schlegel<sup>2</sup>, K. Skovgaard<sup>1</sup>

<sup>1</sup>. National Veterinary Institute, Technical University of Denmark, Frederiksberg C, Denmark

<sup>2</sup>. IDT Biologika GmbH, Dessau-Rosslau, Germany

## Introduction

Substantial morbidity and mortality is caused by secondary bacterial infections occurring in individuals after influenza A virus (IAV) infection. Results from studies in mice suggest that this may in part be due to a lack of responsiveness to Toll-like receptor (TLR) ligands in the post-IAV infected individual. Using the pig as an animal model, we have identified microRNAs (miRNAs) that are differentially expressed in lung tissue two weeks after challenge compared to uninfected controls, i.e. well after the infection has cleared. The role for differential expression of miRNA at this late time point remains unclear. We therefore seek to examine the potential involvement of miRNAs in the translational regulation of TLRs and associated proteins, thus contributing to the lowered responsiveness to bacterial TLR ligands at this late time point, making the individual vulnerable to secondary infections.

## Methods and outcome

Pigs were experimentally challenged with a Danish reassortant IAV strain (A/sw/Denmark/12687/03(H1N2)). Lung tissue was harvested 14 days after challenge, as well as from uninfected control animals. Using RNAseq and high-throughput RT-qPCR, we quantified the expression of relevant miRNAs (e.g. miR-335 and miR-146a-5p) and mRNA levels of relevant miRNA targets.

Transcriptional analysis at the site of infection reveals a set of miRNAs to be regulated one week after the pigs had cleared the IAV infection (i.e. two weeks after challenge). This set included miRNAs experimentally validated or *in silico* predicted to bind to and regulate transcripts of TLRs and relevant co-factors and transcription factors (online tools). The antiviral immune response elicited by IAV infection thus includes late miRNA regulation, which in turn may be at the expense of host responsiveness to bacterial TLR ligands.