



Comparison of the Acidification Activities of Commercial Starter Cultures on Camel and Cow Milk

Berhe, Tesfemariam; Ipsen, Richard; Seifu, Eyassu; Kurtu, Mohammed Y.; Eshetu, Mitiku; Hansen, Egon Bech

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Berhe, T., Ipsen, R., Seifu, E., Kurtu, M. Y., Eshetu, M., & Hansen, E. B. (2015). *Comparison of the Acidification Activities of Commercial Starter Cultures on Camel and Cow Milk*. Poster session presented at 9th NIZO Dairy Conference, Papendal, Netherlands.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Comparison of the Acidification Activities of Commercial Starter Cultures on Camel and Cow Milk

Tesfemariam Berhe^{a*}, Richard Ipsen^b, Eyassu Seifu^c, Mohammed Y. Kurtu^a, Mitiku Eshetu^a, Egon Bech Hansen^d

^aSchool of Animal and Range Sciences, College of Agriculture, Haramaya University, Ethiopia

^bDairy, Meat and Plant Product Technology, Faculty of Science, University of Copenhagen, Denmark

^cDepartment of Food Science and Technology, Botswana College of Agriculture, Gaborone, Botswana

^dDivision of Food Microbiology, National Food Institute, Technical University of Denmark, Denmark

* Lucyselam@gmail.com



Introduction

- Haramaya Camel DairyTM is a project aiming at developing and improving dairy products from camel milk (*Camelus dromedarius*).
- The project focuses on properties, processing and product development of camel milk.
- Camel milk is composed of lactose, fat, and protein in roughly the same proportion as bovine milk. However, the relative composition of the proteins differs and β -lactoglobulin is absent in camel milk. (Hinz et al., 2012).
- The amino acid sequences of the camel caseins and whey proteins are homologous to the cow counterparts but also showing significant differences (Kappeler et al., 1998).
- Camel milk has been reported to be not easily fermentable because of its antibacterial and antiviral properties of the protective proteins (El Agamy et al., 1992)
- Through analysis of the chemical, physical and functional properties of the milk constituents we will be able to design and develop novel products from camel milk



Figure 1. Camels near Haramaya, Ethiopia

Materials and Methods

- Camel and cow milk samples were collected from Babile area and Haramaya University dairy farm in Ethiopia respectively.
- The milk samples were pasteurized at 65 °C for 30 minutes and inoculated with 0.1U/L of the starter cultures and incubated at 30 °C, 37 °C, and 42 °C.
- The 8 cultures were lyophilized cultures from the range of Chr Hansen A/S. R-704 and R-707 are mesophilic homofermentative cultures composed of strains of *Lactococcus lactis*.
- RST-743 is a homofermentative culture composed of strains of *Lactococcus lactis* and *Streptococcus thermophilus*.
- STI-12 is a homofermentative culture composed of strains of *Streptococcus thermophilus*
- XPL-2 is an aromatic LD culture composed of strains of *Lactococcus lactis* subsp. *cremoris*, *Lactococcus lactis* subsp. *lactis*, *Lactococcus lactis* subsp. *lactis* biovar *diacetylactis*, *Leuconostoc* species, and *Streptococcus thermophilus*.

Materials and Methods ...

- CHN-22 is a mesophilic aromatic LD culture containing strains of *Lactococcus lactis* subsp. *cremoris*, *Leuconostoc pseudomesenteroides*, *Lactococcus lactis* subsp. *lactis* biovar *diacetylactis*, *Lactococcus lactis* subsp. *lactis*, *Leuconostoc mesenteroides*.
- YoFlexR Mild 1.0 and YF-L904 are thermophilic yoghurt cultures containing strains of *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus*.
- The acidification of the cultures were followed using an iCinac instrument (Alliance Instruments, Freppillon, France) that measures the pH, oxidation reduction potential and temperature of the culture simultaneously.

Results

Table 1. Comparison of acidification activities of commercial starter cultures on camel and cow milk

| Temp °C | Culture | V _{max} (Max Acidification Rate) upH/min | | |
|---------|-------------|---|----------|------------|
| | | Camel milk | Cow milk | 50:50 milk |
| 30 | R-704 | -0.004 | -0.0085 | |
| | R-707 | -0.0065 | -0.009 | -0.0095 |
| | RST-743 | -0.007 | -0.0075 | |
| | CHN-22 | -0.004 | -0.006 | |
| | XPL-2 | -0.005 | -0.0075 | |
| 37 | XPL-2 | -0.005 | -0.007 | |
| | RST-743 | -0.003 | -0.0055 | |
| 42 | XPL-2 | -0.007 | -0.0075 | |
| | YF Mild 1.0 | -0.0065 | -0.015 | -0.010 |
| | YF-L904 | -0.0085 | -0.015 | |
| | STI-12 | -0.010 | -0.015 | |




Figure 2. Acidification curves of R-707 at 30 °C in camel milk, cow milk, and mixed camel and cow.

— camel milk
— cow milk
— 50:50 milk

Results.....




Figure 3.

Acidification curves of Yoflex® mild 1.0 at 42 °C in camel milk, cow milk, and mixed camel and cow.

— camel milk
— 50:50 milk
— cow milk

Discussion and Conclusion

- Results in the table and graph indicate that all cultures were able to acidify camel milk and reached a final pH at a level similar to cow milk but the speed of acidification was generally lower in camel milk.
- The reduced activity could either be due to inhibitory substances in camel milk or due to reduced availability of nutrients.
- To distinguish between these possibilities we conducted mixing experiments where acidifications were conducted in a 50:50 mixture of cow and camel milk. The acidification in the mixed milk was almost identical to the acidification in cow milk.
- Based on this result we find the most likely explanation to be that the cultures have difficulties in satisfying their nutritional needs in pure camel milk. The proteolytic system of the LAB cultures might show reduced efficiency on camel milk proteins.

References

- El-Agamy EI, R. Ruppner, A. Ismail, C.P. Champagne, R. Assaf (1992). Antibacterial and antiviral activity of camel milk protective proteins. *J. Dairy Res.* 59, 169-175.
- Hinz, K., P. M. O'Connor, T. Huppertz, R. P. Ross and A. L. Kelly (2012). Comparison of the principal proteins in bovine, caprine, buffalo, equine and camel milk. *Journal of Dairy Research* 79, 185-191.
- Kappeler, s, z. farah and z.puhan (1998). Sequence analysis of *Camelus dromedarius* milk caseins. *Journal of Dairy Research* .65, 209-222.

Acknowledgements

Danish International Development Agency, Danida for funding the project