



Differential expression of small RNAs under chemical stress and fed-batch fermentation in *Escherichia coli*

Rau, Martin Holm; Bojanovic, Klara; Nielsen, Alex Toftgaard; Long, Katherine S.

Published in:

Book of Abstracts. DTU's Sustain Conference 2015

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Rau, M. H., Bojanovic, K., Nielsen, A. T., & Long, K. S. (2015). Differential expression of small RNAs under chemical stress and fed-batch fermentation in *Escherichia coli*. In *Book of Abstracts. DTU's Sustain Conference 2015* [B-8] Technical University of Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Differential expression of small RNAs under chemical stress and fed-batch fermentation in *Escherichia coli*

Martin Holm Rau¹, Klara Bojanovič¹, Alex Toftgaard Nielsen¹, Katherine S. Long^{*1}

1: DTU Biosustain

*Corresponding author email: kalon@biosustain.dtu.dk

Abstract

Background: Bacterial small RNAs (sRNAs) are recognized as posttranscriptional regulators involved in the control of bacterial lifestyle and adaptation to stressful conditions. Although chemical stress due to the toxicity of precursor and product compounds is frequently encountered in microbial bioprocessing applications, the involvement of sRNAs in this process is not well understood. We have used RNA sequencing to map sRNA expression in *E. coli* under chemical stress and high cell density fermentation conditions with the aim of identifying sRNAs involved in the transcriptional response and those with potential roles in stress tolerance.

Results: RNA sequencing libraries were prepared from RNA isolated from *E. coli* K-12 MG1655 cells grown under high cell density fermentation conditions or subjected to chemical stress with twelve compounds including four organic solvent-like compounds, four organic acids, two amino acids, geraniol and decanoic acid. We have discovered 253 novel intergenic transcripts with this approach, adding to the roughly 200 intergenic sRNAs previously reported in *E. coli*. There are eighty-four differentially expressed sRNAs during fermentation, of which the majority are novel, supporting possible regulatory roles for these transcripts in adaptation during different fermentation stages. There are a total of 139 differentially expressed sRNAs under chemical stress conditions, where twenty-nine exhibit significant expression changes in multiple tested conditions, suggesting that they may be involved in a more general chemical stress response. Among those with known functions are sRNAs involved in regulation of outer membrane proteins, iron availability, maintaining envelope homeostasis, as well as sRNAs incorporated into complex networks controlling motility and biofilm formation.

Conclusions: This study has used deep sequencing to reveal a wealth of hitherto undescribed sRNAs in *E. coli* and provides an atlas of sRNA expression during seventeen different growth and stress conditions. Although the number of novel sRNAs with regulatory functions is unknown, several exhibit specific expression patterns during high cell density fermentation and are differentially expressed in the presence of multiple chemicals, suggesting they may play regulatory roles during these stress conditions. These novel sRNAs, together with specific known sRNAs, are candidates for improving stress tolerance and our understanding of the *E. coli* regulatory network during fed-batch fermentation.