



## Photocatalytic Water Splitting: Using a 2Photon Tandem Approach

Seger, Brian; Mei, Bastian Timo; Bae, Dowon; Kemppainen, Erno ; Pedersen, Thomas; Vesborg, Peter; Hansen, Ole; Chorkendorff, Ib

*Published in:*  
Book of Abstracts. DTU's Sustain Conference 2015

*Publication date:*  
2015

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Seger, B., Mei, B. T., Bae, D., Kemppainen, E., Pedersen, T., Vesborg, P., Hansen, O., & Chorkendorff, I. (2015). Photocatalytic Water Splitting: Using a 2Photon Tandem Approach. In *Book of Abstracts. DTU's Sustain Conference 2015* [E-1] Technical University of Denmark.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Photocatalytic Water Splitting: Using a 2-Photon Tandem Approach

Brian Seger<sup>\*1</sup>, Bastian Timo Mei<sup>1</sup>, Dowon Bae<sup>1</sup>, Erno Kempainen, Thomas Pedersen<sup>2</sup>, Peter Vesborg<sup>1</sup>, Ole Hansen<sup>1</sup>, Ib Chorkendorff<sup>\*1</sup>


1: DTU Physics; 2: DTU Nanotech 3: Aalto University

\*Corresponding author email: bseger@dtu.dk

Currently the world uses 17 TW of energy, with ~85% of that coming from fossil fuels. The solar irradiation that strikes earth is 120,000 TW, thus we only need to capture a fraction of this. However the greatest issue with solar energy from a practical and economic standpoint is its intermittency. (The sun doesn't shine at night.) This means storing solar based energy is just as important as harvesting it. As biological processes have proven, a very efficient way of storing energy is in the form of chemical bonds. The electrolysis of water into hydrogen and oxygen is an excellent way to store energy since water is a plentiful resource, hydrogen is an extremely energy dense molecule (by weight), and the oxygen by-product can be discarded to the atmosphere.

Rather than use a solar cell and an electrolyzer separately, we can combine these devices thus minimizing capital costs. This talk will focus on how we construct a photoelectrolyzer that is both highly efficient and highly durable in the corrosive environments needed for water splitting. Our focus is to use 2-photon tandem solar cells with transparent, conductive protective coatings to allow for highly efficient photoelectrolyzers. Figure 1A and 1B shows the general design of our approach to photoelectrolysis. While there are many parts to the device we are focusing on all of these in our quest to make a 20% efficient solar to hydrogen device. This talk will discuss what materials we use for both the large band gap and small band gap photoabsorber as well as our recent improvements in catalytic efficiency for both the H<sub>2</sub> evolution and O<sub>2</sub> evolution reaction.

1 A)



1 B)

