



## Microbial dynamics in anaerobic digestion (AD) reactors with sodium oleate in the substrate

Zhu, Xinyu; De Francisci, Davide; Treu, Laura; Kougias, Panagiotis; Campanaro, S.; Angelidaki, Irini

*Published in:*

Book of Abstracts. DTU's Sustain Conference 2015

*Publication date:*

2015

*Document Version*

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Zhu, X., De Francisci, D., Treu, L., Kougias, P., Campanaro, S., & Angelidaki, I. (2015). Microbial dynamics in anaerobic digestion (AD) reactors with sodium oleate in the substrate. In *Book of Abstracts. DTU's Sustain Conference 2015* Article R-7 Technical University of Denmark.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Microbial dynamics in anaerobic digestion (AD) reactors with sodium oleate in the substrate

X. Zhu<sup>1</sup>, D. De Francisci<sup>1</sup>, L. Treu<sup>1</sup>, P.G. Kougias<sup>1</sup>, S. Campanaro<sup>2</sup> and I. Angelidaki\*<sup>1</sup>

1: DTU Environment 2: Department of Biology, University of Padua

\*Corresponding author email: iria@env.dtu.dk

In AD systems, the accumulation of long chain fatty acids (LCFA) leads to process instability and decrease of the methane production. The understanding of how the microbial communities change in response to LCFA pulses is essential to optimize the overall process. In this study two lab-scale continuously stirred reactors used to characterize, via 16s rRNA gene analysis, the microbial shifts due to LCFA increase in the feedstock composition. Genera that had relative abundance higher than 1% or increased more than 2 folds after sodium oleate addition are chosen for discussion. The result shows that the average methane yield was  $150 \pm 4$  ml CH<sub>4</sub>/gVS for cattle manure and  $380 \pm 37$  ml CH<sub>4</sub>/gVS for manure with sodium oleate addition. 75% of the added sodium oleate was utilized to produce methane. The addition of sodium oleate caused a slight decrease on the methane yield with a concomitant increase of VFA. In average, 1.6 million sequencing reads were generated from each sample and 20% of the reads were identified using RDP classifier. Results showed that 4 genera had relative abundance higher than 1% after the LCFA addition, suggesting that these members involved in the LCFA degradation process. The relative abundance of *Pseudomonas*, *Clostridium XI* and *Clostridium III* was not significantly affected (changed less than 2 folds) by the addition of LCFA, suggesting that these genera are not directly involved in the  $\beta$ -oxidation process but probably in later steps of the degradation process, i.e. the metabolism of VFA. This observation consists with the previous study where *Pseudomonas* has been reported to have versatility in catabolizing esoteric organic compounds (von Graevenitz 1976), and *Clostridium III* has been found playing important role in the efficient operation of anaerobic reactors (Shiratori, Ikeno et al. 2006). *Syntrophomonas* increased significantly after the addition of sodium oleate and became the most abundant genus. Several members of *Syntrophomonas* genus have been proven to be specialized in performing  $\beta$ -oxidation on LCFA (Sousa, Smidt et al. 2009). The significant increase of *Syntrophomonas* found in this experiment confirmed that this genus is a key player in the LCFA degradation under thermophilic anaerobic conditions. As for methanogens, *Methanoculleus* was found to be the most abundant genus, with no significant change before and after sodium oleate addition.

### Reference:

- Shiratori, H., H. Ikeno, S. Ayame, N. Kataoka, A. Miya, K. Hosono, T. Beppu and K. Ueda (2006). "Isolation and characterization of a new *Clostridium* sp. that performs effective cellulosic waste digestion in a thermophilic methanogenic bioreactor." *Applied and environmental microbiology* 72(5): 3702-3709.
- von Graevenitz, A. (1976). "Genetics and Biochemistry of *Pseudomonas*." *The Yale journal of biology and medicine* 49(3): 308
- Sousa, D. Z., H. Smidt, M. M. Alves and A. J. Stams (2009). "Ecophysiology of syntrophic communities that degrade saturated and unsaturated long - chain fatty acids." *FEMS microbiology ecology* 68(3): 257-272.
- Hori, T., S. Haruta, Y. Ueno, M. Ishii and Y. Igarashi (2006). "Dynamic transition of a methanogenic population in response to the concentration of volatile fatty acids in a thermophilic anaerobic digester." *Applied and environmental microbiology* 72(2): 1623-1630