



Utilization of biomass combustion ash for liming purposes on top of forest soils

Maresca, Alberto; Astrup, Thomas Fruergaard

Published in:
Book of Abstracts. DTU's Sustain Conference 2015

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Maresca, A., & Astrup, T. F. (2015). Utilization of biomass combustion ash for liming purposes on top of forest soils. In *Book of Abstracts. DTU's Sustain Conference 2015* [W-12] Technical University of Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Utilization of biomass combustion ash for liming purposes on top of forest soils

A. Maresca ^{*1}, T.F. Astrup ¹

1: DTU Environment; *Corresponding author email: almar@env.dtu.dk

The use of energy from wood combustion is becoming an increasingly popular source of renewable energy. Ashes are generated and although alternative uses – e.g. for liming and fertilizing purposes - have been proposed (e.g. Steenari et al., 1999), biomass ashes are commonly landfilled (Knapp & Insam, 2011). Benefits and risks owing to ash utilization have to be evaluated based on the ash chemical composition but also on the actual leaching release. Although several studies have focused on the ash composition and potential leaching release (e.g. Mellbo et al., 2008; Steenari et al., 1998; Zhan et al., 1996), the results are still fragmented.

In this study, two biomass combustion ash samples were investigated. The samples were characterized for their chemical composition, thus acid digested and analyzed by ICP-MS/OES. The long term leaching of the ash samples was investigated according to CEN/TS 14405:2004. Overall, higher contents of some plant nutrients – i.e. Ca, K, Mg, Mn and P - as well as of a few heavy metals - i.e. Cd and Ni - were found in the ash samples compared to Danish forest soil (Ingerslev et al., 2014). A long term leaching of alkaline compounds was found and although a decrease in the pH value – initial pH of 12.5 - was detected along the experiment, a pH value above 10.5 was registered at the L/S ratio 1000 l/kg. High releases of soluble compounds, such as K, Na and S were found, and drops of about two orders of magnitudes were shown by the L/S ratio 20 l/kg. The concentrations of P, Cd and Ni were always below detection limit: 15 µg/l, 0.20 µg/l and 0.52 µg/l respectively. Although similar patterns in the release of Cr were found by comparing the two ashes, the higher Cr releases were found for the fly ash sample.

The suitability of wood ash for liming purposes was confirmed. Although high contents of nutrients were found in our ash samples in comparison with the investigated Danish forest soil, their different solubility (e.g. K and P) limits the assessment of the actual fertilizing value of the ash. Considerable amounts of some soluble nutrients are likely to leach out after the first rain event, while others showed relatively small releases according to our results. Although generally low releases of heavy metals were found, their toxicity in relation to the actual soil organisms should be investigated to assess potential risks owing to ash utilization.

REFERENCES

- CEN/TS 14405, 2004. Characterization of waste – Leaching behaviour tests – Up-flow percolation test.
- Ingerslev, M., Hansen, M., Pedersen, L.B., Skov, S., 2014. Effects of wood chip ash fertilization on soil chemistry in a Norway spruce plantation on a nutrient-poor soil. *For. Ecol. Manage.* 334, 10–17.
- Knapp, B.A., Insam, H., 2011. Recycling of Biomass Ashes: Current Technologies and Future Research Needs, in: Insam, H., Knapp, B.A. (Eds.), *Recycling of Biomass Ashes*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 1–16.
- Mellbo, P., Sarenbo, S., Stålnacke, O., Claesson, T., 2008. Leaching of wood ash products aimed for spreading in forest floors - Influence of method and L/S ratio. *Waste Manag.* 28, 2235–44.
- Steenari, B.-M., Karlsson, L.G., Lindqvist, O., 1999. Evaluation of the leaching characteristics of wood ash and the influence of ash agglomeration. *Biomass and Bioenergy* 16, 119–136.
- Steenari, B.-M., Marsic, N., Karlsson, L.-G., Tomsic, A., Lindqvist, O., 1998. Long term leaching of stabilized wood ash. *Doktorsavhandlingar Vid Chalmers Tek. Hogsk.*
- Zhan, G., Erich, M.S., Ohno, T., 1996. Release of trace elements from wood ash by nitric acid. *Water, Air, Soil Pollut.* 88, 297–311.