



Microbial interference with hatch and survival of European eel larvae

Sørensen, Sune Riis; Lauesen; Tomkiewicz, Jonna; de Schryver, P.

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Sørensen, S. R., Lauesen, Tomkiewicz, J., & de Schryver, P. (2013). *Microbial interference with hatch and survival of European eel larvae*. Abstract from DAFINET Workshop : Diagnosis and Control of Fish Diseases, Frederiksberg, Denmark.

http://www.dafinet.dk/DAFINET/Abstract_books_files/DAFINET%20April%202013%20Abstracts.pdf

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Microbial interference with hatch and survival of European eel larvae

Sørensen S.R.¹, Lauesen², Tomkiewicz J.¹ and de Schryver P.³

¹*National Institute of Aquatic Resources,
Technical University of Denmark, Denmark*

²*Billund Aquaculture Service Aps., Kløvermarken 27, Billund, 7190, Denmark*

³*Laboratory of Aquaculture & Artemia Reference Center,
Ghent University, Rozier 44, 9000 Gent, Belgium*

Recent research has significantly improved our knowledge and capabilities in the field of *in vitro* production of yolk sac larvae from European eel (*Anguilla anguilla*). Female broodstock European eels are matured by weekly administration of pituitary extract and male eels with hCG (human chorionic gonadotropin), which afford gametes for *in vitro* fertilization studies. The maturing process may lead to mass hatchings of up to ½ million larvae of which some survive the entire yolk sac phase. However, the rearing of larvae suffers from high larval mortalities, and water quality might be a crucial factor for larval survival in rearing systems.

By applying antibiotic treatment as a research tool, it was possible to determine the extent of microbial interference in the production of high numbers of good quality larvae. By controlling microbiota during egg and larval incubation, the egg hatching success and larval longevity more than doubled. Using scanning electron microscopic analysis it was observed that microbe inhibiting treatments reduced bacterial colonization of the eggs surface, which possibly cause reduced gas and ionic exchange across chorionic membrane.

These results suggest that future eel larviculture should not only focus on optimizing physical incubation conditions, but certainly also on the control over microbial interference.

Presenting author: Sune Riis Sørensen; srs@aqu.dtu.dk