



## Identification of the sansalvamide non-ribosomal peptide synthetase in *Fusarium solani*

Sørensen, Jens Laurids; Fuertes, Patricia Romans ; Sondergaard, Teis Esben; Nielsen, Kristian Fog; Hansen, Frederik Teilfeldt; Giese, Henriette; Brodersen, Ditlev Egeskov

*Published in:*

Book of abstracts from the 13th European Conference on Fungal Genetics

*Publication date:*

2016

*Document Version*

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Sørensen, J. L., Fuertes, P. R., Sondergaard, T. E., Nielsen, K. F., Hansen, F. T., Giese, H., & Brodersen, D. E. (2016). Identification of the sansalvamide non-ribosomal peptide synthetase in *Fusarium solani*. In *Book of abstracts from the 13th European Conference on Fungal Genetics* (pp. 437-437). [CS5T51]

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

POSTER SESSION ABSTRACTS  
Session CS5 Applied genomics and biotechnology  
CS5T51

**Tuesday 5th April**  
**14:00 - 16:00**

**SØRENSEN Jens Laurids** (1), **ROMANS-FUERTE**s Patricia (2), **SONDERGAARD** Teis Esben (3), **NIELSEN** Kristian Fog (4), **HANSEN** Frederik T (1), **GIESE** Henriette (3), **BRODERSEN** Ditlev Egeskov (2)

(1) Aalborg University, Esbjerg, Denmark

(2) Aarhus University, Aarhus, Denmark

(3) Aalborg University, Aalborg, Denmark

(4) Technical University of Denmark, Kgs. Lyngby, Denmark

**Identification of the sansalvamide non-ribosomal peptide synthetase in *Fusarium solani***

Members of the *Fusarium* genus have a huge genetic potential for production of secondary metabolites. One of the most interesting compound classes is the non-ribosomal peptides (NRPs), which are synthesized by huge multi-domain synthetases (NRPSs). In a comparative analysis of the genomes sequences from ten different *Fusarium* species we have previously identified 52 NRPS orthology groups of which only 6 produce a known compound [1]. To fill the missing pieces we set out to identify the biosynthetic pathway responsible for production of the NRP sansalvamide. This cyclic pentadepsipeptide was originally isolated from an unidentified *Fusarium* species [2] and subsequently several strains belonging to the *Fusarium solani* species complex [3]. Sansalvamide contains an A-hydroxyisocaproic acid (HICA) unit, which is also found in the cyclic hexadepsipeptide destruxin produced by *Metarhizium* species. The gene cluster responsible for destruxin biosynthesis has been identified in *M. robertsii*, which consists of non-ribosomal peptide synthetase (NRPS; DtxS1), an aldo-keto reductase (DtxS2), a cytochrome P450 monooxygenase and a decarboxylase (DtxS4) [4]. A BlastP analysis of the synthetase DtxS1 against *F. solani* sequences resulted in NRPS30 as the best hit (total score 18001; identity: 45%). An orthologue of DtxS3, which provides the HICA unit from reduction of A-ketoisocaproic acid, was furthermore identified directly downstream of NRPS30. To verify that NRPS30 is responsible for biosynthesis of sansalvamide in *F. solani* we applied an *Agrobacterium tumefaciens*-mediated transformation (ATMT) approach to generate knock-out mutants. Comparative studies of secondary metabolites in the resulting deletion mutants and wild type confirmed the absence of sansalvamide in the NRPS30 deletion mutant, implicating this synthetase in the biosynthetic pathway for sansalvamide.

[1] Hansen, F. T., Gardiner, D. M., Lysøe, E., Fuertes, P. R., Tudzynski, B., Wiemann, P., Sondergaard, T. E., Giese, H., Brodersen, D. E. and Sørensen, J. L. Fungal Genet. Biol. 2015 75, 20-29.

[2] Belofsky, G. N., Jensen, P. R. and Fenical, W. Tetrahedron Lett. 1999 40, 2913-2916.

[3] Short, D. P. G., O'Donnell, K., Thrane, U., Nielsen, K. F., Zhang, N., Juba, J. H. and Geiser, D. M. Fungal Genet Biol 2013 53, 59-70.

[4] Donzelli, B. G. G., Krasnoff, S. B., Sun-Moon, Y., Churchill, A. C. L. and Gibson, D. M. Curr. Genet. 2012 58, 105-116.