



Perturbation of Neonatal Microbial Gut Community by Peripartum Antibiotics in Wistar Rats Lead to Decreased Weight Gain

Tulstrup, Monica Vera-Lise; Roager, Henrik Munch; Clement Thaarup, Ida; Licht, Tine Rask; Bahl, Martin Iain

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Tulstrup, M. V.-L., Roager, H. M., Clement Thaarup, I., Licht, T. R., & Bahl, M. I. (2016). *Perturbation of Neonatal Microbial Gut Community by Peripartum Antibiotics in Wistar Rats Lead to Decreased Weight Gain*. Poster session presented at 10th INRA-Rowett Symposium: Gut Microbiology, Clermont-Ferrand, France.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.




Perturbation of Neonatal Microbial Gut Community by Peripartum Antibiotics in Wistar Rats Leads to Decreased Weight Gain

Monica Vera-Lise Tulstrup¹, Henrik Munch Roager¹, Ida Clement Thaarup¹, Tine Rask Licht¹ and Martin Iain Bahl¹

Introduction

Antibiotics are frequently administered orally to either mothers or young children to treat or prevent bacterial infections not necessarily related to the gastrointestinal system. This has adverse effects on the commensal gut microbial community, as it disrupts the intricate balance between specific bacterial groups within this ecosystem, potentially leading to dysbiosis.

We hypothesize that modulation of community composition and function induced by peripartum antibiotics affects intestinal microbial composition and general health of the offspring.




Methods

Pregnant Wistar rats (n=33) were dosed by oral gavage with either amoxicillin (AMX), vancomycin (VAN), or water (CON) daily from 8 days before delivery until weaning of the offspring. Offspring weight-gain was recorded during the entire study period and dissections were performed at four time points (2 days, 2 weeks, 4 weeks and 14 weeks). Bacterial abundance was determined by plating of fresh fecal samples from dams right before birth. Bile acids levels were determined in the blood serum by UPLC-MS.

Results

1. Bacterial load in antibiotic treated dams, shortly before giving birth, is significantly higher than in control animals.

Figure 1: Bacterial abundance in feces of dams shortly before birth of offspring. (A) Abundance of anaerobic bacteria (Wilkins-Chalgren agar) and (B) aerobic Bacteria (Plate-count agar). Each dot represents an individual animal. Horizontal lines and error bars show means and SEM, respectively. Significant differences from CON group are indicated by asterisks (***) (P < 0.001).



2. Significant differences in liver, spleen, cecum, cecum pH and epididymal fat between Groups.




Figure 2: Significant changes in body composition. (A) Liver percentage of bodyweight in 4 week old offspring, (B) Spleen percentage of bodyweight in 14 week old offspring, (C) Cecum weight of 14 week old offspring, (D) Epididymal fat weight of 14 week old offspring, (E) Spleen weight of dams, (F) Spleen percentage of bodyweight in dams, (G) Cecum weight of dams, and (H) Cecum pH in Cecum. Each point represents an individual animal. Horizontal lines and error bars show means and SEM, respectively. Significant differences from CON group are indicated by asterisks (*; P < 0.05, **; P < 0.01, ***; P < 0.001).

3. Significantly lower weight gain and food intake was observed in offspring from antibiotic treated dams compared to controls.




Figure 3: Weight gain and food intake in offspring. (A) Average Growth of Offspring, (B) Average weight gain difference of offspring of antibiotic treated compared to control animals and (C) Average food intake in Offspring after weaning. Each point represents a group average. Horizontal lines and error bars show means and SEM, respectively. Significant differences from CON group are indicated by asterisks (*; P < 0.05).

4. Antibiotic treatment causes significant changes of the bile acid (BA) concentration and composition in the blood serum of dams.




Figure 5: Sum of Primary, Conjugated Primary, Secondary and Conjugated Secondary Bile Acids. (A) Primary bile acids, (B) Conjugated primary bile acids, (C) Secondary bile acids, and (D) Conjugated secondary bile acids. Horizontal lines and error bars show means and SEM, respectively. Significant differences from CON group are indicated by asterisks (*; P < 0.05, **; P < 0.01, ***; P < 0.001).

Conclusion

We observed reduced weight-gain in offspring of antibiotic treated dams compared to controls, even though offspring never received antibiotics directly. Additionally peripartum antibiotic treatment of rats affect liver, spleen, cecum and epididymal fat pad weight at least 14 weeks after birth. The antibiotic treated dams show a significant increase in both aerobic and anaerobic bacteria, which constitute the inoculum for the offspring. Bile acid profiles are also changed significantly in the dams, indicating a markedly different bacterial composition and activity in the gastrointestinal tract that may explain the weight difference in offspring.

Future work

- 16S rRNA gene sequencing:** Bacterial community composition in feces and caecum
- UPLC-MS of bile acids:** Analyze bile acid composition in blood (and feces) of offspring
- BOMB calorimeter:** Analyze energy content in feces
- GC-MS of SCFA:** Analysis of short chain fatty acids in caecum
- qPCR of gene expression:** Examining effects of AB on satiety, tight junction proteins, Bile acid reg., immune system reg.
- qPCR/ plating:** Bacterial abundance in offspring
- Liver fat:** Analyze percentage of fat in liver
- Behavioral study:** Analyze social and locomotive behavior in offspring
- Elisa of gut hormones, blood lipids and inflammation factors:** PYY, GLP-1, Haptoglobin, Leptin