



Development and validation of extensive growth and growth boundary models for psychrotolerant pseudomonads in seafood, meat and vegetable products

Martinez Rios, Veronica; Dalgaard, Paw

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Martinez Rios, V., & Dalgaard, P. (2016). *Development and validation of extensive growth and growth boundary models for psychrotolerant pseudomonads in seafood, meat and vegetable products*. Abstract from 25th International ICFMH Symposium, Dublin, Ireland.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Food Micro 2016

Food Spoilage

FOODMICRO2016-267

DEVELOPMENT AND VALIDATION OF EXTENSIVE GROWTH AND GROWTH BOUNDARY MODELS FOR PSYCHROTOLERANT PSEUDOMONADS IN SEAFOOD, MEAT AND VEGETABLE PRODUCTS

Veronica Martinez-Rios¹, Paw Dalgaard*¹

¹Microbiology and Production, National Food Institute, DTU Food, Kgs. Lyngby, Denmark

Developing Scientist Competition: No

Presentation Preference: Oral

Abstract Content: Extensive growth and growth boundary models were developed and validated for psychrotolerant pseudomonads growing in seafood, meat and vegetable products. The new models were developed by expanding an existing cardinal parameter-type model for growth of pseudomonads in milk (Martinez-Rios *et al.*, Int. J. Food Microbiol. 216. 110-120, 2016). MIC-values for acetic-, benzoic- and citric acids were determined in broth and terms modelling their antimicrobial effect were added to the model. Cardinal parameter values for CO₂ and a_w were obtained from literature. The new model included 9 environmental parameters and their interactive effects. It was successfully validated using 319 growth rates (μ_{max} -values) for psychrotolerant pseudomonads in seafood and meat products. These data from literature (n=291) or own experiments (n=28) resulted in bias and accuracy factor values of 1.14 and 1.28, respectively, when observed and predicted μ_{max} -values were compared. Thus, on average μ_{max} -values for seafood and meat products were overestimated by 14%. Additionally, the reference growth rate parameter $\mu_{ref25^{\circ}C}$ was calibrated by fitting the model to 21 μ_{max} -values in vegetable products. This resulted in a $\mu_{ref25^{\circ}C}$ -value of 0.54 1/h. The calibrated vegetable model was successfully validated using 51 μ_{max} -values for psychrotolerant pseudomonads in vegetables. Average bias and accuracy factor values of 1.24 and 1.38 were obtained, respectively. Lag time models were developed by using relative lag times from literature data. Performance of the new expanded model was equally good for seafood and meat products, and importance of including the effect of acetic, benzoic, citric acids and CO₂ in order to accurately predict growth of psychrotolerant pseudomonads was clearly demonstrated e.g. for brined shrimps. The high number of environmental parameters included in the two models make them flexible and suitable for product development as the effect of substituting one combination of preservatives with another can be predicted.

Disclosure of Interest: None Declared

Keywords: Chilled food chain, Model validation, Product development, Shelf-life prediction, Spoilage