



A detailed view of *Listeria monocytogenes*' adaptation and survival under cold temperature stress

Hingston, P.; Hansen, Lisbeth Truelstrup; Wang, S. ; Allen, K. ; Chen , J.

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Hingston, P., Hansen, L. T., Wang, S., Allen, K., & Chen, J. (2016). *A detailed view of Listeria monocytogenes' adaptation and survival under cold temperature stress*. Abstract from FoodMicro 2016 – 25th International ICFMH Symposium, Dublin, Ireland.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Poster Abstracts

Food Microbiology & Human Health

PO2-SB-193

A DETAILED VIEW OF LISTERIA MONOCYTOGENES' ADAPTATION AND SURVIVAL UNDER COLD TEMPERATURE STRESS

P. Hingston^{1,*}, L. Truelstrup Hansen², S. Wang¹, K. Allen¹, J. Chen¹

¹Food, Nutrition and Health, University of British Columbia, Vancouver, Canada, ²Microbial Food Safety and Environmental Hygiene, Technical University of Denmark, Søborg, Denmark

Abstract Content: The human pathogen *Listeria monocytogenes* (*Lm*) continues to be a challenge for the food industry where it is known to contaminate ready-to-eat foods and grow during refrigerated storage. In order to gain increased control of *Lm* in the food-supply-chain, an improved understanding of low temperature stress adaptation methods is needed. In this study, RNA-seq (strand specific Illumina libraries; 22-39 million 2x100bp reads) and cell membrane fatty acid profiling were used to analyze adaptation mechanisms used by a fast growing, serotype 1/2a, *Lm* food plant isolate at 4°C. Brain heart infusion (BHI) broth pre-tempered to 20 or 4°C, was inoculated with 10⁷ CFU/ml of stationary phase (SP) cells grown at 20°C in BHI, and incubated at 20 and 4°C. RNA and lipids were extracted at T1– early lag phase (LP), T2 – end of LP, T3 – mid-exponential, T4 – transition to SP, and T5 – late SP (T5+4 h or 2 days for 20 and 4°C respectively). The number of coding transcripts upregulated (>2 log₂, p<0.05) at 4°C relative to 20°C was 142, 96, 91, 45, and 388 from T1-T5 respectively, while the number of downregulated genes at T1-T5 was 91, 38, 56, 125, and 256 respectively. Notably, the greatest differential gene expression occurred in *Lm* cells during late SP at 4°C, the most relevant physiological state to *Lm*'s survival in chilled food products. Common among all time points was the upregulation of nine genes required for branched-chain fatty acid (BCFA) synthesis, which was supported by an increase in membrane BCFAs from 77% at T1-4°C to 93% at T5-4°C. Putative cold stress regulatory mechanisms could be observed through negatively correlated expression levels of sense and antisense RNA. This research highlights *Lm*'s response to cold stress and provides deeper insight into how refrigerated storage conditions influence microbial gene expression and physiology.

Disclosure of Interest: None Declared

Keywords: cold stress, Food Safety, *Listeria monocytogenes*, RNA sequencing, time course