



## **Seaweed potentials – evaluation of year-round biomass composition of commercial cultivated sugarkelp- results from project KOMBI**

**Holdt, Susan Løvstad; Silva Marinho, Goncalo; Angelidaki, Irini**

*Publication date:*  
2015

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*

Holdt, S. L., Silva Marinho, G., & Angelidaki, I. (2015). *Seaweed potentials – evaluation of year-round biomass composition of commercial cultivated sugarkelp- results from project KOMBI*. Abstract from 5th Nordic Seaweed Conference, Grenaa, Denmark.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Nordic Seaweed Conference 2015, Oct 7-8**

Kystvejen's Hotel and Conference Center, Kystvej 26, 8500 Grenaa, Denmark

**Seaweed potentials – evaluation of year-round biomass composition of commercial cultivated sugarkelp- results from project KOMBI: Susan L. Holdt, Goncalo S. Marinho, Iriini Angelidaki**

[suho@food.dtu.dk](mailto:suho@food.dtu.dk) Affiliations: Technical University of Denmark, Kgs. Lyngby, Denmark

**ABSTRACT**

In this study, the year-round protein, amino acid, fatty acid, pigments, mineral and vitamin content and profiles were considered to evaluate the nutritional value and harvest time of the *Saccharina latissima* biomass for optimized value and application.

Sugarkelp was cultivated both in close proximity to a blue mussel and fish farm (IMTA) and in a reference site, both outside Horsens fjord in Denmark. Sugarkelp biomass was measured by harvesting sporophytes (deployed in February 2013) from 1m rope droppers (n=3) at 2 m depth in 2013-2014. Biomass was weighed, followed by freeze drying, homogenizing and chemical characterization by various methods for the specific analyses of biomass composition.

Protein content varied throughout the experimental period with the highest values recorded in November (10.8%) and the lowest values recorded in May 2013 (1.3 %). The lipid concentration varied from 0.62-0.88% DW in July to 3.33-3.35% DW in November ( $P<0.05$ ). Polyunsaturated fatty acids (PUFA's) made up more than half of the fatty acids with a maximum in July (52.3-54.0% FAME), including the most appreciated health beneficial EPA and DHA. Mineral content are discussed in relation to legislations, and a few trace metals such as Cd, Pb, Iodine may be problematic in some seasons. The pigment profile did not change during the year, however the concentration did, and with fucoxanthin as the most interesting.

Generally the year-round variations were due season, and no difference between the two locations (reference and IMTA).