



## Combining X-ray and Electron Based in situ Characterization of Catalysts

Damsgaard, Christian Danvad

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*

Damsgaard, C. D. (2016). *Combining X-ray and Electron Based in situ Characterization of Catalysts*. Abstract from 3rd International Conference on In-Situ and Correlative Electron Microscopy, Saarbrücken, Germany.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Combining X-ray and Electron Based *in situ* Characterization of Catalysts

Christian D. Damsgaard<sup>1,2</sup>

<sup>1</sup> Department of Physics, Technical University of Denmark, Kgs. Lyngby, Denmark.

<sup>2</sup> Center for Electron Nanoscopy, Technical University of Denmark, Kgs. Lyngby, Denmark

To understand the function of catalysts for heterogeneous catalysis and improve catalytic properties, a fundamental insight into structure-functionality relationships is required. As catalysts may change their structure with respect to the environment, it is essential to investigate the catalysts under reaction conditions. Furthermore, structural and compositional information have to be acquired on different length scales [1] and such *in situ* studies require dedicated complementary techniques. *In situ* X-ray diffraction (XRD) and X-ray absorption spectroscopy (XAS) are eminent to follow the average catalyst's structural (local and global) and electronic changes during synthesis and reaction conditions [2]. In order to follow the dynamics (stability, topography, etc.) of the surface and interfaces, imaging is often the solution [3]. Nanoscale imaging and spectroscopy of catalysts in a gaseous environment is traditionally performed in an environmental transmission electron microscope (ETEM). TEM gives insight in the atomic changes during reaction, however it is restricted to relatively low pressure (<~1 kPa) and a thin sample (<~100 nm) [4]. Spatially resolved information on the meso scale (50 nm–1  $\mu$ m) can be obtained by X-ray microscopy, which enables *in situ* studies at both ambient and elevated pressure [5]. Furthermore, due to the much higher penetration depth of X-rays compared to electrons, X-ray based characterization techniques are more suited for realistic sample and conditions.


This contribution illustrates how catalyst properties can be elucidated by the combination of X-ray and electron based methods in general. One example is the study of Ga-based (PdGa and NiGa) catalysts for methanol synthesis, where the average local structure and electronic structure obtained with XAS is combined with the average global structural information from XRD. Additionally, single nanoparticle (NP) compositional and structural information was studied by high-resolution ETEM experiments (Figure 1). The catalysts were investigated during formation and methanol synthesis and a full life-cycle model of the catalyst is formulated[6,7].

Another example highlighted in this contribution combines X-ray imaging with ETEM studies. By use of a special designed *in situ* cell (Figure 2A) a bifunctional Cu/ZnO@zeolite core-shell catalyst for direct production of methanol was studied with hard X-ray microscopy (X-ray ptychography). The cell is based on a TEM heater chip from Protochips (Figure 2B), which enables *ex situ* electron microscopy imaging before and after reaction [5]. The complementary nature of *in situ* hard X-ray ptychography and electron microscopy was applied to study the stability of the core-shell catalyst in a hierarchical manner at different length scales during reduction and oxidation treatments. The study reveals a stable core-shell interface at 250°C, although reduction of the Cu containing core material led to a shrinkage of the particles on the nanometer scale. At further heating to 350°C changes on the  $\mu$ m scale were observed.


Several material systems have been investigated and diffraction, imaging and spectroscopic methods have been applied both with x-rays and electrons. The results clearly underlined the need for complementary techniques and highlight the potential of these for application in catalysis.

References:

- [1] J. D. Grunwaldt, J. B. Wagner, and R. E. Dunin-Borkowski, *ChemCatChem* **5** (2013), p. 62.  
 [2] B. S. Clausen, *Catal. Today* **39** (1998), p. 293.  
 [3] Q. Jeangros, T. W. Hansen, J. B. Wagner, R. E. Dunin-Borkowski, C. Hébert, J. Van Herle, and A. Hessler-Wyser, *Chem. Commun. (Camb)*. **50** (2014), p. 1808.  
 [4] T. W. Hansen, J. B. Wagner, P. L. Hansen, S. Dahl, H. Topsøe, and C. J. H. Jacobsen, *Science* **294** (2001), p. 1508.  
 [5] S. Baier, C. D. Damsgaard, M. Scholz, F. Benzi, A. Rochet, R. Hoppe, T. Scherer, J. Shi, A. Wittstock, M. Burghammer, J. B. Wagner, C. G. Schroer, and J.-D. Grunwaldt, *Microsc. Microanal.* **22** (2016), p. 178.  
 [6] E. M. Fiordaliso, I. Sharafutdinov, H. W. P. Carvalho, J.-D. Grunwaldt, T. W. Hansen, I. Chorkendorff, J. B. Wagner, and C. Damsgaard, *ACS Catal.* **5** (2015), p. 5827.  
 [7] I. Sharafutdinov, C. F. Elkjær, H. W. Pereira de Carvalho, D. Gardini, G. L. Chiarello, C. D. Damsgaard, J. B. Wagner, J.-D. Grunwaldt, S. Dahl, and I. Chorkendorff, *J. Catal.* **320** (2014), p. 77.



**Figure 1.** (A) HRTEM of a NiGa NP supported on SiO<sub>2</sub>, acquired in 110Pa H<sub>2</sub> at 700°C. (B) STEM-EELS on NiGa NP acquired in vacuum at room temperature.



**Figure 2.** A) Sketch of in situ cell used for x-ray Ptychography. B) Optical micrograph of a Protochips® Aduro E-Chip with SiN membrane [5].