



## How can we make Fickian dispersion models useful in practice?

Harry Lee, Jonghyun; Rolle, Massimo; Kitanidis, Peter

*Publication date:*  
2016

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Harry Lee, J., Rolle, M., & Kitanidis, P. (2016). *How can we make Fickian dispersion models useful in practice?*. Abstract from AGU Fall meeting 2016, San Francisco, United States.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## H34D-04: How can we make Fickian dispersion models useful in practice?

Wednesday, 14 December 2016

16:45 - 17:00

📍 Moscone West - 3018

Dispersion in porous media originates from the variability of fluid velocity jointly with concentration at scales smaller than the ones resolved in the continuum description of solute transport. The unresolved scales are thus associated with the pore-grain geometry and the larger-scale heterogeneity that are ignored when the composite pore-grain medium is replaced by a homogeneous continuum. This applies whether the formation is modelled as homogeneous or discretized into homogeneous blocks. The process of dispersion is typically described through the Fickian model, *i.e.*, dispersive flux is proportional to the gradient of the resolved concentration, commonly with the Scheidegger parameterization, *i.e.*, a particular way to compute the dispersion coefficients utilizing dispersivity coefficients. Although the Fick-Scheidegger parameterization is by far the most commonly used in solute transport applications, its validity has been questioned.

In this presentation, we will list conditions under which the Fickian dispersion model is justified, then present how the practical Fickian model can approximate what is often thought as “non-Fickian” behavior. Specifically, we will show with several examples that the Fickian dispersion model performs adequately using appropriate dispersion coefficients in a domain that is discretized finely enough for the local equilibrium conditions that the Fickian model requires to be satisfied. Over the last thirty years, advances in numerical linear algebra, adaptive mesh refinement, and high performance computing environments, in combination with a dramatic drop in computational cost, have made it possible to perform fine-resolution simulation. We will also present how upscaled hydraulic conductivity and macrodispersion coefficients change with respect to different grid size and heterogeneity scale and discuss the role of diffusion and mean velocity. From these illustrations, we argue that the predictive ability of transport modeling can be greatly improved by finer-scale simulation because fine-scale modeling captures the physics of dispersion more accurately. New possibilities of high-resolution measurements through advanced sensor technology and data processing techniques have the potential to achieve more accurate modeling of transport.

### Authors

Jonghyun Harry Lee \*

Stanford University

Technical University of  
Denmark

Massimo Rolle

Peter K Kitanidis  
Stanford University

[Find Similar](#)

### View Related Events

**Day:** Wednesday, 14 December 2016

