



## Simulations of filamentary transport in the SOL and estimations of the power deposition profile

Rasmussen, Jens Juul; Nielsen, Anders Henry; Madsen, Jens; Naulin, Volker; Olsen, Jeppe Miki Busk; Magnussen, Michael Løiten; Thrysoe, Alexander Simon; Tophøj, Laust Emil Hjerrild; Coelho, R.

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Rasmussen, J. J., Nielsen, A. H., Madsen, J., Naulin, V., Olsen, J. M. B., Magnussen, M. L., Thrysoe, A. S., Tophøj, L. E. H., & Coelho, R. (2016). *Simulations of filamentary transport in the SOL and estimations of the power deposition profile*. Abstract from 21st EU-US Transport Task Force Meeting, Leysin, Switzerland.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Simulations of filamentary transport in the SOL and estimations of the power deposition profile


J. Juul Rasmussen<sup>1</sup>, A. H. Nielsen<sup>1</sup>, J. Madsen<sup>1</sup>, V. Naulin<sup>1</sup>, J.M.B. Olsen<sup>1</sup>,  
M. Løiten<sup>1</sup>, A.S. Thrysøe<sup>1</sup>, L. Tophøj<sup>1</sup>, and R. Coelho<sup>2</sup>

<sup>1</sup>DTU Physics, Technical University of Denmark, Kgs. Lyngby, Denmark

<sup>2</sup>Instituto de Plasmas e Fusão Nuclear, University of Lisboa, Portugal

Corresponding author: Jens Juul Rasmussen Email: jjra@fysik.dtu.dk

The transport in the edge region and through the scrape-off-layer (SOL) at the outboard mid-plane of magnetically confined toroidal plasmas is known to be strongly intermittent and dominated by filamentary structures of enhanced plasma pressure – plasma blobs. These structures are generated in the edge region and propagate through the SOL with velocities up to a few percent of the ion acoustic speed. They play a significant role in setting up the plasma profiles in the SOL and for the power deposition profiles. We have investigated the transport dynamics, the ensuing density and pressure profiles in the SOL, and estimated power depositions on plasma facing components by applying the HESEL code. HESEL is an energy conserving four-field Braginskii model governing the



dynamical evolution of generalized vorticity, density, electron, and ion pressures [1]. The parallel dynamics in the SOL is parametrized and different divertor conditions - attached or detached - are mimicked by applying sheath boundary conditions to either the full potential field or only its profile, respectively. This significantly affects the evolution of the plasma blobs and thus the transport of energy and particles through the SOL. We further observe an asymmetry between the ion and electron temperature both in the blob structures and in the background SOL plasma. Ion temperatures are typical 2-4 times higher than the electron temperature, which is a result of different parallel and perpendicular dissipation rates. The power deposition profile at the divertor is obtained by mapping the mid-plane pressure profiles to the divertor region along the magnetic field lines [2], see figure (for the “attached” case) where the traces of the individual blob

structures are clearly identified. We have investigated different L-mode scenarios including cases with high upstream SOL density – the so-called “shoulder” formation - recently characterized in JET and ASDEX Upgrade [3]. Within the present model we observe - for the same divertor condition – the formation of a ‘shoulder’ in the density profile by increasing the connection length, thus decreasing the efficiency of the SOL ability to remove plasma from the outboard midplane.

HESEL is embedded into a Kepler workflow developed within the EUROfusion Integrated Modeling framework [2]. This consists of a workflow allowing access to experimental Langmuir probe data as well as discharge parameters.

[1] J. Madsen et al. Phys. Plasmas **23**, 032306 (2016).

[2] A.H. Nielsen et al. 42<sup>nd</sup> EPS Conference on Plasma Physics, Lisbon, Portugal 2015, P1.105, ECA Vol. **39E**.

[3] D. Carralero et al. J. Nucl. Mat. **463**, 123, (2015); Phys. Rev. Lett. **115**, 215002 (2015).