Technical University of Denmark

Laboratory studies on the effect of ozonation on THM formation in swimming pool water

K.M.S. Hansen, A. Spiliotopoulou, W.A. Cheema, H.R. Andersen

6th International Conference Swimming Pool & Spa
17 – 20 March, 2015
Amsterdam, The Netherlands



DTU Environment
Department of Environmental Engineering

Ozone chemistry



t½, 20 °C = 3 days

t½, 20 °C = 20 min

Increased pH

$$O_3 + H_2O \xrightarrow{^-OH} 2\,{}^{\cdot}OH + O_2$$

Contaminants

Low reactivity:

${}^{\cdot}OH + R \rightarrow R_{ox}$

Urea



Organo-chlorine



High reactivity:

$O_3 + R \rightarrow R_{ox}$

double bond

Ozone chemistry

I. Organic matter reactive with ozone

- Direct oxidation by ozone
- Fast consumption of ozone
- Decrease chlorine reactivity of pollutants
- Low ozone life time → no reaction with bromide

II. Organic matter not reactive with ozone

- Slow consumption of ozone
- Ozone converts to hydroxyl radicals with time
- Radical attack of inactive carbon → increased chlorine reactivity
- Long ozone life time → oxidation of bromide to bromate

Systems: DIN (German standard)



Systems: Slip-stream (USA)



DEL Ozone proprietary drawing

Our aim

Optimizing ozone treatment

- Contact time
- Required ozone dose to minimize chlorination DBPs

Ozone kinetics not well described in swimming pools

- Effects of pH
- Effects of organic, repeated ozonation

Effect of ozonation on DBP formation

- Pool water
- Filling water
- Fresh pollutants

Lab-scale experiments



Effect of pH on ozone lifetime



- Very little difference in the pH range swimming pools are operated

Repeated ozonation – ozone lifetime





- Fast removal of 1st ozone dose → ozone reactive material
- 2nd and 3rd → no ozone reactive material

- Little difference → no ozone reactive material in the water

Chlorination of the ozonated samples

Chlorine: 24 h at 25 °C → Chlorine residual +

Purge & trap – GC/MS

Chlorine consumption



I. Direct oxidation by ozone

II. Radical mechanism

- Different chlorine consumption for tap water and swimming pool water

Formation of total trihalomethane



TTHM

Incorporated Br

I. Direct oxidation by ozone

II. Radical mechanism

- Ozonation of filling water → decreased THM
- Ozonation of swimming pool water → increased THM

Fresh pollutants

Why do not we see an decrease in THM formation in pool water?



+

+



- Control
- 0.5 ppm O_3
- 1.0 ppm
- 1.5 ppm
- 3.0 ppm
- +100 ppm

Total organic carbon (TOC) and chlorine consumption





- Ozonation decrease TOC and decreased chlorine consumption

TTHM formation



- Fresh pollutant direct ozone reaction → short ozone life time → no decrease in formation of brominated THM

Conclusions

Kinetic possible

- pH minimal effect
- TOC (ozone reactive vs non-reactive)

Chlorinated pool water

- Reaction via hydroxyl radical, making the organic material more reactive and increased THM formation.

Presence of fresh pollutants

- Direct oxidation by ozone → decreased THM formation

- Still need some more information on ozone dose required

Thanks for your attention

See you at the next pool conference with results from ozonation of full scale system

DTU Environment
Department of Environmental Engineering