



## Acclimation of continuous biomethanation process to extremely high ammonia levels

Tian, Hailin; Mancini, Enrico; Fotidis, Ioannis; Angelidaki, Irini

*Publication date:*  
2016

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Tian, H., Mancini, E., Fotidis, I., & Angelidaki, I. (2016). *Acclimation of continuous biomethanation process to extremely high ammonia levels*. Abstract from Biogas Science 2016, Szeged, Hungary.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Acclimation of continuous biomethanation process to extremely high ammonia levels

**Hailin Tian; Enrico Mancini; Ioannis A. Fotidis\*; Iriini Angelidaki**

(Department of Environmental Engineering, Technical University of Denmark,  
Building 113, DK-2800 Kgs. Lyngby, Denmark)

(\*Corresponding author: [ioanf@env.dtu.dk](mailto:ioanf@env.dtu.dk))

## Abstract

Anaerobic digestion (AD) is an effective technology to recover energy (i.e. biogas) from biomass. However, when protein-rich bioproducts are used as feedstocks, could lead to low methane yields and even to complete process failure. The reason is that high ammonia levels from protein degradation could inhibit the microbes mediating the AD process. Acclimatized methanogenic communities to high ammonia levels can offer a solution to this problem. Nevertheless, the acclimation of ammonia tolerant methanogenic communities for continuous AD of protein-rich substrates still poses serious problems (e.g. washout effect). Thus, the present study investigates the acclimation of methanogenic communities during AD of protein-rich bioproducts. Microalgae *Chlorella vulgaris* (>50% protein in dry matter), was co-digested with cattle manure (80/20 VS microalgae/VS manure) in two identical continuous stirred tank reactors (CSTR) under mesophilic conditions ( $37\pm 1^\circ\text{C}$ ). The reactors had a hydraulic retention time of 23 days and organic loading rate of  $1.95\pm 0.1 \text{ g VS L}^{-1}\text{d}^{-1}$ . Acclimation strategy was followed a stepwise increasing ( $0.5 \text{ g NH}_4^+-\text{N L}^{-1}$  each step) the total ammonia (TAN) concentration starting from 4 up to  $10 \text{ g NH}_4^+-\text{N L}^{-1}$  using  $\text{NH}_4\text{Cl}$ . During the baseline period ( $4 \text{ g NH}_4^+-\text{N L}^{-1}$ ) the methane yield was  $335 \text{ NmL CH}_4 \text{ g}^{-1} \text{ VS}$ . Throughout the acclimation period, the methane yields of the CSTR reactors were stable with an average variation less than 10% compared to the yield at baseline period. Additionally, pH decreased from 8.4 to 7.8 during the ammonia acclimation process. Owing to the pH drop, the free ammonia (FAN) concentration was kept in high but stable levels ( $920\pm 150 \text{ mg NH}_3\text{-N L}^{-1}$ ), even though the TAN increased 1.5-fold. This indicates that only TAN increase was not enough to cause inhibition and that FAN plays the most crucial role in the ammonia inhibitory mechanism. The present study showed that it is possible to acclimatize methanogenic communities to extremely high TAN levels. Finally, the results derived from this study could offer a practical approach to digest protein-rich bioproducts.

Keywords:

Adaptation; Ammonia inhibition; Anaerobic digestion; CSTR.