



Direct current (DC) resistivity and induced polarization (IP) monitoring of active layer dynamics at high temporal resolution

Doetsch, Joseph ; Ingeman-Nielsen, Thomas; Christiansen, Anders V.; Fiandaca, Gianluca; Auken, Esben; Adamson, Kathryn; Lane, Timothy; Elberling, Bo

Published in:
Geophysical Research Abstracts

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Doetsch, J., Ingeman-Nielsen, T., Christiansen, A. V., Fiandaca, G., Auken, E., Adamson, K., Lane, T., & Elberling, B. (2014). Direct current (DC) resistivity and induced polarization (IP) monitoring of active layer dynamics at high temporal resolution. *Geophysical Research Abstracts*.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Direct current (DC) resistivity and induced polarization (IP) monitoring of active layer dynamics at high temporal resolution

Joseph Doetsch (1), Thomas Ingeman-Nielsen (2), Anders V. Christiansen (1), Gianluca Fiandaca (1), Esben Auken (1), Kathryn Adamson (3), Timothy Lane (4), and Bo Elberling (5)

(1) HydroGeophysics Group, Department of Geoscience, Aarhus University, Aarhus, Denmark (joseph.doetsch@geo.au.dk), (2) Technical University of Denmark, Copenhagen, Denmark, (3) University of London, United Kingdom, (4) Université Paris, France, (5) Center for Permafrost (CENPERM), University of Copenhagen, Denmark

With climatic changes, permafrost thawing and changes in active layer dynamics influencing microbial activity and greenhouse gas feedbacks to the climate system, understanding of the interaction between biogeochemical and thermal processes in the ground is of increasing interest. Here we present results of from an on-going field experiment, where the active layer dynamics are monitored using direct current (DC) resistivity and induced polarization (IP) measurements at high temporal resolution. These DC/IP measurements are supplemented by pore water analysis, continuous ground temperature monitoring (0-150 cm depth) and structural information from ground penetrating radar (GPR).

The study site (N69°15', W53°30', 30 m a.s.l.) is located at a Vaccinium/Empetrum heath tundra area near the Arctic Station on Qeqertarsuaq on the west coast of Greenland. Mean air temperatures of the warmest (July) and the coldest (February–March) months are 7.1 and –16.0°C, respectively. The DC/IP monitoring system was installed in July 2013 and has since been acquiring at least 6 data sets per day on a 42-electrode profile with 0.5 m electrode spacing. Recorded data include DC resistivity, stacked full-decay IP responses and full waveform data at 1 kHz sampling frequency. The monitoring system operates fully automatic and data are backed up locally and uploaded to a web server.

Time-lapse DC resistivity inversions of data acquired during the freezing period of October – December 2013 clearly image the soil freezing as a strong increase in resistivity. While the freezing horizon generally moves deeper with time, some variations in the freezing depth are observed along the profile. Comparison with soil temperature measurements at different depths indicates a linear relationship between the logarithm of electrical resistivity and temperature. Preliminary time-lapse inversions of the full-decay induced polarization (IP) data indicate a decrease of chargeability with freezing of the ground. Once laboratory analysis of the pore water is available, we will assess if these changes in chargeability are caused by geochemical changes of the pore water.