



Low nitrous oxide production in intermittent-feed high performance nitrifying reactors

Su, Qingxian; Jensen, Malene M.; Smets, Barth F.

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Su, Q., Jensen, M. M., & Smets, B. F. (2017). *Low nitrous oxide production in intermittent-feed high performance nitrifying reactors*. Abstract from Frontiers International Conference on Wastewater Treatment (FICWTM2017), Palermo, Italy.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Low nitrous oxide production in intermittent-feed high performance nitritating reactors

Qingxian Su*, Marlene M. Jensen*, Barth F. Smets*

* Department of Environmental Engineering, Technical University of Denmark, 2800, Lyngby, Denmark

Nitrous oxide, nitrification, ammonia-oxidizing bacteria, nitrous oxide production pathways, intermittent feeding, pH

Summary

Nitrous oxide (N₂O) production from autotrophic nitrogen removal processes, especially nitritating systems, is of growing concern. N₂O dynamics were characterized and N₂O production factors were quantified in two lab-scale intermittent-feed nitritating SBRs. 93 ± 14% of the oxidized ammonium was converted to nitrite, with the average total net N₂O production of 2.1 ± 0.7% of the ammonium oxidized. Operation with intermittent feeding appears an effective optimization approach to mitigate N₂O emissions from nitritating systems. Net N₂O production rates transiently increased with a rise in pH after each feeding, indicating a potential role of pH in N₂O production.