



## Synthesis of branched–backbone oligosaccharides of the pectic RG-I plant cell wall polysaccharide

Awan, Shahid Iqbal; Clausen, Mads Hartvig

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Awan, S. I., & Clausen, M. H. (2017). *Synthesis of branched–backbone oligosaccharides of the pectic RG-I plant cell wall polysaccharide*. Abstract from 19th European Carbohydrate Symposium, Barcelona, Spain.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.




## SYNTHESIS OF BRANCHED-BACKBONE OLIGOSACCHARIDES OF THE PECTIC RG-I PLANT CELL WALL POLYSACCHARIDE

Shahid I. Awan<sup>2</sup>, Mads H. Clausen<sup>1,2</sup>

<sup>1</sup> Center for Nanomedicine and Theranostics, <sup>2</sup> Department of Chemistry, Technical University of Denmark, Kemitorvet 207, 2800 Kgs. Lyngby, Denmark.  
E-Mail: shia@kemi.dtu.dk

Plants are an essential part of life on earth. They are the primary food producers, climate regulators and provide habitats for other organisms [1]. The dependence of industrialized nations on plant cell walls due to their industrial applications has rapidly increased. Cellulose, hemicelluloses, and pectin polysaccharides are the main structural components of the plant cell wall. Among plant carbohydrates, pectins are highly heterogeneous polysaccharides. They are mainly found in the primary plant cell wall and contribute to various cell functions, including support, defense, signaling, and cell adhesion [2]. Rhamnogalacturonan I (RG-I) is one of the structural classes of pectic polysaccharides, along with homogalacturonan, rhamnogalacturonan II, and xylogalacturonan. The chemical structure of RG-I is complex having a backbone consisting of alternating  $\alpha$ -linked L-rhamnose and D-galacturonic acid units with numerous branches of galactan, arabinan, or arabinogalactan positioned at C-4 of the rhamnose residues [3].



**Figure 1.** Branched-backbone synthetic target of the pectic RG-I polysaccharide.

The use of defined oligosaccharides rather than isolated polysaccharides can aid in obtaining detailed information about biosynthetic pathways, plant evolution, and agronomical properties. Furthermore, biological testing can provide new insight into plant biology; important for plant preservation, engineering, and utilization of plants as a source of bioenergy. Present work towards defined RG-I substructures involves a [4+3]-coupling to furnish a heptasaccharide backbone unit (see Figure 1). Moreover, installation of side chains of different lengths and nature can be installed at the C-4 position of rhamnose unit. Finally, these oligosaccharides will be deprotected to furnish target molecules to pursue biological studies.

[1] Hipp, A., Gibson, J.P., Gibson, T.R. *Plant Diversity (The Green World)*. **2007**, Chelsea House Pub. ISBN-10: 0791089606.

[2] a) Mohnen, D. *Curr. Opin. Plant Biol.* **2008**, 11, 266–277. Harholt, J.; Suttangkakul, A.; Scheller, H. V. *Plant Physiol.* 2010, 153, 384–395; b) Willats, W. G. T.; Knox, J. P.; Mikkelsen, J. D. *Trends Food Sci. Technol.* **2006**, 17, 97–104.

[3] Zakharova, A. N.; Madsen, R.; Clausen, M. H. *Org. Lett.* **2013**, 15, 1826-1829.