



Simulating control of paratuberculosis in Danish dairy herds

Kirkeby, Carsten Thure; Græsbøll, Kaare; Nielsen, Søren S.; Toft, Nils; Hisham Beshara Halasa, Tariq

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Kirkeby, C. T., Græsbøll, K., Nielsen, S. S., Toft, N., & Hisham Beshara Halasa, T. (2017). *Simulating control of paratuberculosis in Danish dairy herds*. Abstract from Copenhagen Cattle 2017, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Simulating control of paratuberculosis in Danish dairy herds

Carsten Kirkeby¹, Kaare Græsbøll¹, Søren S. Nielsen², Nils Toft¹ and Tariq Halasa¹

¹*Simulation modeling group, DTU Veterinary Institute, Technical University of Denmark.*

²*Section for Animal Welfare and Disease Control, Department of Veterinary and Animal Sciences (IVH), University of Copenhagen*

Paratuberculosis

Paratuberculosis (PTB) is a chronic disease caused by *Mycobacterium avium* ssp. *paratuberculosis* (MAP). It causes economic losses and reduced animal welfare in dairy cattle herds worldwide. Animals are infected through ingested MAP shed in the feces and milk of infected animals in the same herd. Infected animals remain subclinical for a long period before they get clinical, often resulting in fatal diarrhea. Simulation models of dairy herds can be used to predict the effect of control and prevention strategies, and are especially suited for this type of slowly developing disease.

Control actions

The median within-prevalence of PTB in Danish dairy herds is around 6% and 50% of the Danish dairy cattle herds are closed (i.e. they do not purchase animals into the herd). We here simulated a closed Danish dairy herd with 200 lactating cows and a normal within-herd prevalence of PTB using the stochastic mechanistic bio-economic simulation model "iCull". Additionally, we simulated a herd with lower hygiene and thus with a higher within-herd prevalence. In the simulations we evaluated the control actions recommended in the Danish control program for PTB. These actions were: removing calves from the dam right after birth; avoiding feeding calves with colostrum from infected cows; avoiding feeding calves with colostrum from infected cows; and a strict test-and-cull strategy. From repeated simulations, the impact of each recommended action was estimated from the results. The three first-mentioned recommended actions did not have much impact on the prevalence alone, but we found a synergistic effect when combining these effects. The test-and-cull strategy was by far the most efficient action, resulting in the economically most optimal situations.

Purchase of animals

We also explored the consequences of purchasing cattle from herds possibly infected with PTB, representing the other 50% of the Danish herds that are not closed. Herds that were already infected did not have a high risk of increasing their prevalence. Herds without infected animals were highly susceptible to PTB and should avoid purchasing animals from infected herds. We also estimated the probability of fadeout of PTB once a herd was infected. This depends on the number of infected animals. If only one infected animal was purchased into a disease-free herd, we found 18% probability that the herd would still be infected after ten simulated years.