



## **Genome Sequence of *Enterococcus mundtii* EM01, Isolated from *Bombyx mori* Midgut and Responsible for Flacherie Disease in Silkworms Reared on an Artificial Diet**

**de Diego-Diaz, Beatriz; Treu, Laura; Campanaro, Stefano; da Silva Duarte, Vinicius; Saviane, Alessio; Cappelozza, Silvia; Squartini, Andrea**

*Published in:*  
Genome Announcements

*Link to article, DOI:*  
[10.1128/genomeA.01495-17](https://doi.org/10.1128/genomeA.01495-17)

*Publication date:*  
2018

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
de Diego-Diaz, B., Treu, L., Campanaro, S., da Silva Duarte, V., Saviane, A., Cappelozza, S., & Squartini, A. (2018). Genome Sequence of *Enterococcus mundtii* EM01, Isolated from *Bombyx mori* Midgut and Responsible for Flacherie Disease in Silkworms Reared on an Artificial Diet. *Genome Announcements*, 6(3), Article e01495-17. <https://doi.org/10.1128/genomeA.01495-17>

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Genome Sequence of *Enterococcus mundtii* EM01, Isolated from *Bombyx mori* Midgut and Responsible for Flacherie Disease in Silkworms Reared on an Artificial Diet

Beatriz de Diego-Diaz,<sup>a,b</sup>  Laura Treu,<sup>a</sup> Stefano Campanaro,<sup>c</sup> Vinicius da Silva Duarte,<sup>d</sup> Alessio Saviane,<sup>e</sup> Silvia Cappelozza,<sup>e</sup> Andrea Squartini<sup>f</sup>

<sup>a</sup>Department of Environmental Engineering, Technical University of Denmark, Kongens Lyngby, Denmark

<sup>b</sup>Department of Chemistry, University of Navarra, Navarra, Spain

<sup>c</sup>Department of Biology, University of Padua, Padua, Italy

<sup>d</sup>Department of Microbiology, Federal University of Viçosa, Viçosa, Brazil

<sup>e</sup>CREA Centro di Ricerca Agricoltura e Ambiente, Padua, Italy

<sup>f</sup>Department of Agronomy, Food, Natural Resources, Animals and Environment (DAFNAE), Legnaro, PD, Italy

**ABSTRACT** The whole genome sequence of *Enterococcus mundtii* strain EM01 is reported here. The isolate proved to be the cause of flacherie in *Bombyx mori*. To date, the genomes of 11 other *E. mundtii* strains have been sequenced. EM01 is the only strain that displayed active pathological effects on its associated animal species.

*Enterococcus mundtii* is a taxon stemming from the reclassification of different enterococcal bacteria that formerly included *Streptococcus (Enterococcus) faecalis* and *Enterococcus faecium* (1). Its role as a potential pathogen has been signaled (2–4), as has its apparently commensal presence within the guts of different vertebrate and invertebrate animals (5). Being capable of lactic fermentation, its occurrence in fermented dairy products is also documented (6, 7).

There are already 11 available genome sequences from this species. The sequenced strains were isolated from different sources and are QU25 (8, 9), EMB156 (GenBank accession number NZ\_CP022340), ATCC 882 (10), CRL1656 (11), CRL35 (6), QAUEM2808 (7), SL\_16 (12), DSM 4838 (NZ\_JXKV01000001), CGB1038-1\_S1 (NZ\_MSTR01000001), 6B1\_DIV0119 (NZ\_NGMS01000001), and C2 (NZ\_FOUC01000001). The genome sequences are complete for two of these strains, QU25 and EMB156.

Strain EM01 was isolated in Italy in 2008 from 5th-instar larvae of the silkworm, *Bombyx mori*, reared on a nonsterile artificial diet based on mulberry leaves (13). EM01 was shown to be the causal agent of flacherie disease in silkworms. The only other available genome sequence of an *E. mundtii* isolate from the same host (strain EMB156) came instead from the guts of healthy *B. mori* silkworms reared in China. In fact, reports of *E. mundtii* isolation from silkworms in that country judged its occurrence as potentially beneficial for the same host (14). This makes the present report particularly relevant, as it allows inspection of whether distinctive genetic traits could possibly differentiate an ascertained pathogenic version of the taxon from its supposedly nonthreatening counterpart.

An Illumina MiSeq sequencer (Ramaciotti Centre, Australia) was used for the whole-genome sequencing of strain EM01. The genomic libraries used for this purpose were obtained with the Nextera XT kit (Illumina, Inc., USA). Up to 1,918,184 paired-end reads (2 × 250 bp) were generated, and these yielded 153-fold coverage of the studied genome. The assembly of 99.4% of these reads generated 54 scaffolds. All the aforementioned analyses were performed with version 10.1.1 of the CLC Genomics Workbench software (Qiagen Bioinformatics, Germany) (5). The draft genome of

Received 30 November 2017 Accepted 5 December 2017 Published 18 January 2018

**Citation** de Diego-Diaz B, Treu L, Campanaro S, da Silva Duarte V, Saviane A, Cappelozza S, Squartini A. 2018. Genome sequence of *Enterococcus mundtii* EM01, isolated from *Bombyx mori* midgut and responsible for flacherie disease in silkworms reared on an artificial diet. *Genome Announc* 6:e01495-17. <https://doi.org/10.1128/genomeA.01495-17>.

**Copyright** © 2018 de Diego-Diaz et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Laura Treu, [latr@env.dtu.dk](mailto:latr@env.dtu.dk).

*E. mundtii* EM01 is 3,134,567 bp in length, with a mean G+C content of 38.2%. The  $N_{50}$  for this isolate reached 136,907 bp.

Rapid Annotations using Subsystems Technology (RAST) (15) was used for genome annotation, and 2,994 coding sequences were detected, as were 53 structural RNAs. Up to 51 of the genes detected are related to virulence, disease, and defense, and 33 are phage-associated sequences. No clusters of regularly interspaced short palindromic repeats were detected. When comparing the sequence of this genome with the other whole-genome sequences reported in the literature, it can be seen how the gene identification for EM01 is slightly better than that for all others except QU25 (3,229).

Finally, a BLAST search was performed using the 16S rRNA gene sequence, and 100% identity and query coverages were found with *E. mundtii* ATCC 43186 (GenBank accession number NR\_024906). The genome sequence was used as input for PHASTER software. PHASTER analysis yielded two intact and four incomplete prophage regions, which is consistent with analysis results from the other reference genomes.

**Accession number(s).** This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number PHIL00000000. The version described in this paper is version PHIL01000000.

## REFERENCES

- Collins MD, Farrow JAE, Jones D. 1986. *Enterococcus mundtii* sp. nov. Int J Syst Bacteriol 36:8–12. <https://doi.org/10.1099/00207713-36-1-8>.
- Higashide T, Takahashi M, Kobayashi A, Ohkubo S, Sakurai M, Shirao Y, Tamura T, Sugiyama K. 2005. Endophthalmitis caused by *Enterococcus mundtii*. J Clin Microbiol 43:1475–1476. <https://doi.org/10.1128/JCM.43.3.1475-1476.2005>.
- Kauffhold A, Ferrieri P. 1991. Isolation of *Enterococcus mundtii* from normally sterile body sites in two patients. J Clin Microbiol 29:1075–1077.
- Raja NS, Ul Haq RN, Janjua KA. 2006. Management of vancomycin-resistant enterococci infections. Infect Dis J Pak 15:13–19.
- Ferreira AE, Canal N, Morales D, Fuentefria DB, Corção G. 2007. Characterization of enterococci produced by *Enterococcus mundtii* isolated from humans feces. Braz Arch Biol Technol 50:249–258. <https://doi.org/10.1590/S1516-89132007000200010>.
- Bonacina J, Saavedra L, Suárez NE, Sesma F. 2014. Draft genome sequence of the nonstarter bacteriocin-producing strain *Enterococcus mundtii* CRL35. Genome Announc 2:e00444-14. <https://doi.org/10.1128/genomeA.00444-14>.
- Farah N, Mehdi A, Soomro SI, Soomro NI, Tareb R, Desmasures N, Vernoux JP, Bakhtiar SM, Imran M. 2016. Draft genome sequence of *Enterococcus mundtii* QAUEM2808, isolated from dahi, a fermented milk product. Genome Announc 4:e00995-16. <https://doi.org/10.1128/genomeA.00995-16>.
- Abdel-Rahman MA, Tashiro Y, Zendo T, Shibata K, Sonomoto K. 2011. Isolation and characterisation of lactic acid bacterium for effective fermentation of cellobiose into optically pure homo L-(+)-lactic acid. Appl Microbiol Biotechnol 89:1039–1049. <https://doi.org/10.1007/s00253-010-2986-4>.
- Shiwa Y, Yanase H, Hirose Y, Satomi S, Araya-Kojima T, Watanabe S, Zendo T, Chibazakura T, Shimizu-Kadota M, Yoshikawa H, Sonomoto K. 2014. Complete genome sequence of *Enterococcus mundtii* QU 25, an efficient L-(+)-lactic acid-producing bacterium. DNA Res 21:369–377. <https://doi.org/10.1093/dnares/dsu003>.
- Linden BA, Turner WR, Thom C. 1926. Public health weekly reports for August 6, 1926. Publ Health Rep 41:1647–1694.
- Magni C, Espeche C, Repizo GD, Saavedra L, Suárez CA, Blancato VS, Espariz M, Esteban L, Raya RR, Font de Valdez G, Vignolo G, Mozzi F, Taranto MP, Hebert EM, Nader-Macías ME, Sesma F. 2012. Draft genome sequence of *Enterococcus mundtii* CRL1656. J Bacteriol 194:550. <https://doi.org/10.1128/JB.06415-11>.
- Chen B, Sun C, Liang X, Lu X, Gao Q, Alonso-Pernas P, Teh BS, Novoselov AL, Boland W, Shao Y. 2016. Draft genome sequence of *Enterococcus mundtii* SL 16, an indigenous gut bacterium of the polyphagous pest *Spodoptera littoralis*. Front Microbiol 7:1676. <https://doi.org/10.3389/fmicb.2016.01676>.
- Cappellozza S, Saviane A, Tettamanti G, Squadrin M, Vendramin E, Paolucci P, Franzetti E, Squartini A. 2011. Identification of *Enterococcus mundtii* as a pathogenic agent involved in the “flacherie” disease in *Bombyx mori* L. larvae reared on artificial diet. J Invertebr Pathol 106:386–393. <https://doi.org/10.1016/j.jip.2010.12.007>.
- Fei C, Lu XM, Qian YH, Zhang H, Mahmood Q. 2006. Identification of *Enterococcus* sp. from midgut of silkworm based on biochemical and 16S rDNA sequencing analysis. Ann Microbiol 56:201–205. <https://doi.org/10.1007/BF03175006>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. BMC Genomics 9:75. <https://doi.org/10.1186/1471-2164-9-75>.