



Data Requirements and Modeling for Gas Hydrate-Related Mixtures and a Comparison of Two Association Models

Liang, Xiaodong; Aloupis, Georgios; Kontogeorgis, Georgios M.

Published in:
Journal of Chemical and Engineering Data

Link to article, DOI:
[10.1021/acs.jced.7b00081](https://doi.org/10.1021/acs.jced.7b00081)

Publication date:
2017

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Liang, X., Aloupis, G., & Kontogeorgis, G. M. (2017). Data Requirements and Modeling for Gas Hydrate-Related Mixtures and a Comparison of Two Association Models. *Journal of Chemical and Engineering Data*, 62(9), 2592-2605. <https://doi.org/10.1021/acs.jced.7b00081>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Data requirements and modeling for gas hydrate related mixtures and a comparison of two association models

*Xiaodong Liang, Georgios Aloupis, Georgios M. Kontogeorgis**

Center for Energy Resources Engineering (CERE),

Department of Chemical and Biochemical Engineering,

Technical University of Denmark, 2800 Kgs. Lyngby, Denmark

Email: gk@kt.dtu.dk

Abstract

The association theory based advanced thermodynamic models gain more and more attention and applications in many industries. The Cubic Plus Association (CPA) and the simplified Perturbed-Chain Statistical Associating Fluid Theory (sPC-SAFT) equations of state (EOS) are two of the most widely used association models in the chemical and petroleum industries. The CPA model is extensively used in flow assurance, in which gas hydrate is one of the central topics. Experimental data play a vital role in validating models and obtaining model parameters. In this work, we will compare the performance of the CPA and sPC-SAFT EOS for modeling the fluid phase equilibria of gas hydrate related systems, and try to explore how the models can help in suggesting experimental measurements. These systems contain water, hydrocarbon (alkane or aromatic) and either methanol or mono-ethylene glycol. It is well-known that the determination of SAFT-type model parameters for associating fluids remains a challenge, since there are at least five pure component parameters for these compounds, and there is no property combination found yet to be enough to ensure the best parameter set. Therefore, in this work two parameter sets have been chosen for the sPC-SAFT EOS for a fair comparison. The comparisons are made for pure fluid properties, vapor-liquid equilibria and liquid-liquid equilibria of binary and ternary mixtures, as well as vapor-liquid-liquid equilibria of quaternary mixtures. The results show, from an overall point of view, that these two models have equally good performance, and the two parameter sets with the sPC-SAFT EOS are also comparable, especially for the vapor-liquid equilibria systems. Moreover, the modeling results suggest that some data are less reliable than others, which indicates the need for more measurements to further validate the models, especially for multicomponent systems.

Keywords: Gas hydrate, inhibitor, CPA, sPC-SAFT, Vapor-liquid-liquid equilibria

1. Introduction

Fair comparisons of thermodynamic models are rare in literature. When comparisons are only done for the correlative performance, e.g. binary phase equilibria, differences can be masked by the adjustable parameters which often play an important role. This is the case even when classical thermodynamic models like cubic equations of state (EOS) are compared against modern Statistical Associating Fluid Theory (SAFT)-type or similar EOS. Thus, the true value of thermodynamic models should be based on predictive calculations, e.g. multicomponent phase equilibria or prediction of properties other than those used in the parameter estimation, both of which are also important for engineering applications. In this work we focus on the multicomponent multiphase equilibria of systems of relevance to the petroleum industry (water-alcohols/glycols-hydrocarbons). While such multicomponent systems are of great importance, not many data are available, possibly due to the experimental difficulties, e.g. the detection of low solubilities in some of the phases. The corresponding author recalls a communication with Kenneth Marsh some years ago discussing a manuscript published in Journal of Chemical & Engineering Data. That manuscript contained experimental data for many water-glycol-hydrocarbon systems, but unfortunately they were not accurate. The corresponding author appreciated Kenneth Marsh's interference in this case and with his continuous effort the authors repeated some of the measurements and today we have some of the best water-monoethylene glycol (MEG)-alkane LLE data available in the literature.¹ This is highly appreciated as such data are extremely rare and only very few laboratories can measure them.

This work focuses on the fluid phase equilibria of gas hydrate related systems. A combination of appropriate temperature and pressure conditions as well as the presence of light gases may lead water molecules to reform into coplanar poly-membered (5 or 6) rings and create a 3D polyhedral shaped formations with the light gases trapped within it. In this way, the water acts like a cage. This

phenomenon is called gas hydrate formation, which is considered as one of the major concerns in production, transportation and processing in the petroleum industry. This is because, when it occurs, it can reduce the flow of the gas in the pipeline and eventually block the pipeline stopping the production, and it might also affect and damage the equipment.² Addition of chemicals like alcohols and glycols, called thermodynamic gas hydrate inhibitors, is one widely used technique to prevent gas hydrate formation in the oil and gas industry. Methanol and MEG are common choices, while ethanol is an alternative. Methanol and MEG are both effective and reliable in preventing gas hydrate formation. MEG gains more attention nowadays, because it can be reused and regenerated with insignificant losses, and decreases the possibility of creation and occurrence of corrosion in the pipelines.³

Thermodynamic models play a vital role in determining the minimum necessary amount of chemicals used as gas hydrate inhibitors. The Cubic Plus Association (CPA) EOS⁴ has been shown to be very useful in predicting gas hydrate formation, based on extensive validations for many gas hydrate related systems.⁵⁻⁷ The CPA EOS uses the same association term of the SAFT models,⁸⁻¹³ and one of the most widely used SAFT models, the Perturbed Chain-SAFT (PC-SAFT) EOS^{14,15} and its simplified version (sPC-SAFT),¹⁶ has gained more and more attention in the petroleum and chemical industries.⁵ Therefore, a systematic investigation on the performance of the sPC-SAFT EOS in modeling the fluid phase equilibria of gas hydrate related systems might further contribute to the acceptance of this model in the petroleum industry, and it might provide an alternative to cubic EOS for engineers. Moreover, a thorough and fair comparison is believed necessary to conclude which model or modeling approach may be better, and it will also help in giving more confidence of existing and/or suggesting more experimental measurements, when necessary.

Many works have been done in modeling systems containing hydrocarbon, water, methanol or MEG with the association theory based models.¹⁷⁻²² However, only few comparisons among these models have been published for multicomponent multiphase equilibria, and one of the most recent studies was for water-methanol-hydrocarbons with the CPA and sPC-SAFT EOS.²³ This work will focus on a fair comparison of these two models on their performance in modeling gas hydrate related systems in both correlation and prediction manners. The rest of the work is organized as (1) brief introduction of the two models in section 2; (2) presentation of the experimental data in section 3; (3) modeling results and discussion in section 4 and (4) conclusion and suggestions in section 5.

2. Models

In the past three decades, the association theory based thermodynamic models have been gaining attention and applications in many areas, including oil & gas, chemicals, environments and pharmaceuticals. The CPA and sPC-SAFT EOS are, among the association models, two of the most widely used models in the chemical and petroleum industries.⁵

2.1 CPA EOS

The CPA EOS was proposed by Kontogeorgis et al.⁴ and it is a combination of the SRK (or other cubic) EOS, widely used in the petroleum industry, and the association term of the SAFT type models. The CPA reduces to SRK in the absence of hydrogen bonding compounds, leading to a good balance between accuracy and simplicity, thus it helps in gaining easy acceptance in the oil, gas and chemical industries. In terms of pressure, the CPA EOS can be expressed for mixtures as:

$$P = \frac{RT}{V_m - b} - \frac{a(T)}{V_m(V_m + b)} - \frac{1}{2} \frac{RT}{V_m} \left(1 + \rho \frac{\partial \ln g}{\partial \rho} \right) \sum_i x_i \sum_{A_i} (1 - X^{A_i}) \quad (1)$$

where ρ is the molar density ($\rho = 1/V_m$).

2.2 sPC-SAFT EOS

The PC-SAFT EOS was developed by Gross and Sadowski.¹⁴ In terms of the reduced residual Helmholtz free energy, it can be expressed as:

$$a^r = (a^{hs} + a^{chain}) + a^{disp} + a^{assoc} \quad (2)$$

where a^{hs} and a^{chain} are the contributions of the hard sphere segment-segment interaction and the chain formation, respectively. The dispersion force a^{disp} is built by extending the perturbation theory of Barker and Henderson²⁴ to use the hard-chain as the reference term. The term a^{assoc} represents the contributions of association forces of sites, which can be expressed as:

$$a^{assoc} = \sum_i x_i \left[\sum_{A_i} \left(\ln X^{A_i} - \frac{X^{A_i}}{2} \right) + \frac{M_i}{2} \right] \quad (3)$$

where M_i is the association site number of molecule i , and X^{A_i} is the fraction of molecules i not bonded at site A, given by:

$$X^{A_i} = \left[1 + \sum_j \sum_{B_j} \rho_j X^{B_j} \Delta^{A_i B_j} \right]^{-1} \quad (4)$$

where $\Delta^{A_i B_j}$ is the association strength between the site A_i and site B_j , which is obtained as:

$$\Delta^{A_i B_j} = N_{av} \sigma_{ij}^3 g^{hs} \kappa^{A_i B_j} \left[\exp \left(\frac{\varepsilon^{A_i B_j}}{kT} \right) - 1 \right] \quad (5)$$

In this work, the sPC-SAFT EOS proposed by von Solms et al.¹⁶ with both modifications is used. It simplifies the original PC-SAFT EOS and reduces the computational time, without loss of accuracy for many applications. The same pure component parameters can be used for both the original and simplified PC-SAFT EOS. The readers are referred to the literature^{5,14,16} for more details on equations and symbols. It is relevant to mention the combining rule of the association volume, which will be used

in some cases in the following discussion. For a binary system consisting of two associating compounds, e.g. water and methanol, the following combining rule is used

$$(\sigma_{ij}^3 \kappa^{A_i B_j}) = \sqrt{(\sigma_{ii}^3 \kappa^{A_i B_i})(\sigma_{jj}^3 \kappa^{A_j B_j})} \quad (6)$$

Between an associating compound and an aromatic one, e.g. water-benzene, however, the following combining rule is used

$$(\sigma_{ij}^3 \kappa^{A_i B_j}) = \kappa^{cross} \left(\frac{\sigma_{ii}^3 + \sigma_{jj}^3}{2} \right) \quad (7)$$

This combining rule is volume based, not the one commonly used for the diameters. It is inspired by the CPA EOS, in which the term σ_{ii}^3 is replaced by the co-volume parameter b . However, people may expect that similar results are obtained after fitting the cross association volume to the experimental data. More details of the models, equations and symbols are referred to the literature.^{4,5,14}

3. Experimental data

In this work, we focus on the relationship among experimental data, models and modeling approaches, instead of providing a complete literature review on all pertinent experimental data. Therefore, we selected the mixtures of water, methanol or MEG and hydrocarbon(s), as the performances of the models for these systems (water-gas hydrate inhibitors-oil) may be crucial in determining their acceptance by the petroleum industry. When the pure compound model parameters are available, the phase equilibrium data of binary mixtures are helpful in the models' correlative performance and testing the predictive capabilities of these pure component parameters, as well as in determining the binary interaction parameters which will be used in the next step for predictions of multicomponent systems. The collected data of binary and multicomponent mixtures of relevance to this study are listed in Table 1 and 2, respectively. Not all of these data are going to be used in the

modeling part, but they are useful in the discussion and in future studies. Some of the references contain data collections and recommendations, which may be useful as well.

Table 1. Experimental data review of binary mixtures of relevance to this study (polar compounds like water, alcohols and MEG with hydrocarbons and cross-associating systems of the polar compounds)

1 st comp.	2 nd comp.	T range (K)	P range (bar)	Ref.
Methane	Water	274.19-483.15	0.07-1334	25-32
Methane	Methanol	200.0-502.2	6.9-2935	25,32-35
Methane	Ethanol	298.15-498.15	5.0-364.2	36-38
Methane	MEG	283.2-398.15	1.1-396.2	32,39,40
Ethane	Water	274.26-373.15	3.2-49.5	28,32,41,42-44
Ethane	Methanol	283.2-303.2	5-67.7	32,45
Ethane	Ethanol	298.15-498.15	2.5-117.7	36,38
Ethane	MEG	283.2-303.2	5-40	32
n-Hexane	Water	270.0-490.0	3.4-29.6	46,47
n-Hexane	Methanol	245.15-339.60	0.1-1500	48-54
n-Hexane	Ethanol	328.15-351.45	0.5-1.0	55,56
n-Hexane	MEG	307.95-330.35	1.0	57
n-Octane	Water	310.9-539.1	0.1-784.5	58,59
n-Octane	Methanol	288.91-372.3	0.1-1500	60-62
n-Octane	Ethanol	313.15-348.15	0.04-0.95	63,64
Benzene	Water	279.15-453.15	0.06-20.2	65,66
Benzene	Methanol	298.15-363.15	0.1-3.1	67-72
Benzene	Ethanol	293.15-353.26	0.06-1.0	56,65,73-76
Benzene	MEG	279.2-342.1	1.0	77
Toluene	Water	273.15-458.15	0.02-17.0	66
Toluene	Methanol	313.15	0.08-0.37	76
Toluene	Ethanol	308.15-328.15	0.06-0.41	78,79
Toluene	MEG	279.1-361.0	1.0	77
Water	Methanol	298.15-523.15	0.04-85.1	80-83
Water	MEG	343.15-363.15	0.06-0.7	84

Table 2. Experimental data review of multicomponent mixtures of type water-alcohol/MEG/TEG-hydrocarbons and methanol/MEG-hydrocarbons.

1 st comp.	2 nd comp.	3 rd comp.	4 th comp.	T range (K)	P range (bar)	Ref.
Water	Methanol	Methane		267.55-313.45	42-400.5	25,28,85
Water	Methanol	Ethane		283.2-303.2	5-40	28
Water	Methanol	Propane		269.35-293.15	2.3-63.2	85,86
Water	Methanol	n-Butane		273.15-293.15	1.0	85
Water	Methanol	Hexane		293.2-323.15	1.0	87-89
Water	Methanol	Heptane		298.2	1.0	90
Water	Methanol	Benzene		293.15-333.15	1.0	87,91
Water	Methanol	Toluene		303.15	1.0	92
Water	MEG	Methane		283.2-303.2	50-400.6	28
Water	MEG	Ethane		283.2-303.2	5-40	28
Water	MEG	Hexane		283.15-323.15	1.0	1,93
Water	MEG	Nonane		313.15	1.0	94
Water	MEG	Benzene		298.2-323.2	1.0	77
Water	MEG	Toluene		298.2-323.2	1.0	77
Water	MEG	Ethylbenzene		313.15	1.0	94
Water	Ethanol	Propane		313-403	99.05	95
Water	Ethanol	Hexane		329.26-349.66	1.0	96
Water	Ethanol	Heptane		298.2	1.0	90
Water	Ethanol	Benzene		293.15-328.15	1.0	97,98
Water	Ethanol	Toluene		303.15	1.0	92
Water	TEG	Toluene		368.75-412.55	0.85	77,99
Methanol	Methane	Ethane		275.2-283.2	50-400	28
Methanol	Benzene	Heptane		298.15	1.0	61
Methanol	Benzene	Octane		298.15	1.0	61
Methanol	Benzene	Nonane		298.15	1.0	61
Methanol	Benzene	Decane		298.15	1.0	61
Methanol	Toluene	Heptane		298.15	1.0	61
Methanol	Toluene	Octane		298.15	1.0	61
Methanol	Toluene	Nonane		298.15	1.0	61
Methanol	Toluene	Decane		298.15	1.0	61
MEG	Methane	Ethane		275.2-283.2	49.9-400	28
Water	Methanol	Methane	Ethane	275.2-283.2	13-400	28
Water	Methanol	Methane	Propane	281.37-310.93	34.47-208.77	100
Water	Methanol	Methane	Heptane	263.15-323.15	69.02-70.74	101
Water	Methanol	Methane	Toluene	263.15-323.15	80.05-96.11	101
Water	MEG	Methane	Ethane	275.2-283.2	5-400	28
Water	MEG	Methane	Propane	273.15-283.15	69.02-70.39	102
Water	MEG	Methane	Hexane	263.15-283.15	68.95-69.02	102
Water	MEG	Methane	Octane	281.1-363.2	31-150	103
Water	MEG	Methane	Toluene	263.15-323.15	69.22-69.5	101

4. Modeling results and discussion

4.1 Pure component parameters

The first step in modeling fluid phase equilibria using EOS is to obtain the pure component parameters. The CPA and sPC-SAFT EOS have the same number of pure component parameters, i.e. three and five parameters for non-associating and associating fluids, respectively. In general, these parameters are obtained by fitting to vapor pressure and liquid density. It is well-known that the determination of SAFT parameters for associating fluids is a challenge, since there are at least five pure component parameters for these compounds, and there is no property combination found yet to be enough to ensure the best parameter set.¹⁰⁴⁻¹⁰⁶ Several parameter sets have been published for water, methanol and MEG¹⁰⁷ with the sPC-SAFT EOS. Two of these sets, while having rather different parameter values, provide equally good representation of vapor pressures and liquid densities, have been chosen for the sPC-SAFT EOS for a fair comparison. Based on our experience, the two-site association scheme 2B¹⁰⁸ is used for methanol in this work, while more sophisticated developments may be needed for the cross association for alcohol containing mixtures.¹⁰⁹ The pure component parameters are given in Tables 3 and 4 for the CPA and sPC-SAFT EOS, respectively.

Firstly, these parameters of the associating fluids are validated against the vapor pressure and liquid density data from the DIPPR¹¹⁰ and NIST^{111,112} databases in Table 5. The percentage average absolute deviations (%AAD), defined in the following equation

$$\%AAD(\Omega) = \frac{1}{Np} \sum_{i=1}^{Np} \left| \frac{\Omega^{cal}}{\Omega^{exp}} - 1 \right| \times 100\% \quad (8)$$

where Ω is vapor pressure, liquid density or solubility used in the following sections. Np is the number of experimental data points.

In addition, an average %AAD is defined for each model and each phase

$$\text{average \%AAD} = \frac{\sum \%AAD}{\text{Number of \%AAD}} \quad (9)$$

The vapor pressures of water from the CPA and sPC-SAFT EOS are also plotted in Figure 1. It can be seen from Table 5 and Figure 1 that these two models and the two parameter sets of the sPC-SAFT EOS have quite similar performance, and in general the data from the two databases are consistent with each other.

Table 3. The CPA parameters of relevant compounds

Comp.	b (L/mol)	$\Gamma = a_0/Rb$ (K)	c_1 (-)	$\varepsilon^{A_i B_i}/k$ (K)	$\kappa^{A_i B_i}$ (-)	Scheme (-)	Ref.
Methane	0.0291	959.028	0.44718	-	-	-	113
Ethane	0.04288	1544.55	0.58463	-	-	-	113
n-Hexane	0.10789	2640.03	0.8313	-	-	-	114
Benzene	0.07499	2867.19	0.7576	-	-	-	115
Toluene	0.09214	3051.36	0.8037	-	-	-	115
Methanol	0.030978	1573.71	0.43102	2957.78	16.1	2B	114
MEG	0.0514	2531.71	0.6744	2375.75	14.1	4C	116
Water	0.014515	1017.34	0.67359	2003.25	69.2	4C	114

Table 4. The sPC-SAFT parameters of relevant compounds

Comp.	m (-)	σ (Å)	ε/k (K)	$\varepsilon^{A_i B_i}/k$ (K)	$\kappa^{A_i B_i}$ (-)	Scheme (-)	Ref.
Methane	1.0	3.7039	150.03	-	-	-	14
Ethane	1.6069	3.5206	191.42	-	-	-	14
n-Hexane	3.0576	3.7983	236.77	-	-	-	14
Benzene	2.4653	3.6478	287.35	-	-	-	14
Toluene	2.8149	3.7169	285.69	-	-	-	14
Methanol (#1)	1.5255	3.23	188.9	2899.50	0.035176	2B	15
Methanol (#2)	1.8824	3.002	181.77	2738.03	0.0547	2B	117
MEG (#1)	1.90878	3.5914	325.23	2080.03	0.0235	4C	118
MEG (#2)	2.3535	3.3186	280.29	2006.60	0.04539	4C	119
Water (#1)	1.5	2.6273	180.3	1804.22	0.18	4C	120
Water (#2)	2.0	2.3449	171.67	1704.06	0.1596	4C	106

Table 5. %AAD of vapor pressure and liquid density against the data from DIPPR and NIST*

Comp.	Tr range	Vapor pressure			Liquid density		
		CPA	sPC-SAFT (#1)	sPC-SAFT (#2)	CPA	sPC-SAFT (#1)	sPC-SAFT (#2)
Water	0.5-0.95	0.72 (0.73)	0.87 (0.96)	1.39 (1.48)	1.19 (1.10)	3.40 (3.36)	2.20 (2.12)
Methanol	0.5-0.95	1.18 (0.75)	1.72 (1.60)	1.32 (1.23)	0.53 (0.61)	0.65 (0.73)	0.36 (0.50)
MEG	0.5-0.65	1.45 (1.65)	0.35 (0.58)	1.28 (1.07)	0.65 (-)	1.76 (-)	1.76 (-)

* The values inside the parentheses is the %AAD calculated based on the data from NIST.¹¹¹




Figure 1. Vapor pressure of water versus temperature ($1/T$) from the CPA and sPC-SAFT EOS. The experimental data are from NIST.¹¹¹

4.2 Binary Vapor-Liquid Equilibria (VLE)

The modeling results of selected binary mixtures are presented in Table 6. The prediction result of each system is given inside parentheses after the correction one, which needs the binary interaction parameter k_{ij} . The average %AADs of each model are also given in the last row of the table for the correlation results. It can be concluded that the three models show quite similar performance in

correlating data, even though they predict different results in some cases. It is interesting to notice that negative values of k_{ij} are needed for both models to correlate the systems of water-methanol/MEG.

Table 6. Binary interaction parameters and both the correlated and predicted %AAD Pressure*

Np	T (K)	System	$k_{ij}/\kappa^{\text{cross}}$ & %AAD Pressure					
			CPA		sPC-SAFT (#1)		sPC-SAFT (#2)	
287	274.19- 444.26	Methane Water	0.7988 $-236.5/T$	6.3(42)	0.2967 $-89.61/T$	11(37)	0.2306 $-92.62/T$	9.3(>100)
165	274.19- 344.26	Methane Water	0.7988 $-236.5/T$	7.0(33)	0.3768 $-114.0/T$	5.6(30)	0.2846 $-109.2/T$	6.0(>100)
97	220.00- 338.20	Methane Methanol	0.053	12(27)	0.041	8.0(39)	0.01	11(19)
85	298.15- 398.15	Methane MEG	0.18	8.0(56)	0.070	5.4(49)	0.064	7.2(47)
153	274.26- 444.26	Ethane Water	0.5437 $-143.3/T$	27(64)	0.2224 $-52.42/T$	27(75)	0.1773 $-53.97/T$	27(55)
33	283.2- 303.20	Ethane Methanol	0.028	6.1(16)	0.031	6.1(24)	0.020	5.9(19)
162	293.15- 333.15	Hexane Methanol	0.01	4.3(6.4)	0.027	4.9(12)	0.021	4.4(11)
128	298.15- 363.15	Benzene Methanol	0.0068	3.2(3.7)	0.0093	5.9(6.6)	0.011	5.2(6.3)
128	298.15- 363.15	Benzene Methanol	0.02/0.01	1.9(3.7)	0.045/0.11	1.4(6.6)	0.04/0.11	1.3(6.3)
78	313.15- 318.15	Toluene Methanol	-0.01	9(10)	-0.0067	15(15)	-0.0013	13(13)
78	313.15- 318.15	Toluene Methanol	0.034/0.029	3.3(10)	0.045/0.13	3.0(15)	0.040/0.13	3.0(13)
130	298.05- 523.15	Water Methanol	-0.09	2.7(13)	-0.059	2.8(14)	-0.066	2.5(19)
40	343.15- 363.15	Water MEG	-0.115	2.8(26)	-0.05	2.5(20)	-0.057	3.5(27)
average %AAD (only for correlation)				7.2		7.6		7.6

* The value inside the parenthesis is from prediction, i.e. $k_{ij} = 0$ and $\kappa^{\text{cross}} = 0$; Np is number of data point.

Figure 2 presents the correlated results of the solubility of methane in water using CPA and sPC-SAFT EOS, which clearly shows similar performance from these models. Since the gas hydrate formation usually occurs below 310K, we developed also a temperature dependent k_{ij} by correlating the data below 350K only. These two k_{ij} sets are compared in Figure 3, together with a constant k_{ij} , which are calculated from the correlations in the first row of Table 6 at temperature 313.45K. It can be seen from Figure 3 that a temperature dependent k_{ij} is necessary for this binary mixture, while the two temperature dependencies show similar performance. We have previously observed that a temperature independent k_{ij} works reasonably for the solubility of methane in methanol or MEG.¹⁰⁷




Figure 2. The correlated solubility of methane in water from the CPA and sPC-SAFT EOS. The binary interaction parameters are shown in Table 6. The experimental data are from Frost et al.,²⁵ Lekvam et al.,²⁷ Wang et al.,²⁸ Gao et al.²⁹ and Culberson et al.³⁰




Figure 3. The solubility of methane in water from the sPC-SAFT EOS with three different k_{ij} approaches. The experimental data are from Frost et al.,²⁵ Lekvam et al.,²⁷ Wang et al.,²⁸ Gao et al.²⁹ and Culberson et al.³⁰

The CPA and PC-SAFT EOS are not able to satisfactorily correlate the solubility of ethane in water, as seen in Table 6. A typical example is given in Figure 4 for three temperatures. The results at 293K are reasonable, while the results at 310K and 344K are not satisfactory, if compared with Figure 2, especially for the sPC-SAFT EOS. This may appear, at a first glance, rather surprising. It should be pointed out that the binary interaction parameter is tuned to the total pressure for all the data together, and it might be possible to correlate the data with more sophisticated temperature dependent binary interaction parameters, which might lead to worse predictive capabilities. The experimental solubilities of methane and ethane in water are compared in Figure 5. It can be seen that the pressure has little

impact on the solubility of ethane in water above 50bar, while the solubility of methane increases gradually. There are not many data available for the solubility of ethane in water under high pressures. This suggests the need for more experimental measurements of this system under high pressures, which will help us understanding the different behaviors of methane and ethane in water.




Figure 4. The solubility of ethane in water from the CPA and sPC-SAFT EOS. The binary interaction parameters are shown in Table 6. The experimental data are from Wang et al.,²⁸ Culberson et al.⁴¹ and Mohammadi et al.⁴²




Figure 5. Comparison of experimental solubilities of methane and ethane in water at two temperatures. The experimental data are from Culberson et al.^{30,41}

Physically the interaction between methanol and benzene is stronger than that of methanol and n-hexane, because we do not see liquid-liquid phase split of methanol and benzene (as shown in Figure 6) while we have LLE of methanol and n-hexane under some circumstances (see below e.g. Figure 7). It can be seen from Table 6 that an extra adjustable parameter for the cross association volume can slightly reduce the %AAD for the binary of methanol and benzene. With or without the cross association volume, however, it can be seen from Figure 6 that the modeling results are quite similar. In this sense, we may consider that solvation might not be necessary for modeling this binary.




Figure 6. VLE of methanol-benzene from the CPA EOS with or without solvation. The binary interaction parameters are given in Table 6. The experimental data are from Toghiani et al.⁷⁰ and Scatchard et al.⁷¹

4.3 Binary Liquid-Liquid Equilibria (LLE)

It is well-established that it is difficult to predict the LLE of binary polar + hydrocarbons systems. The results of some representative binary mixtures with correlated binary interaction parameters are presented in Table 7. Two adjustable binary interaction parameters are used when the hydrocarbon is aromatic. The %AAD corresponds to compounds in the order listed in Table 7, i.e. the first row is for the deviation of the solubility of the hydrocarbon in the aqueous phase, and the second row is for the deviation of the solubility of the polar compound in the organic phase. The average %AADs for each model and each phase are given in the last row of the table. Apparently the correlations of these binary mixtures are not as good as what have been seen in the VLE systems in Table 6. The results, however, can still be considered satisfactory. Again the CPA and sPC-SAFT EOS show quite similar performance in an overall point of view, while the first parameter set #1 of the sPC-SAFT EOS gives a slightly larger overall deviation.

Table 7. LLE of polar-hydrocarbon systems with the CPA and sPC-SAFT models

Np	Temp. (K)	System	$\frac{k_{ij}}{\kappa^{cross}}$		%AAD (mutual solubility)			
			CPA	sPC-SAFT (#1)	sPC-SAFT (#2)			
23	270.0-490.0	n-Hexane	0.0355	35	0.0488	49	0.0021	46
		Water		9.9		13		6.1
6	278.15-303.15	n-Hexane	0.01	47	0.0272	43	0.022	42
		Methanol		12		33		12
5	307.95-330.35	n-Hexane	0.059	5.9	0.043	9.7	0.040	3.7
		MEG		11		19		4.2
15	279.15-453.15	Benzene	0.0355	23	0.058	13	0.023	11
		Water	0.079	8.9	0.15	16	0.15	17
6	279.2-342.1	Benzene	0.049	14	0.020	9.1	0.029	9.4
		MEG	0.0393	3.8	0.043	9.6	0.080	12
16	273.15-458.15	Toluene	0.019	22	0.049	16	0.014	14
		Water	0.071	4.7	0.145	17	0.145	18
10	279.1-361.0	Toluene	0.049	11	0.026	3.4	0.030	4.4
		MEG	0.039	4.3	0.051	6.1	0.083	8.6
average %AAD				15		18		15

The LLE data of methanol-n-hexane have been reported by many groups.⁴⁸⁻⁵⁴ Figure 7 presents three data sets from different sources and the modeling results with the CPA and sPC-SAFT EOS. The data set 1 was measured under pressure 14bar,⁵² and the other two sets are at atmospheric pressure.^{51,53} It can be seen that the data are consistent with each other on the methanol rich polar phase, while there are significant differences on the n-hexane rich phase, especially when the methanol concentration is lower than 0.15. This suggests a need for more systematic evaluation of the experimental data. It can also be seen that the CPA EOS presents larger deviations on the methanol rich side close to the cloud point, while the first parameter set #1 of the sPC-SAFT EOS has difficulties in modeling the methanol lean branch.




Figure 7. LLE of methanol-n-hexane from the CPA and sPC-SAFT EOS. The experimental data are from Blanco et al.,⁵² Hradetzky et al.⁵¹ and Matsuda et al.⁵³ The binary interaction parameters are shown in Table 7.

Figure 8 presents the LLE of the binary mixture of water and benzene. The CPA EOS performs better in modeling the solubility of water in the organic phase, while PC-SAFT shows smaller deviations in modeling the solubility of benzene in the aqueous phase, especially at low to moderate temperatures with the currently investigated parameters. It should be pointed out that neither model can have correctly capture the minimum in the solubility of benzene, and the sPC-SAFT EOS shows larger deviations at higher temperatures.




Figure 8. LLE of water-benzene from the CPA and sPC-SAFT EOS. The binary interaction parameters are shown in Table 7. The experimental data are from Jou et al.⁶⁶

The models show similar behavior for the LLE of MEG-hydrocarbon, and one typical example for MEG-toluene is given in Figure 9. It shows that toluene is more soluble in MEG than MEG in toluene. With the given parameters, the CPA EOS performs better in modeling the solubility of MEG in toluene, while the sPC-SAFT EOS seems to be better for the other phase. The overall performance of the two models is again considered to be rather similar and quite satisfactory.




Figure 9. LLE of MEG-toluene from the CPA and sPC-SAFT EOS. The binary interaction parameters are shown in Table 7. The experimental data are from Folas et al.⁷⁷

4.4 Multicomponent fluid phase equilibria

When the pure component parameters and binary interaction parameters are available, we can investigate the predictive capabilities of the models, i.e. the predictions of multicomponent multiphase equilibria using solely binary parameters estimated from binary data. The predicted results of the VLE of water-methanol/MEG-methane are given in Table 8. Apparently the two models show quite similar performance with all parameter sets.

Table 8. %AAD of composition in the VLE of ternary mixtures

T (K) (Np)	Systems	Aqueous phase			Vapor phase			
		CPA	sPC-SAFT (#1)	sPC-SAFT (#2)	CPA	sPC-SAFT (#1)	sPC-SAFT (#2)	
280.25- 313.45 (Np=105)	Methane	35	26	32	0.009	0.011	0.007	
	Water	0.4	0.3	0.3	22	19	15	
	Methanol	0.7	0.5	0.7	2.8	6.6	1.2	
	k_{12}	the first case in Table 6						
	Methane	35	30	35	0.009	0.011	0.007	
	Water	0.4	0.3	0.4	22	18	14	
	Methanol	0.7	0.6	0.7	2.8	6.7	1.2	
	k_{12}	the second case in Table 6						
	Methane	18	14	18	0.009	0.0112	0.007	
	Water	0.2	0.2	0.2	25	20	17	
	Methanol	0.5	0.4	0.5	3.1	6.5	1.2	
	k_{12}	0.0443	0.0108	-0.0649				
	283.2- 303.2 (Np=92)	Methane	24	30	29	-	-	-
		Water	0.1	0.1	0.1	-	-	-
MEG		0.1	0.1	0.1	-	-	-	
k_{12}		the first case in Table 6						
Methane		24	33	31	-	-	-	
Water		0.1	0.2	0.1	-	-	-	
MEG		0.1	0.1	0.1	-	-	-	
k_{12}		second case in Table 6						
Methane		15	20	18	-	-	-	
Water		0.1	0.1	0.1	-	-	-	
MEG	0.1	0.1	0.1	-	-	-		
k_{12}	0.0443	0.0108	-0.0649	-	-	-		
average %AAD		8.6	8.7	9.2	8.6	8.5	5.5	

Three options of the binary interaction parameter between water and methane, i.e. a temperature dependent k_{ij} covering data up to 450K, a temperature dependent k_{ij} covering data only up to 350K and a constant k_{ij} , have been investigated. The two temperature dependent k_{ij} sets show similar performance as seen for this binary mixture. However, we have surprisingly found that a constant k_{ij} significantly improves the predictions of the solubility of methane in the aqueous phase without deteriorating the other phase. It is important to recall that a constant k_{ij} does not describe the binary mixture of water and methane well, as discussed in Figure 3 above. This might be a coincidence, as we see from Figure 10 that the constant k_{ij} underpredicts and overpredicts the solubility of methane in the liquid phase at 283.2K and 303.2K, respectively. Therefore, a temperature dependent k_{ij} is still recommended if wide temperature applications are concerned.




Figure 10. The solubility of methane in the solution of methanol and water from the sPC-SAFT EOS (set #2) with different k_{ij} approaches. The experimental data are from Wang et al.²⁸

The deviations of the predicted LLE results of water-methanol/MEG-n-hexane/benzene/toluene are presented in Table 9. The deviations are given corresponding to the component list as in Table 8. The average %AADs for each model and each phase are added in the last row of the table as well. The models again show comparable performances, even though a slightly larger deviation is seen for the first parameter set #1 of the sPC-SAFT EOS.

Table 9. %AAD of composition in the LLE of ternary mixtures

T (K)	Systems	Aqueous phase			Organic phase		
		CPA	sPC-SAFT (#1)	sPC-SAFT (#2)	CPA	sPC-SAFT (#1)	sPC-SAFT (#2)
293.15-	n-Hexane	29	22	22	2.0	2.1	1.6
318.15	Water	8.0	4.5	5.8	69	75	68
Np=81	Methanol	1.9	1.2	1.4	56	30	35
283.15-	n-Hexane	37	23	20	0.017	0.019	0.010
323.15	Water	0.012	0.009	0.015	43	50	31
Np=24	MEG	0.029	0.021	0.031	17	42	23
	Benzene	31	40	39	6.5	6.3	5.7
	Water	7.9	9.2	9.8	33	35	34
	Methanol	4.8	6.3	6.4	42	44	44
		Solvation is used for methanol and benzene					
293.15-	Benzene	28	28	30	6.9	9.1	7.6
333.15	Water	6.2	4.2	5.2	32	37	33
Np=62	Methanol	3.6	2.3	2.7	44	54	48
		No solvation is used for methanol and benzene					
298.2-	Benzene	19	27	21	0.012	0.013	0.016
323.2	Water	0.2	0.3	0.2	8.0	6.3	5.2
Np=10	MEG	0.2	0.2	0.2	14	7.9	19
298.2-	Toluene	18	15	15	0.006	0.016	0.014
323.2	Water	0.044	0.026	0.025	3.9	4.6	2.6
Np=8	MEG	0.11	0.07	0.066	7.6	9.1	12
	average %AAD	11	10	10	21	23	21

The performances of the models are generally considered to be satisfactory, but it can be seen that deviations of the solubilities of water and methanol in the n-hexane rich organic phase are relatively large. The modeling results of this system at two conditions are presented in Figures 11 and 12, for which the data are from different sources. The models give very satisfactory predictions for all the components in Figure 11, while they poorly predict the solubilities of water and methanol in the organic phase in Figure 12. It can be recognized that the solubility of water shown in Figure 12 is almost one order magnitude higher than those shown in Figure 11, and it is closer to the solubility of water in benzene, as given in Figure 13. This suggests once again the need for further data validation, e.g. for making new and systematic measurements for this system under these conditions.




Figure 11. LLE of water-methanol-n-hexane at 293.2K from the CPA EOS. The experimental data are from Skrzecz et al.⁸⁷ The binary interaction parameters are shown in Tables 6 and 7.




Figure 12. LLE of water-methanol-n-hexane at 318.15K from the CPA EOS. The experimental data are from Liu et al.⁸⁸ The interaction parameters are shown in Tables 6 and 7.

As discussed above (e.g. in Figure 6), solvation between methanol and benzene has insignificant effects on the VLE of the binary mixture. It is shown in Figure 13 that the solvation can slightly move the solubility lines, i.e. bringing more methanol into the organic phase and less benzene into the aqueous phase. As indicated by the %AAD given in Table 9 solvation between methanol and benzene has a small impact on modeling the system of water-methanol-benzene as well from an overall point of view. Therefore, as an alternative, solvation could be used for water and benzene only. As shown in Figure 6, more measurements for the binary methanol and benzene at low concentration range of methanol may be useful and may actually lead to somewhat different conclusions.






Figure 13. LLE of water-methanol-benzene from the CPA EOS with or without solvation between benzene and methanol at 318.15K. The interaction parameters are shown in Tables 6 and 7.

The performance of these models and parameters are further tested in predicting the fluid phase equilibria of quaternary mixtures. The results for two mixtures are presented in Figures 14 and 15 and the deviations are listed in Table 10. Both systems contain water, MEG and methane. The last component is n-hexane in the first mixture and toluene in the second one. Solvation is used for both MEG-toluene and water-toluene. The two models show similar performance for these systems, especially for the vapor phase and the aqueous phase, and the only exception might be an underprediction of the solubility of MEG in the organic phase of the second quaternary mixture from the CPA EOS. The modeling results, in general terms, are satisfactory, except both models underpredict the solubility of n-hexane in the aqueous phase, as shown in Figure 14. Both models overpredict the solubility of toluene in the aqueous phase at 263.15K in Figure 15, in which case the qualitative behavior is opposite against the data. This suggests that more measurements are needed to further verify the data and validate the model.

Table 10. Vapor-liquid-liquid equilibria of two quaternary mixtures

Temperature (K)	Comp.	%AAD of mole fraction					
		Vapor		Organic		Aqueous	
		CPA	sPC-SAFT (#2)	CPA	sPC-SAFT (#2)	CPA	sPC-SAFT (#2)
263.15- 283.15	H2O	-	-	38	14	0.06	0.05
	MEG	-	-	52	34	0.49	0.49
	C1	0.04	0.03	1.4	5.3	33	35
	nC6	9.8	4.2	2.4	4.5	94	88
263.15- 323.15	H2O	44	38	18	13	0.06	0.06
	MEG	-	-	90	23	0.12	0.02
	C1	0.02	0.03	13	17	11	14
	Toluene	3.0	10	2.9	3.6	1121	2152
average %AAD*		11	10	27	14	20	20

* The average %AADs of the aqueous phase do not count the solubility of toluene, since one point largely overpredicted.








Figure 14. VLLE of water-MEG-methane-n-hexane at 263.15K and 283.15K. Solid and dash-dot lines are modeling results from the CPA and sPC-SAFT EOS, respectively. The parameter set #2 is used for the sPC-SAFT EOS.






Figure 15. VLLE of water-MEG-methane-toluene from 263.15K to 323.15K. Solid and dash-dot lines are modeling results from the CPA and sPC-SAFT EOS, respectively. The parameter set #2 is used for the sPC-SAFT EOS.

5. Conclusion

In this work, a systematic and fair comparison has been conducted for the CPA and sPC-SAFT EOS by applying them into modeling the fluid phase equilibria of gas hydrate related systems. The results show that these two models perform quite similarly in both correlating binary mixtures and predicting multicomponent multiphase equilibria, and the modeling results are satisfactory in most cases. Experimental data are ultimate criteria in process and product design, and they are also necessary for obtaining model parameters and validating model's performance. We show in this work that thermodynamic models can in reverse help in suggesting experimental measurements. For example, more measurements are needed for ethane-water system under high pressures, for methanol-n-hexane LLE data at low concentration of methanol, for water-methanol-n-hexane LLE data and for water-MEG-methane-toluene at low temperatures.

Author Information

Corresponding Author

* Tel.: 045-45252859. Fax: 045-45882258. Email: gk@kt.dtu.dk

List of Symbols

EOS = Equation(s) of State

CPA = Cubic Plus Association

SAFT = Statistical Associating Fluid Theory

PC-SAFT = Perturbed-Chain Statistical Associating Fluid Theory

sPC-SAFT = Simplified Perturbed-Chain Statistical Associating Fluid Theory

%AAD = Percent Average Absolute Deviation

T = Temperature

P = Pressure

V_m = Molar volume

R = Ideal Gas constant

Tr = Reduced temperature (T/critical temperature)

a^r = Reduced residual Helmholtz free energy

a^{HS} = Reduced residual Helmholtz free energy from hard-sphere

a^{HC} = Reduced residual Helmholtz free energy from chain formation

a^{disp} = Reduced residual Helmholtz free energy from dispersion

a^{assoc} = Reduced residual Helmholtz free energy from association (hydrogen-bonding)

b = The co-volume parameter

$a(T)$ = The van der Waals energy parameter

g = Radial distribution function

m = Segment number

x_i = Molar fraction of component i

A_i = Association site type A of component i

X^{A_i} = Free site fraction of molecules i not bonded at site A

Greeks Letters

ρ = Molar density

σ = Segment diameter (T independent)

ε = Segment energy

$\Delta^{A_i B_j}$ = Association strength

$\varepsilon^{A_i B_j}$ = Association energy between sites A_i and B_j

$\kappa^{A_i B_j}$ = Association volume between sites A_i and B_j

κ^{cross} = Adjustable cross association volume

Reference

- (1) Mokbel, I.; Lindemann, C.; Duchet-Suchaux, P.; Jose, J. Liquid-liquid equilibria of binary and ternary systems involving monoethyleneglycol, water, n-alkanes at three temperatures: 283.15, 303.15 and 333.15K. *Fuel*. **2016**, *163*, 17-24.
- (2) Hydrates. <http://petrowiki.org/Hydrates> (accessed March 21, **2017**).
- (3) Jonassen, A. H. Distribution of hydrate inhibitors in processing of gas-condensate. NTNU-Trondheim, Norway. **2013**.
- (4) Kontogeorgis, G. M.; Voutsas, E. C.; Yakoumis, I. V.; Tassios, D. P. An equation of state for associating fluids. *Ind. Chem. Eng. Res.* **1996**, *35*, 4310-4318.
- (5) Kontogeorgis, G. M.; Folas, G. K. Thermodynamic models for industrial applications – from classical and advanced mixing rules to association theories. John Wiley and Sons: New York, **2010**.
- (6) Riaz, M. Distribution of Complex Chemicals in Oil-Water Systems. PhD Thesis, Technical University of Denmark, Kgs. Lyngby, Denmark, **2011**.
- (7) Frost, M. Measurement and Modelling of Phase Equilibrium of Oil-Water-Polar Chemicals. PhD Thesis, Technical University of Denmark, Kgs. Lyngby, Denmark, **2014**.
- (8) Wertheim, M. S. Fluids with highly directional attractive forces. I. Statistical thermodynamics. *J. Stat. Phys.* **1984**, *35*, 19-34.
- (9) Wertheim, M. S. Fluids with highly directional attractive forces. II. Thermodynamic perturbation theory and integral equations. *J. Stat. Phys.* **1984**, *35*, 35-47.
- (10) Wertheim, M. S. Fluids with highly directional attractive forces. III. Multiple attraction sites. *J. Stat. Phys.* **1986**, *42*, 459-476.
- (11) Wertheim, M. S. Fluids with highly directional attractive forces. IV. Equilibrium polymerization. *J. Stat. Phys.* **1986**, *42*, 477-492.
- (12) Chapman, W. G.; Gubbins, K. E.; Jackson, G.; Radosz, M. SAFT: Equation-of-state solution model for associating fluids. *Fluid Phase Equilib.* **1989**, *52*, 31-38.
- (13) Chapman, W. G.; Gubbins, K. E.; Jackson, G.; Radosz, M. New reference equation of state for associating liquids. *Ind. Eng. Chem. Res.* **1990**, *29*, 1709-1721.
- (14) Gross, J.; Sadowski, G. Perturbed-chain SAFT: an equation of state based on a perturbation theory for chain molecules. *Ind. Eng. Chem. Res.* **2001**, *40*, 1244-1260.
- (15) Gross, J.; Sadowski, G. Application of the perturbed-chain SAFT equation of state to associating systems. *Ind. Eng. Chem. Res.* **2002**, *41*, 5510-5515.
- (16) von Solms, N.; Michelsen, M. L.; Kontogeorgis, G. M. Computational and physical performance of a modified PC-SAFT equation of state for highly asymmetric and associating mixtures. *Ind. Eng. Chem. Res.* **2003**, *42*, 1098-1105.
- (17) Galindo, A.; Whitehead, P. J.; Jackson, G.; Burgess, A. N. Predicting the high-pressure phase equilibria of water + n-alkanes using a simplified SAFT theory with transferable intermolecular interaction parameters. *J. Phys. Chem.* **1996**, *100*, 6781-6792.
- (18) Voutsas, E. C.; Boulougouris, G. C.; Economou, I. G.; Tassios, D. P. Water/hydrocarbon phase equilibria using the thermodynamic perturbation theory. *Ind. Eng. Chem. Res.* **2000**, *39*, 797-804.
- (19) Vega, L. F.; Llovel, F.; Blas, F. J. Capturing the solubility minima of n-alkanes in water by soft-SAFT. *J. Phys. Chem. B*, **2009**, *113*, 7621-7630.
- (20) Dufal, S.; Galindo, A.; Jackson, G.; Haslam, A. J. Modelling the effect of methanol, glycol inhibitors and electrolytes on the equilibrium stability of hydrates with the SAFT-VR approach. *Mol. Phys.* **2012**, *110*, 1223-1240.

- (21) Papaioannou, V.; Calado, F.; Lafitte, T.; Dufal, S.; Sadeqzadeh, M.; Jackson, G.; Adjiman, C. S.; Galindo, A. Application of the SAFT- γ Mie group contribution equation of state to fluids of relevance to the oil and gas industry. *Fluid Phase Equilib.* **2016**, *416*, 104-119.
- (22) Ahmed, S.; Ferrando, N.; de Hemptinne, J. C.; Simonin, J. P. Bernard, O.; Baudouin, O. A new PC-SAFT model for pure water, water-hydrocarbons, and water-oxygenates systems and subsequent modeling of VLE, VLLE, and LLE. *J. Chem. Eng. Data*, **2016**, *61*, 4178-4190.
- (23) Tybjerg, P. C. V.; Kontogeorgis, G. M.; Michelsen, M. L.; Stenby, E. H. Phase equilibria modeling of methanol-containing systems with the CPA and sPC-SAFT equations of state. *Fluid Phase Equilib.* **2010**, *288*, 128-138.
- (24) Barker, J. A.; Henderson, D. Perturbation theory and equation of state for fluids: the square-well potential. *J. Chem. Phys.* **1967**, *47*, 2856-2861.
- (25) Frost, M.; Karakatsani, E.; von Solms, N.; Richon, D.; Kontogeorgis, G. M. Vapor-liquid equilibrium of methane with water and methanol. Measurements and modeling. *J. Chem. Eng. Data*. **2014**, *59*, 961-967.
- (26) Kiepe, J.; Horstmann, S.; Fischer K.; Gmehling, J. Experimental determination and prediction of gas solubility data for methane + water solutions containing different monovalent electrolytes. *Ind. Eng. Chem. Res.* **2003**, *42*, 5392-5398.
- (27) Lekvam, K.; Bishnoi, P. R. Dissolution of methane in water at low temperatures and intermediate pressures. *Fluid Phase Equilib.* **1997**, *131*, 297-309.
- (28) Wang, L. K.; Chen, G. J.; Han, G. H.; Guo, X. Q.; Guo, T. M. Experimental study on the solubility of natural gas components in water with or without hydrate inhibitor. *Fluid Phase Equilib.* **2003**, *207*, 143-154.
- (29) Gao, J.; Zheng, D. Q.; Guo, T. M. Solubilities of methane, nitrogen, carbon dioxide, and a natural gas mixture in aqueous sodium bicarbonate solutions under high pressure and elevated temperature. *J. Chem. Eng. Data*. **1997**, *42*, 69-73.
- (30) Culberson, O. L.; McKetta Jr., J. J. Phase equilibria in hydrocarbon-water systems III - The solubility of methane in water at pressures to 10,000 PSIA. *Petro. Trans. AIME.* **1951**, *192*, 223-226.
- (31) Tabasinejad, F.; Gordon Moore, R.; Mehta, S. A.; Van Fraassen, K. C.; Barzin, Y. Water solubility in supercritical methane, nitrogen, and carbon dioxide: measurement and modeling from 422 to 483 K and pressures from 3.6 to 134 MPa. *Ind. Eng. Chem. Res.* **2011**, *50*, 4029-4041.
- (32) Hong, J. H.; Malone, P. V.; Jett, M. D.; Kobayashi, R. The measurement and interpretation of the fluid-phase equilibria of a normal fluid in a hydrogen bonding solvent: the methane-methanol system. *Fluid Phase Equilib.* **1987**, *38*, 83-96.
- (33) Yarym-Agaev, N. L.; Sinyavskaya, R. P.; Koliushko, I. I. Phase equilibrium in the binary systems water-methane and methanol-methane under high pressure. *Zh. Prikl. Khim.* **1985**, *58*, 165-168.
- (34) Francesconi, A. Z.; Lentz, H.; Franck, E. U. Phase equilibria and PVT data for the methane-methanol system to 300 MPa and 240 °C. *J. Phys. Chem.* **1981**, *85*, 3303-3307.
- (35) Hong, J. H.; Malone, P. V.; Jett, M. D.; Kobayashi, R. The measurement and interpretation of the fluid-phase equilibria of a normal fluid in a hydrogen bonding solvent: the methane-methanol system. *Fluid Phase Equilib.* **1987**, *38*, 83-96.
- (36) Suzuki, K.; Sue, H.; Itou, M.; Smith, R. L.; Inomata, H.; Arai, K.; Saito, S. Isothermal vapor-liquid equilibrium data for binary systems at high pressures: carbon dioxide-methanol, carbon dioxide-ethanol, carbon dioxide-1-propanol, methane-ethanol, methane-1-propanol, ethane-ethanol, and ethane-1-propanol systems. *J. Chem. Eng. Data*. **1990**, *35*, 63-66.

- (37) Frolich P. K.; Tauch E. J.; Hogan J. J.; Peer A. A. Solubilities of gases in liquids at high pressure. *Ind. Eng. Chem.* **1931**, *23*, 548-550.
- (38) Brunner, E.; Hültenschmidt, W. Fluid mixtures at high pressures VIII. Isothermal phase equilibria in the binary mixtures: (ethanol+hydrogen or methane or ethane). *J. Chem. Thermodynamics.* **1990**, *22*, 73-84.
- (39) Jou, F. Y.; Otto, F. D.; Mather, A. E. Solubility of methane in glycols at elevated pressures. *Can. J. Chem. Eng.* **1994**, *72*, 130-133
- (40) Zheng, D. Q.; Ma, W.D.; Wei, R.; Guo, T. M. Solubility study of methane, carbon dioxide and nitrogen in ethylene glycol at elevated temperatures and pressures. *Fluid Phase Equilib.* **1999**, *155*, 277-286.
- (41) Culberson, O. L.; McKetta Jr., J. J. Phase equilibria in hydrocarbon-water systems II - The solubility of ethane in water at pressures to 10,000 psi. *Petro. Trans. AIME.* **1950**, *189*, 319-322.
- (42) Mohammadi, A. H.; Chapoy, A.; Tohidi, B.; Richon, D. Measurements and thermodynamic modeling of vapor-liquid equilibria in ethane-water systems from 274.26 to 343.08 K. *Ind. Eng. Chem. Res.* **2004**, *43*, 5418-5424.
- (43) King Jr., A. D.; Coan, C. R. Solubility of water in compressed carbon dioxide, nitrous oxide, and ethane. Evidence for hydration of carbon dioxide and nitrous oxide in the gas phase. *J. Am. Chem. Soc.* **1971**, *93*, 1857-1862.
- (44) Chapoy, A.; Coquelet, C.; Richon, D. Measurement of the water solubility in the gas phase of the ethane + water binary system near hydrate formatting conditions. *J. Chem. Eng. Data*, **2003**, *48*, 957-966.
- (45) Ohgaki, K.; Sano, F.; Katayama, T. Isothermal vapor-liquid equilibrium data for binary systems containing ethane at high pressures. *J. Chem. Eng. Data*, **1976**, *21*, 55-58.
- (46) Barrufet, M. A.; Liu, K.; Rahman, S.; Wu, C. Simultaneous vapor-liquid-liquid equilibria and phase molar densities of a quaternary system of propane + pentane + octane + water, *J. Chem. Eng. Data.* **1996**, *41*, 918-922.
- (47) Tsonopoulos, C.; Wilson, G. M. High-temperature mutual solubilities of hydrocarbons and water. Part I. Benzene, cyclohexane and n-hexane. *AIChE J.* 1983, *29*, 990.
- (48) Vapor-liquid equilibrium data of hexane + methanol from Dortmund data bank. <http://www.ddbst.com/en/EED/VLE/VLE%20Hexane%3BMethanol.php> (accessed March 12, **2017**).
- (49) Ferguson J. B. The system methyl alcohol-n-hexane at 45 degrees. *J. Phys. Chem.* **1932**, *36*, 1123-1128.
- (50) Góral, M.; Oracz, P.; Warycha, S. Vapour-liquid equilibria XIV. The ternary system cyclohexane-methanol-hexane at 293.15K. *Fluid Phase Equilib.* **2000**, *169*, 85-99.
- (51) Hradetzky, G.; Lempe, D. A. Phase equilibria in binary and higher systems methanol + hydrocarbons(s): Part I. Experimental determination of liquid-liquid equilibrium data and their representation using the NRTL equation. *Fluid Phase Equilib.* **1991**, *69*, 285-301.
- (52) Blanco, A. M.; Ortega, J. Experimental study of miscibility, density and isobaric vapor-liquid equilibrium values for mixtures of methanol in hydrocarbons (C5, C6). *Fluid Phase Equilib.* **1996**, *122*, 207-222.
- (53) Matsuda, H.; Kurihara, K.; Ochi, K.; Kojima, K. Prediction of liquid-liquid equilibria at high pressure for binary systems using EOS-GE models: methanol+hydrocarbon systems. *Fluid Phase Equilib.* **2002**, *203*, 269-284.

- (54) Hölscher, I. F.; Schneider, G. M.; Ott, J. B. Liquid-liquid phase equilibria of binary mixtures of methanol with hexane, nonane and decane at pressures up to 160 MPa. *Fluid Phase Equilib.* **1986**, *27*, 153-169.
- (55) Sinor, J. E.; Weber J. H. Vapor-liquid equilibria at atmospheric pressure. Systems containing ethyl alcohol, n-hexane, benzene and methylcyclopentane. *J. Chem. Eng. Data*, **1960**, *5*, 243-247.
- (56) Yuan, K. S.; Ho, J. C. K.; Koshpande, A. K.; Lu, B. C. Y. Vapor-liquid equilibria. *J. Chem. Eng. Data*, **1963**, *8*, 549-559.
- (57) Derawi, S. O.; Kontogeorgis, G. M.; Stenby, E. H.; Haugum, T.; Fredheim, A. O. Liquid-liquid equilibria for glycol hydrocarbons data and correlation. *J. Chem. Eng. Data*. **2002**, *47*, 169-173.
- (58) Shaw, D.G.; Maczynski, A.; Goral, M.; Wisniewska-Goclovska, B.; Skrzecz, A.; Owczarek, I.; Blazej, K.; Haulait-Pirson, M.C.; Hefter, G.T.; Kapuku, F.; Maczynska, Z. IUPAC-NIST Solubility Data Series. 81. Hydrocarbons with water and seawater revised and updated. Part 7. C₈H₁₂–C₈H₁₈ hydrocarbons with water. *J. Phys. Chem. Ref. Data*. **2005**, *34*, 2261-2298.
- (59) Heidman, J. L.; Tsonopoulos, C.; Brady, C. J.; Wilson, G. M. High-temperature mutual solubilities of hydrocarbons and water II: ethylbenzene, ethylcyclohexane, and n-octane. *AIChE J.* **1985**, *31*, 376-384.
- (60) Ott, J. B.; Hölscher, I. F.; Schneider, G. M. (Liquid + liquid) phase equilibria in (methanol + heptane) and (methanol + octane) at pressures from 0.1 to 150 MPa. *J. Chem. Thermodyn.* **1986**, *18*, 815-826.
- (61) Higashiuchi, H.; Sakuragi, Y.; Iwai, Y.; Arai, Y.; Nagatani, M. Measurement and Correlation of Liquid-Liquid Equilibria of Binary and Ternary Systems Containing Methanol and Hydrocarbons. *Fluid Phase Equilib.* **1987**, *36*, 35-47.
- (62) Kurihara, K.; Midorikawa, T.; Hashimoto, T.; Kojima, K.; Ochi, K. Liquid-liquid solubilities for the binary systems of methanol with octane and 2,2,4-trimethylepentane. *J. Chem. Eng. Japan*, **2002**, *35*, 360-364.
- (63) Boublikova, L.; Lu, B. C. Y. Isothermal vapour-liquid equilibria for the ethanol-n-octane system. *J. Chem. Technol. Biotech.* **1969**, *19*, 89-92.
- (64) Janaszewski, B.; Oracz, P.; Goral, M.; Warycha, S. Vapour-liquid equilibria. I. An apparatus for isothermal total vapour pressure measurements: binary mixtures of ethanol and t-butanol with n-hexane, n-heptane and n-octane at 313.15 K. *Fluid Phase Equilib.* **1982**, *9*, 295-310.
- (65) Vapor-liquid equilibrium data of benzene + water from Dortmund data bank. <http://www.ddbst.com/en/EED/VLE/VLE%20Benzene%3BWater.php> (accessed March 12, **2017**).
- (66) Jou, F. Y.; Mather, A. E. Liquid-liquid equilibria for binary mixtures of water + benzene, water + toluene and water + p-xylene from 273 K to 458 K. *J. Chem. Eng. Data*. **2003**, *48*, 750-752.
- (67) Scatchard, G.; Ticknor, L. B. Vapor-liquid equilibrium. IX. The methanol-carbon tetrachloride-benzene system. *J. Am. Chem. Soc.* **1952**, *74*, 3724-3729.
- (68) Vapor-liquid equilibrium data of benzene + methanol from Dortmund data bank. <http://www.ddbst.com/en/EED/VLE/VLE%20Benzene%3BMethanol.php> (accessed March 12, **2017**).
- (69) Miyano Y.; Hayduk, W. Solubilities of isobutane in methanol + benzene, methanol + cyclohexane and benzene + cyclohexane mixed solvent solutions at 298 K and 40-102 kPa. *Fluid Phase Equilib.* **1993**, *38*, 277-268.
- (70) Toghiani, H.; Toghiani, R. K.; Viswanath, D. S. Vapor-liquid equilibria for the methanol-benzene and methanol-thiophene systems. *J. Chem. Eng. Data*, **1994**, *39*, 63-67.
- (71) Scatchard, G.; Wood, S. E.; Mochel, J. M. Vapor-liquid equilibrium. VI. Benzene-methanol mixtures. *J. Am. Chem. Soc.* **1946**, *68*, 1957-1960.

- (72) Nagata, I. Vapor-liquid equilibrium data for the binary systems methanol-benzene and methyl acetate-methanol. *J. Chem. Eng. Data.* **1969**, *14*, 418-420.
- (73) Brown, I.; Smith, F. Liquid-vapour equilibria. IV. The system ethanol + benzene at 45°C. *Austra. J. Chem.* **1954**, *7*, 264-268.
- (74) Zielkiewicz, J.; Oracz, P.; Warycha, S. Total vapour pressure measurements and excess Gibbs energies for the binary systems methanol + ethanol, ethanol + 2-propanol, benzene + cyclohexane, benzene + carbon tetrachloride and benzene + ethanol at 303.15 and 313.15 K. *Fluid Phase Equilib.* **1990**, *58*, 191-209.
- (75) Udovenko, V. V.; Fatkulina, L. G. Vapor pressure of three-component systems. I. The system ethyl alcohol-1, 2-dichloroethane-benzene. *Zh. Fiz. Khim.* **1952**, *26*, 719-730.
- (76) Góral, M.; Skrzecz, A.; Bok, A.; Mączyński, A. Recommended vapor-liquid equilibrium data. Part 3. Binary alkanol-aromatic hydrocarbon systems. *J. Phys. Chem. Ref. Data*, **2004**, *33*, 959-997.
- (77) Folas, G. K.; Kontogeorgis, G. M.; Michelsen, M. L.; Stenby, E. H.; Solbraa, E. Liquid-liquid equilibria for binary and ternary systems containing glycols, aromatic hydrocarbons and water: experimental measurements and modeling with the CPA EoS. *J. Chem. Eng. Data.* **2006**, *51*, 977-983.
- (78) Kretschmer, C. B.; Wiebe, R. Liquid-Vapor Equilibrium of Ethanol – Toluene Solutions. *J. Am. Chem. Soc.* 1949, *71*, 1793-1797.
- (79) Van Ness, H. C.; Soczek, C. A.; Peloquin, G. L.; Machado, R. L. Thermodynamic excess properties of three alcohol-hydrocarbon systems. *J. Chem. Eng. Data.* **1967**, *12*, 217-224.
- (80) Butler, J. A. V.; Thomson, D. W.; MacLennan, W. H. The free energy of the normal aliphatic alcohols in aqueous solution. Part I. The partial vapour pressures of aqueous solutions of methyl, n-propyl, and n-butyl alcohols. Part II. The solubilities of some normal aliphatic alcohols in water. Part III. The theory of binary solutions, and its application to aqueous-alcoholic solutions. *J. Chem. Soc.* **1933**, 674-686.
- (81) Griswold, J.; Wong, S. Y. Phase equilibria of the acetone-methanol-water systems from 100°C into the critical region. *Chem. Eng. Prog. Symp. Ser.* **1952**, *48*, 18-34.
- (82) Kurihara, K.; Minoura, T.; Takeda, K.; Kojima, K. Isothermal vapor-liquid equilibria for methanol + ethanol + water, methanol + water, and ethanol + water. *J. Chem. Eng. Data.* **1995**, *40*, 679-684.
- (83) Ratcliff, G. A.; Chao, K. C. Prediction of thermodynamic properties of polar mixtures by a group solution model. *Can. J. Chem. Eng.* **1969**, *47*, 148-153.
- (84) Chiavone-Filho, O.; Proust, P.; Rasmussen, P. Vapor-liquid equilibria for glycol ether + water systems. *J. Chem. Eng. Data.* **1993**, *38*, 128-131.
- (85) Ng, H. J.; Robinson, D. B. Equilibrium phase composition and hydrating conditions in systems containing methanol, light hydrocarbons, carbon dioxide, and hydrogen sulfide. GPA Research Report 66. Gas Processors Association: Tulsa, Okla. **1983**.
- (86) Noda, K.; Sato, K.; Nagatsuka, K.; Ishida, K. Ternary liquid-liquid equilibria for the systems of aqueous methanol solutions and propane or n-butane. *J. Chem. Eng. Japan.* **1975**, *8*, 492-493.
- (87) Skrzecz, A.; Shaw, D.; Maczynski, A. IUPAC-NIST Solubility Data Series. 69. Ternary alcohol-hydrocarbon-water systems. *J. Phys. Chem. Ref. Data.* **1999**, *28*, 983-1235.
- (88) Liu, J.; Qin, Z.; Wang, J. Liquid-liquid equilibria for methanol + water + hexane ternary mixtures. *J. Chem. Eng. Data.* **2002**, *47*, 1243-1245.
- (89) Lindemann, C.; Duchet-Suchaux, P.; Razzouk, A.; Mokbel, I.; Jose, J. Liquid-liquid equilibria of binary and ternary systems methanol/water + n-hexane, + n-octane, + n-dodecane, and + n-

- hexadecane in the temperature range between $T=283.15\text{K}$ and $T=333.15\text{K}$. *J. Chem. Eng. Data.* **2016**, *61*, 2412-2418.
- (90) Letcher, T. M.; Wootton, S.; Shuttleworth, B.; Heyward, C. Phase equilibria for (n-heptane+ water+ an alcohol) at 298.2 K. *J. Chem. Thermodyn.* **1986**, *18*, 1037-1042.
- (91) Triday, J. O. Liquid-liquid equilibria for the system benzene-water-methanol, *J. Chem. Eng. Data.* **1984**, *29*, 321-324.
- (92) de Doz, M. B. G.; Bonatti, C. M.; Sólamo, H. N. Liquid-liquid equilibria of ternary and quaternary systems with two hydrocarbons, an alcohol, and water at 303.15 K System containing 2,2,4-trimethylpentane, toluene, methanol, and water, or 2,2,4-trimethylpentane, toluene, ethanol, and water. *Fluid Phase Equilib.* **2003**, *205*, 53-67.
- (93) Razzouk, A.; Naccoul, R. A.; Mokbel, I.; Duchet-Suchaux, P.; Jose, J.; Rauzy, E.; Berro, C. Liquid-Liquid Equilibria for Monoethylene Glycol + Hexane and 2,2,4-Trimethylpentane, Water + Hexane and 2,2,4-Trimethylpentane, Monoethylene Glycol + Water + Hexane, and Monoethylene Glycol + Water + 2,2,4-Trimethylpentane in the Temperature Range between $T = 283.15\text{ K}$ and $T = 323.15\text{ K}$. *J. Chem. Eng. Data.* **2010**, *55*, 1468-1472.
- (94) Riaz, M.; Yussuf, M. A.; Kontogeorgis, G. M.; Stenby, E. H.; Yan, W.; Solbraa, E. Distribution of MEG and methnaol in well-defined hydrocarbon and water systems: Experimental measurement and modeling using CPA EoS. *Fluid Phase Equilib.* **2013**, *337*, 298-310.
- (95) Horizoe, H.; Tanimoto, T.; Yamamoto, I.; Kano, Y. Phase Equilibrium study for the separation of ethanol-water solution using subcritical and supercritical hydrocarbon solvent extraction. *Fluid Phase Equilib.* **1993**, *84*, 297-320.
- (96) Gomis, V.; Font, A.; Pedraza, R.; Saquete, M. D. Isobaric vapor-liquid and vapor-liquid-liquid equilibrium data for the water-ethanol-hexane system. *Fluid Phase Equilib.* **2007**, *259*, 66-70.
- (97) Ross, S.; Patterson, R. E. Surface and Interfacial Tensions of Conjugate Solutions in Ternary Systems. *J. Chem. Eng. Data.* **1979**, *24*, 111-115.
- (98) Brandani, V.; Chianese, A.; Rossi, M. Ternary liquid-liquid equilibrium data for the water-ethanol-benzene system. *J. Chem. Eng. Data.* **1985**, *30*, 27-29.
- (99) Mostafazadeh, A. K.; Rahimpour, M. R.; Shariati, A. Vapor-liquid equilibria of water + triethylene glycol (TEG) and water + TEG + toluene at 85 kPa. *J. Chem. Eng. Data.* **2009**, *54*, 876-881.
- (100) Ng, H. J.; Chen, C. J.; Robinson, D. B. Vapour-liquid and vapour-liquid-liquid equilibria for H₂S, CO₂, Selected light hydrocarbons and a gas condensate in aqueous methanol or ethylene glycol solutions. GPA Research Report 149. Gas Processors Association: Tulsa, Okla. Gas Research Institute: Chicago, IL. **1995**.
- (101) Chen, C. J.; Ng, H. J.; Robinson, D. B. The solubility of methanol or glycol in water-hydrocarbon systems. GPA Research Report 117. Gas Processors Association: Tulsa, Okla. 1988.
- (102) Ng, H. J.; Chen, C. J.; Robinson, D. B. The influence of high concentrations of methanol on hydrate formation and the distribution of glycol in liquid-liquid mixtures. GPA Research Report 106. Gas Processors Association: Tulsa, Okla. **1987**.
- (103) Ashcroft, S. J.; Brunner, G.; Vollmer, H.; Sweeney, C. W. Three-phase vapor-liquid-liquid equilibria for methane + n-octane + monoethylene glycol + water at pressures from (3.1 to 15.0) MPa and temperatures from (281 to 363) K. *J. Chem. Eng. Data.* **2009**, *54*, 1613-1618.
- (104) Clark, G. N. I.; Haslam, A. J.; Galindo, A.; Jackson, G. Developing optimal Wertheim-like models of water for use in Statistical Associating Fluid Theory (SAFT) and related approaches. *Mol. Phys.* **2006**, *104*, 3561-3581.

- (105) Grenner, A.; Kontogeorgis, G. M.; Michelsen, M. L.; Folas, G. K. On the estimation of water pure compound parameters in association theories. *Mol. Phys.* **2007**, *105*, 1797-1801.
- (106) Liang, X. D.; Tsivintzelis, I.; Kontogeorgis, G. M. Modeling water containing systems with the simplified PC-SAFT and CPA equations of state. *Ind. Eng. Chem. Res.* **2014**, *53*, 14493-14507.
- (107) Liang, X. D. Thermodynamic modeling of complex systems. PhD Thesis, Technical University of Denmark, Kgs. Lyngby, Denmark, **2014**.
- (108) Huang, S. H.; Radosz, M. Equation of state for small, large, polydisperse, and associating molecules. *Ind. Eng. Chem. Res.* **1990**, *29*, 2283-2294.
- (109) Fouad, W. A.; Wang, L.; Haghmoradi, A.; Asthagiri, D.; Chapman, W. G. Understanding the thermodynamics of hydrogen bonding in alcohol-containing mixtures: cross-association. *J. Phys. Chem. B*, **2016**, *120*, 3388-3402.
- (110) DIPPR Database, Design Institute for Physical Property Data; DIADEM, Information and Data Evaluation Manager, version 6.0.1, **2012**.
- (111) REFPROP, Reference Fluid Thermodynamic and Transport Properties, NIST Standard Reference Database 23, Version 9.0, **2010**.
- (112) NIST Chemistry WebBook. <http://webbook.nist.gov/chemistry/> (accessed March 14, **2017**).
- (113) Tsivintzelis, I.; Kontogeorgis, G. M.; Michelsen, M. L.; Stenby, E. H. Modeling phase equilibria for acid gas mixtures using the CPA equation of state. I. Mixtures with H₂S. *AIChE J.* **2010**, *56*, 2965-2982.
- (114) Kontogeorgis, G. M.; Yakoumis, I. V.; Meijer, H.; Hendriks, E.; Moorwood, T. Multicomponent phase equilibrium calculations for water-methanol-alkane mixtures. *Fluid Phase Equilib.* **1999**, *158-160*, 201-209.
- (115) Yakoumis, I. V.; Kontogeorgis, G. M.; Voutsas, E. C.; Tassios, D. P. Vapor-liquid equilibria for alcoholhydrocarbon systems using the CPA equation of state. *Fluid Phase Equilib.* **1997**, *130*, 31-47.
- (116) Derawi, S. O.; Michelsen, M. L.; Kontogeorgis, G. M.; Stenby, E. H. Application of the CPA equation of state to glycol/hydrocarbons liquid-liquid equilibria. *Fluid Phase Equilib.* **2003**, *209*, 163-184.
- (117) Liang, X. D.; Maribo-Mogensen, B.; Thomsen, K.; Yan, W.; Kontogeorgis, G. M. Approach to improve speed of sound calculation within PC-SAFT framework. *Ind. Eng. Chem. Res.* **2012**, *51*, 14903-14914.
- (118) Tsivintzelis, I.; Grenner, A.; Economou, I. G.; Kontogeorgis, G. M. Evaluation of the nonrandom hydrogen bonding (NRHB) theory and the simplified perturbed-chain – statistical associating fluid theory (sPC-SAFT). 2. Liquid-liquid equilibria and prediction of monomer fraction in hydrogen bonding systems. *Ind. Eng. Chem. Res.* **2008**, *47*, 5651-5659.
- (119) Liang, X. D.; Thomsen, K.; Yan, W.; Kontogeorgis, G. M. Modeling the liquid-liquid equilibrium of petroleum fluid and polar compounds containing systems with the PC-SAFT equation of state. *Fluid Phase Equilib.* **2015**, *406*, 147-155.
- (120) Grenner, A.; Kontogeorgis, G. M.; von Solms, N.; Michelsen, M. L. Modeling phase equilibria of alkanols with the simplified PC-SAFT equations of state and generalized pure compound parameters. *Fluid Phase Equilib.* **2007**, *258*, 83-94.

For Table of Contents Only

