



Complete Genome Sequence of *Shewanella* sp. WE21, a Rare Isolate with Multiple Novel Large Genomic Islands

Castillo, Daniel; Gram, Lone; Dailey, Frank E.

Published in:
Genome Announcements

Link to article, DOI:
[10.1128/genomeA.00277-18](https://doi.org/10.1128/genomeA.00277-18)

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Castillo, D., Gram, L., & Dailey, F. E. (2018). Complete Genome Sequence of *Shewanella* sp. WE21, a Rare Isolate with Multiple Novel Large Genomic Islands. *Genome Announcements*, 6(16), [e00277-18].
<https://doi.org/10.1128/genomeA.00277-18>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Complete Genome Sequence of *Shewanella* sp. WE21, a Rare Isolate with Multiple Novel Large Genomic Islands

Daniel Castillo,^a Lone Gram,^b Frank E. Dailey^c

^aMarine Biological Section, University of Copenhagen, Helsingør, Denmark

^bDepartment of Biotechnology and Biomedicine, Technical University of Denmark, Kongens Lyngby, Denmark

^cDepartment of Biology, University of Wisconsin–Sheboygan, Sheboygan, Wisconsin, USA

ABSTRACT We present here the whole-genome sequence of *Shewanella* sp. WE21, an unusual omega-3 fatty acid-producing bacterium isolated from the gastrointestinal tract of the freshwater fish *Sander vitreus* (walleye). This genome contains a number of unique, large genomic islands with genes not present in other *Shewanella* bacteria.

Shewanella spp. are Gram-negative, motile bacilli that are found in many environments but most often in aquatic environments, particularly in marine ones. They are important for the turnover of organic material, and they are capable of dissimilatory reduction of various metals and other substances. (1). Also, several *Shewanella* strains have been reported as candidates for biotechnological applications such as omega-3 fatty acid production (2) and bioremediation of heavy metals (1). In contrast to their beneficial roles, *Shewanella* strains have been identified as major agents of fish spoilage (3) and recently as fish pathogens responsible for outbreaks of disease (4). Furthermore, they are increasingly being implicated as opportunistic human pathogens (5). *Shewanella* sp. WE21, isolated from the gastrointestinal tract of the freshwater fish species *Sander vitreus* (walleye) in Lac des Mille, Ontario, Canada (48°84'N, 90°51'W) (6), represents a novel omega-3 fatty acid-producing freshwater isolate of *Shewanella*. It appears most closely related to *S. putrefaciens* strains by phylogenetic analysis (6). However, *in silico* DNA-DNA hybridization suggests that *Shewanella* sp. WE21 is more closely related to (but distinct from) *S. baltica* species (F. Dailey, unpublished results). Here, we report the whole-genome sequence of this strain.

PacBio sequencing with 100× coverage was performed as previously described (6) for *Shewanella* sp. strain WE21. Annotation was performed by using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (7). Additionally, the genomes were analyzed on the Rapid Annotations using Subsystems Technology (RAST) server (8). Acquired antibiotic resistance genes were identified by using ResFinder version 2.1 (9), virulence factors by using VirulenceFinder version 1.2 (10), clustered regularly interspaced short palindromic repeat (CRISPR) arrays by using CRISPRFinder (11), genomic islands by using IslandViewer version 4 (12), and prophage-related sequences by using PHASTER (13).

The final assembly for *Shewanella* sp. strain WE21 resulted in one large closed contig with a total length of 5,266,746 bp and a G+C content of 45.3%. Genome annotation resulted in 4,416 coding sequences (CDSs) plus one potential overlapping CDS, 105 tRNAs, 176 pseudogenes, and 31 rRNAs; 96.2% of the CDSs were classified as hypothetical proteins. Five prophage-like elements between 33.7 to 43.4 kb in length were detected. No virulence factors were identified in these prophage sequences. In addition, four genomic islands of 131.4 kb, 106.5 kb, 28.7 kb, and 9.2 kb, which have GC% contents of 40.6, 42.3, 38.3, and 57.9, respectively, were identified. In the first genomic island, we detected genes encoding putative virulence factors found in *Vibrio anguil-*

Received 5 March 2018 Accepted 22 March 2018 Published 19 April 2018

Citation Castillo D, Gram L, Dailey FE. 2018. Complete genome sequence of *Shewanella* sp. WE21, a rare isolate with multiple novel large genomic islands. *Genome Announc* 6:e00277-18. <https://doi.org/10.1128/genomeA.00277-18>.

Copyright © 2018 Castillo et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Frank E. Dailey, frank.dailey@uw.edu.

larum (14) and other fish pathogens, as well as a type III secretion system with large overlapping genes. Analysis of the second reveals not only additional genes encoding vibrio-like virulence factors, such as *cnf2*, but also genes encoding insecticidal toxins (15) and polyketides. The third genomic island putatively encodes very novel pili, and the fourth encodes copper efflux genes. Antibiotic resistance genes penicillin (*BL*) and fluoroquinolones (*parC*, *parE*, *gyrA*, and *gyrB*) were detected. Also, cobalt-zinc-cadmium resistance genes (*CzcD*, *CzcA*, *CusA*, *CusB*, and *TR*) and arsenic resistance genes were identified. No CRISPR arrays or plasmids were detected. This genome sequence can facilitate the understanding of symbiosis and virulence in fish intestinal bacteria.

Accession number(s). The draft genome sequence of *Shewanella* sp. strain WE21 can be accessed under the GenBank accession number [CP023019](https://ncbi.nlm.nih.gov/nucl/CP023019). The version described in this paper is the first version.

ACKNOWLEDGMENTS

We acknowledge the Institute of Genome Sciences (IGS), University of Maryland, for performing the genomic sequencing. We thank the Leibniz Institute DSMZ for the use of the Genome-to-Genome Distance Calculator version 2.1 for the *in silico* analyses and Michael Thomas (Bacteriology, University of Wisconsin) for the initial analysis of the polyketide-encoding genes.

We also acknowledge the support of the Department of Biology, University of Wisconsin Colleges (F. E. Dailey), as well as previous support from Concordia University, Wisconsin, for the genome sequencing. This work was also supported by grant 12-132390 (ProAqua project) from the Innovation Fund Denmark (L. Gram and D. Castillo).

REFERENCES

1. Fredrickson JK, Romine MF, Beliaev AS, Auchtung JM, Driscoll ME, Gardner TS, Nealson KH, Osterman AL, Pinchuk G, Reed JL, Rodionov DA, Rodrigues JLM, Saffarini DA, Serres MH, Spormann AM, Zhulin IB, Tiedje JM. 2008. Towards environmental systems biology of *Shewanella*. *Nat Rev Microbiol* 6:592–603. <https://doi.org/10.1038/nrmicro1947>.
2. Amiri-Jami M, Griffiths MW. 2010. Recombinant production of omega-3 fatty acids in *Escherichia coli* using a gene cluster isolated from *Shewanella baltica* MAC1. *J Appl Microbiol* 109:1897–1905. <https://doi.org/10.1111/j.1365-2672.2010.04817.x>.
3. Vogel BF, Venkateswaran K, Satomi M, Gram L. 2005. Identification of *Shewanella baltica* as the most important H₂S-producing species during iced storage of Danish marine fish. *Appl Environ Microbiol* 71:6689–6697. <https://doi.org/10.1128/AEM.71.11.6689-6697.2005>.
4. Pękala A, Kozińska A, Paździor E, Glowacka H. 2015. Phenotypical and genotypical characterization of *Shewanella putrefaciens* strains isolated from diseased freshwater fish. *J Fish Dis* 38:283–293. <https://doi.org/10.1111/jfd.12231>.
5. Janda JM, Abbott SL. 2014. The genus *Shewanella*: from the briny depths below to human pathogen. *Crit Rev Microbiol* 40:293–312. <https://doi.org/10.3109/1040841X.2012.726209>.
6. Dailey FE, McGraw JE, Jensen BJ, Bishop SS, Lokken JP, Dorff KJ, Ripley MP, Munro JB. 2016. The microbiota of freshwater fish and freshwater niches contain omega-3 fatty acid-producing *Shewanella* species. *Appl Environ Microbiol* 82:218–231. <https://doi.org/10.1128/AEM.02266-15>.
7. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
8. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parelllo B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
9. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
10. Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM, Aarestrup FM. 2014. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J Clin Microbiol* 52:1501–1510. <https://doi.org/10.1128/JCM.03617-13>.
11. Grissa I, Vergnaud G, Pourcel C. 2007. CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res* 35:W52–W57. <https://doi.org/10.1093/nar/gkm360>.
12. Langille MG, Brinkman FS. 2009. IslandViewer: an integrated interface for computational identification and visualization of genomic islands. *Bioinformatics* 25:664–665. <https://doi.org/10.1093/bioinformatics/btp030>.
13. Arndt D, Grant J, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
14. Castillo D, Alvisé PD, Xu R, Zhang F, Middelboe M, Gram L. 2017. Comparative genome analyses of *Vibrio anguillarum* strains reveal a link with pathogenicity traits. *mSystems* 2:e00001-17. <https://doi.org/10.1128/mSystems.00001-17>.
15. Bowen D, Rocheleau TA, Blackburn M, Andreev O, Golubeva E, Bhartia R, Ffrench-Constant RH. 1998. Insecticidal toxins from the bacterium *Photobacterium luminescens*. *Science* 280:2129–2132. <https://doi.org/10.1126/science.280.5372.2129>.