



Physical activity monitors to enhance the daily amount of physical activity in elderly-a protocol for a systematic review and meta-analysis

Larsen, Rasmus Tolstrup; Christensen, Jan; Juhl, Carsten Bøgh; Andersen, Henning Boje; Langberg, Henning

Published in:
Systematic Reviews

Link to article, DOI:
[10.1186/s13643-018-0733-6](https://doi.org/10.1186/s13643-018-0733-6)

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Larsen, R. T., Christensen, J., Juhl, C. B., Andersen, H. B., & Langberg, H. (2018). Physical activity monitors to enhance the daily amount of physical activity in elderly-a protocol for a systematic review and meta-analysis. *Systematic Reviews*, 7(1), [69]. <https://doi.org/10.1186/s13643-018-0733-6>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

PROTOCOL

Open Access



Physical activity monitors to enhance the daily amount of physical activity in elderly—a protocol for a systematic review and meta-analysis

Rasmus Tolstrup Larsen^{1*} , Jan Christensen^{1,2}, Carsten Bogh Juhl^{4,5}, Henning Boje Andersen³ and Henning Langberg¹

Abstract

Background: To investigate the use of physical activity monitors (PAMs) for the elderly, the scientific literature should be systematically reviewed and the effect quantified, as the evidence seems inconclusive.

Methods and design: Randomized controlled trials and randomized crossover trials, with participants with a mean age above 65 years, comparing any PAM intervention with other control interventions or no intervention, will be included. This protocol is detailed according to the recommendations of the Cochrane Handbook, and it is reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols statement.

Results: We will present results from the search in a flow diagram. The results from the analyses will include regular meta-analyses, stratified analyses, and meta-regressions. The results on each outcome of interest will be presented in a summary of findings table.

Discussion: This paper will explore and analyze the heterogeneity of the results and try to identify variables that will enhance the effect of PAMs in elderly. The results will be useful to researchers working with elderly and/or PAMs, health care professionals working with elderly, and relatives together with the elderly themselves.

Systematic review registration: PROSPERO [CRD42018083648](https://doi.org/10.1186/1745-7189-4-2018083648).

Keywords: Elderly, Physical activity, Walking, Physical activity monitors, Feedback, Motivation, Behavioral change, Randomized controlled trials, Systematic review, Meta-analysis

Background

Rationale

According to the World Health Organization, elderly who are physically active have lower rates of all-cause mortality and exhibit higher levels of functional health [1]. Even a small change in the daily amount of physical activity may be beneficial on hard outcomes such as all-cause mortality and life expectancy [2]. The American College of Sports Medicine describes walking as the most common

type of physical activity among elderly [3] and walking may play a key role in prevention of cardiovascular disease [4].

Physical activity monitors (PAMs) were originally used to quantify the level of physical activity through the amount of steps taken [5] and have been used in research since the 1960s [6]. However, PAMs are also used effectively to motivate and facilitate to an enhanced level of physical activity, as meta-analyses have reported PAM-based interventions to reduce participant weight in weight loss programs [7], increase the level of physical activity [8], and reduce sedentary time significantly [9]. PAMs provide feedback on physical activity [5]. This

* Correspondence: rala@sund.ku.dk

¹CopenRehab, Section of Social Medicine, Department of Public Health, University of Copenhagen, Copenhagen, Denmark

Full list of author information is available at the end of the article



feedback might facilitate result-driven behavioral change and, hence, increase the level of physical activity [10]. Especially in elderly, PAM-based interventions have been shown to be feasible and effective in enhancing the level of physical activity [11–16]. However, some studies report no significant differences between PAM groups and control groups [17–19]. This could be explained by the devices being difficult and troublesome for elderly to use, and a survey has reported that a third of PAM consumers have stopped using the devices after 6 months [20]. It seems very relevant to investigate the use of PAMs, specifically in elderly as they are expected to behave differently than younger adults. To investigate the above, the scientific literature should be systematically reviewed; the effect quantified and possible factors explaining differences in effect size should be identified.

Research questions and objective

Research questions

1. What is the effect of a PAM-based intervention on physical activity behavior in elderly?
2. What are the potential effects on other outcomes such as changes in body mass index, physical capacity, and health-related quality of life?
3. Which factors explain heterogeneity of the results?

Objective

The objective of this systematic review and meta-analysis is to review the literature and estimate the effect on daily level of physical activity, when using PAMs as an intervention, compared to control interventions in participants aged 65 years and over. Furthermore, to investigate if potential physical activity effects can result in changes in secondary outcomes such as body mass index, physical capacity, and health-related quality of life. Lastly, possible factors explaining heterogeneity will be investigated.

Methods

This protocol is detailed according to the recommendations of the Cochrane Handbook [21], and it is reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols (PRISMA-P) statement.

Eligibility criteria

Types of studies

Randomized controlled trials (RCTs) and randomized crossover trials will be included.

Types of participants

Participants included in eligible studies must have a mean age above 65 years. The participants must be independent walkers with or without walking aids.

Types of interventions

For this review, we will include studies comparing any PAM intervention with other control interventions or no intervention. The PAMs may be portable or wearable, electronic or mechanical, and driven by accelerometers, pedometers, or global positioning system (GPS).

Types of outcome measures

Studies must report at least one of the primary or secondary outcomes in order to be included.

Primary outcome The primary outcome is changed in daily amount of physical activity. If more than one relevant outcome is reported within a study, the outcome will be extracted or calculated favoring daily number of steps, followed by daily number of meters walked, daily amount of energy expenditure measured as calories, daily metabolic equivalent of task (minutes or hours), and finally, if no objective measure is available, self-reported physical activity.

Secondary outcomes Secondary outcomes include (a) meeting the study-specific recommended level of physical activity, (b) change in time spent sedentary, (c) change in time spent in moderate activity, (d) change in time spent in vigorous activity, (e) change in physical capacity, (f) changes in body mass index, and (g) changes in self-reported outcomes. All secondary outcomes (a to g) are independent from each other and will be extracted and analyzed accordingly.

- a. Meeting the study-specific recommended level of physical activity, measured objectively by PAMs will be extracted from any study that provides results on this.
- b. Time spent sedentary will be extracted favoring PAM measured behavior. If no PAM measured

Abstract]) OR “accelerometer-based tracker” [Title/Abstract]) OR “moov now” [Title/Abstract]) OR “misfit ray” [Title/Abstract]) OR “nokia go” [Title/Abstract]) OR “activity monitor” [Title/Abstract]) OR Fitbit [Title/Abstract]) OR Pedometer [Title/Abstract]) OR “step monitor” [Title/Abstract]) OR “physical activity monitor” [Title/Abstract]) OR “Step counter” [Title/Abstract])) AND (((((((“older people” [Title/Abstract]) OR (“older adults”[Title/Abstract]) OR (“old adults”[Title/Abstract]) OR (“residents”[Title/Abstract]) OR elderly[Title/Abstract]) OR “Above 60 years” [Title/Abstract]) OR Seniors [Title/Abstract]) OR Aged [MeSH]) OR frail elderly [MeSH]) OR ((Aged, 80 and over [MeSH]))))

Search string for CINAHL (AB “older adults” OR AB “older people” OR “residents” OR “frail elderly” AB “old adults” OR AB elderly OR AB “Above 60 years” OR AB Seniors OR AB Aged OR AB (Aged, 80 and over)) AND (AB “Step counter” OR AB “physical activity monitor” OR AB “step monitor” OR AB Pedometer OR AB Fitbit OR AB “activity monitor” OR AB “nokia go” OR AB “misfit ray” OR AB “moov now” OR AB “accelerometer-based tracker” OR AB “xiaomi mi band” OR AB “tomtom” OR AB “vivoactive” OR AB “jawbone” OR AB “movement counter” OR AB “quantified movement” OR AB “fitness tracker” OR AB “activity monitoring device” OR AB (physic* AND activit* AND monitor*) OR AB (PAM AND monitor*)) AND (AB “randomized controlled trial” OR AB “controlled clinical trial” OR AB “cross-over trial” OR AB “cross over trial” OR AB “randomized” OR AB “clinical trial” OR AB “randomly”)

Search string for EMBASE

1. (old adults or elderly or seniors or residents or older adults or older people or frail elderly).ab.
2. ((PAM and monitor*) or (physic* and activit* and monitor*) or “activity monitoring device” or “fitness tracker” or “quantified movement” or “movement counter” or “jawbone” or “vivoactive” or “tomtom” or “xiaomi mi band” or “accelerometer-based tracker” or “moov now” or “misfit ray” or “nokia go” or “activity monitor” or Fitbit or Pedometer or “step monitor” or “physical activity monitor” or “Step counter”).ab.
3. (randomized controlled trial or controlled clinical trial or cross over trial or randomized or clinical trial).ab.
4. aged/
5. controlled clinical trial/ or “randomized controlled trial (topic)”/
6. 1 or 4
7. 3 or 5
8. 2 and 6 and 7

Search string for SPORTDiscus (AB “older adults” OR AB “older people” OR “residents” OR “frail elderly” AB “old adults” OR AB elderly OR AB “Above 60 years” OR AB Seniors OR AB Aged OR AB (Aged, 80 and over)) AND (AB “Step counter” OR AB “physical activity monitor” OR AB “step monitor” OR AB Pedometer OR AB Fitbit OR AB “activity monitor” OR AB “nokia go” OR AB “misfit ray” OR AB “moov now” OR AB “accelerometer-based tracker” OR AB “xiaomi mi band” OR AB “tomtom” OR AB “vivoactive” OR AB “jawbone” OR AB “movement counter” OR AB “quantified movement” OR AB “fitness tracker” OR AB “activity monitoring device” OR AB (physic* AND activit* AND monitor*) OR AB (PAM AND monitor*)) AND (AB “randomized controlled trial” OR AB “controlled clinical trial” OR AB “cross-over trial” OR AB “cross over trial” OR AB “randomized” OR AB “clinical trial” OR AB “randomly”)

Search string for CENTRAL

1. Mesh descriptor:[Aged] explode all trees
2. Mesh descriptor:[Frail Elderly] explode all trees
3. “old adults” or elderly or “older adults” or “older people” or “frail elderly” or seniors: ti,ab,kw
4. (PAM and monitor*) or (physic* and activit* and monitor*) or “activity monitoring device” or “fitness tracker” or “quantified movement” or “movement counter” or “jawbone” or “vivoactive” or “tomtom” or “xiaomi mi band” or “accelerometer-based tracker” or “moov now” or “misfit ray” or “nokia go” or “activity monitor” or Fitbit or Pedometer or “step monitor” or “physical activity monitor” or “Step counter”:ti,ab,kw
5. (#1 or #2 or #3) and #4

Searching other resources

Searching references of eligible studies and relevant journals by pearl growing will be conducted independently by two reviewers (RTL and JC) in order to include relevant articles not captured by the search strings. The database Clinicatrials.gov will be used to locate ongoing relevant studies.

Data collection and analysis

Data management

The technology platform, *Covidence*, will be used to import citations from the literature searches, screening of title and abstracts, screening of full text, assessing risk of bias in included studies, and extracting the data. The analyses will be conducted in *Stata Statistical Software, version 15*.

Selection of studies

The selection of studies will be done by merging search results from the databases, removing duplicates, examining the titles, and examining full-text reports according to the inclusion criteria.

Two authors (RTL, JC) will independently screen titles and abstracts and assess eligible articles in full text. Any inconsistencies between authors will be discussed and solved with consultation of a third author (CJ).

Data extraction and management

Data on the following items will be extracted from all included studies.

Source Study ID, protocol ID, review author, citation, and contact details

Methods Study design, aim of study, number of arms or groups, funding source, informed consent obtained, and ethical approval

Participants Total number of participants, setting, possible diagnostic criteria, age, sex, country, comorbidities, education length, and marriage status

Interventions Duration of intervention, specific intervention, and intervention details sufficient for replication

Outcomes All outcomes specified in the “Types of outcome measures” section and specific time points, outcome definitions, and unit of measurement

Results Number of participants allocated to each group, summary data for each intervention, and control groups (as reported) including adverse events

Miscellaneous Funding sources, key conclusions, miscellaneous comments from authors and if correspondence was required

Assessment of risk of bias in included studies

Two review authors (RTL and JC) will independently assess the risk of bias in included studies, using the *RoB 2.0 tool* [28]. Disagreement between review authors will be solved by including a third reviewer (CJ). The risk of bias assessment for each study will be presented using a table with judgment and support for judgment.

Measures of treatment effect

Treatment effect, measured as continuous data, will be expressed as mean difference with 95% confidence intervals for outcomes measured with the same outcome measurement instrument, or as standardized

mean difference with 95% confidence intervals, when different measurement instruments are used in the included studies. Dichotomous outcomes, such as adverse events or meeting the recommended level of physical activity, will be analyzed and expressed as risk ratios with 95% confidence intervals.

Unit of analysis issues

Studies with multiple treatment groups If a study has more than one intervention group and both seem relevant for inclusion in the systematic review, the intervention groups will be included as two separate studies and the control group from the study will be separated [29].

Assessment of heterogeneity

The heterogeneity of the study results will be examined using Cochran Q test and quantified with I^2 values and the between study variance τ^2 .

Assessment of small study bias

Small study bias will be assessed by calculating an *Egger's test* score and illustrated with a funnel plot. If small study bias is found, by a positive *Egger's test*, a *metatrim* analysis will be conducted and an adjusted effect size will be calculated.

Data synthesis

If two or more included studies allows for it, the effect size will be calculated using a random-effects meta-analysis adjusting to Hedges' g . If the included studies need network meta-analysis to be pooled this will be performed, as described in Chapter 16.6.3 of the *Cochrane Handbook* [30]. An alpha level of 0.05 will be considered statistical significant. If it is not possible to conduct a meta-analysis, we will describe the data narratively specific to each outcome.

Subgroup analysis and investigation of heterogeneity

We will explore heterogeneity by conducting sub-group analyses and stratified analyses on the following nominal variables:

- Placement of PAM (ankle worn, wrist worn, hip worn)
- Type of PAM (pedometer versus accelerometer)
- Diagnoses (healthy, cancer patients, pulmonary patients, cardiovascular patients, etc.)
- Feedback frequency (real time, daily, weekly or monthly)

We will explore heterogeneity on continuous data by performing univariate meta-regressions on the following variables:

- Mean age
- Sex distribution
- Number (or percent) of participants with walking aids
- Intervention length
- Mean baseline physical activity

Sensitivity analysis

Sensitivity analyses will be performed on all outcomes of interest by stratifying on overall risk of bias, defined by the RoB 2.0 tool [28] (low/some concerns/high). Furthermore, we will perform sensitivity analyses on how the primary outcome has been extracted, by performing meta-analyses with mean differences on (1) change in daily number of steps, (2) change in daily number of meters walked, (3) change in daily amount of energy expenditure measured as calories, (4) change in daily metabolic equivalent of task (minutes or hours), and (5) self-reported physical activity.

Summary of findings table

We will create a Summary of findings table with effect sizes on the outcomes earlier presented. Two reviewers (RTL and JC) will independently rate the quality of the evidence using the GRADE approach. Downgrading and upgrading the quality of the evidence will be described in the footnotes in the table. An empty example of a summary of findings table is listed in Additional file 1.

Discussion

The aim of this systematic review and meta-analysis is to systematically locate, evaluate, summarize, and analyze available evidence regarding the use of PAMs to enhance the level of physical activity in elderly. This paper will explore and analyze the heterogeneity of the results and try to identify variables that will enhance the effect of PAMs in elderly. The discussion will contain a cost-effective evaluation of the effect size of PAMs and the expected prize of the monitors. We anticipate that this review and the results will be useful to researchers working with elderly and/or PAMs, health care professionals working with elderly, and relatives together with the elderly themselves.

Strength of this systematic review and meta-analysis

We have not identified any systematic reviews investigating the effect of PAM-based interventions in elderly. This paper will be the first systematic review and meta-analysis directly calculating the effect size and exploring heterogeneity of the results. Furthermore, this systematic review and meta-analysis will only include RCTs and randomized crossover trials and hence, the level of evidence is likely to be high.

Additional file

Additional file 1: Table S1. Summary of findings. (DOCX 17 kb)

Abbreviations

GPS: Global positioning system; PAMs: Physical activity monitors; PRISMA-P: Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols; RCTs: Randomized controlled trials

Funding

The content presented within this paper was produced as part of the project REACH: this project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement no. 690425 [31].

Authors' contributions

RTL has been the lead author on this protocol and has participated in all steps if the writing process. JC has together with CJ and RTL designed the search strategy and has participated in all steps of the writing process. CJ together with RTL has designed the "Data collection and analysis" section and has participated in all steps of the writing process. HBA and HL together with RTL have participated in the preliminary design of the study and have participated in all steps of the writing process. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹CopenRehab, Section of Social Medicine, Department of Public Health, University of Copenhagen, Copenhagen, Denmark. ²Department of Occupational and Physiotherapy, Copenhagen University Hospital, Copenhagen, Denmark. ³DTU Management Engineering Inst., Technical University of Denmark, Diplomvej 372 office 226, 2800 Lyngby, Denmark. ⁴Research Unit of Musculoskeletal Function and Physiotherapy, Institute of Sports Science and Clinical Biomechanics, Faculty of Health Sciences, University of Southern Denmark, Odense, Denmark. ⁵Department of Rehabilitation, Copenhagen University Hospital, Herlev and Gentofte, Denmark.

Received: 3 January 2018 Accepted: 13 April 2018

Published online: 02 May 2018

References

1. World Health Organization. Physical Activity and Older Adults. World Health Organization; 2015. www.who.int/dietphysicalactivity/factsheet_olderadults/en/. Accessed 18 Dec 2017.
2. Wen CP, Wai JPM, Tsai MK, Yang YC, Cheng TYD, Lee M-C, et al. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lancet Lond Engl*. 2011;378:1244–53.
3. Chodzko-zajko WJ, Proctor DN, Singh MAF, Minson CT, Nigg CR, Salem GJ, et al. Exercise and physical activity for older adults. *Med Sci Sports Exerc*. 2009;41:1510–30.
4. Murtagh EM, Murphy MH, Boone-Heinonen J. Walking: the first steps in cardiovascular disease prevention. *Curr Opin Cardiol*. 2010;25:490–6.
5. Bassett DR, Toth LP, LaMunion SR, Crouter SE. Step counting: a review of measurement considerations and health-related applications. *Sports Med*. 2017;47:1303–15.
6. Stunkard A. A method of studying physical activity in man. *Am J Clin Nutr*. 1960;8:595–601.
7. Richardson CR, Newton TL, Abraham JJ, Sen A, Jimbo M, Swartz AM. A meta-analysis of pedometer-based walking interventions and weight loss. *Ann Fam Med*. 2008;6:69–77.

8. Kang M, Marshall SJ, Barreira TV, Lee J-O. Effect of pedometer-based physical activity interventions: a meta-analysis. *Res Q Exerc Sport*. 2009;80:648–55.
9. Qiu S, Cai X, Ju C, Sun Z, Yin H, Zügel M, et al. Step counter use and sedentary time in adults. *Medicine (Baltimore)*. 2015;94 <https://doi.org/10.1097/MD.0000000000001412>.
10. Patel MS, Asch DA, Volpp KG. Wearable devices as facilitators, not drivers, of health behavior change. *JAMA*. 2015;313:459–60.
11. Harris T, Kerry SM, Victor CR, Ekelund U, Woodcock A, Iliffe S, et al. A primary care nurse-delivered walking intervention in older adults: PACE (pedometer accelerometer consultation evaluation)-lift cluster randomised controlled trial. *PLoS Med*. 2015;12:e1001783.
12. McMahon SK, Lewis B, Oakes M, Guan W, Wyman JF, Rothman AJ. Older adults' experiences using a commercially available monitor to self-track their physical activity. *JMIR MHealth UHealth*. 2016;4:e35.
13. Kawagoshi A, Kiyokawa N, Sugawara K, Takahashi H, Sakata S, Satake M, et al. Effects of low-intensity exercise and home-based pulmonary rehabilitation with pedometer feedback on physical activity in elderly patients with chronic obstructive pulmonary disease. *Respir Med*. 2015;109:364–71.
14. Nishiguchi S, Yamada M, Tanigawa T, Sekiyama K, Kawagoe T, Suzuki M, et al. A 12-week physical and cognitive exercise program can improve cognitive function and neural efficiency in community-dwelling older adults: a randomized controlled trial. *J Am Geriatr Soc*. 2015;63:1355–63.
15. Peel NM, Paul SK, Cameron ID, Crotty M, Kurrle SE, Gray LC. Promoting activity in geriatric rehabilitation: a randomized controlled trial of accelerometry. *PLoS One*. 2016;11:e0160906.
16. Croteau KA, Richeson NE, Farmer BC, Jones DB. Effect of a pedometer-based intervention on daily step counts of community-dwelling older adults. *Res Q Exerc Sport*. 2007;78:401–6.
17. Nolan CM, Maddocks M, Canavan JL, Jones SE, Delogu V, Kalaraju D, et al. Pedometer step count targets during pulmonary rehabilitation in chronic obstructive pulmonary disease. A randomized controlled trial. *Am J Respir Crit Care Med*. 2017;195:1344–52.
18. Takahashi PY, Quigg SM, Croghan IT, Schroeder DR, Ebbert JO. Effect of pedometer use and goal setting on walking and functional status in overweight adults with multimorbidity: a crossover clinical trial. *Clin Interv Aging*. 2016;11:1099–106.
19. McMurdo MET, Sugden J, Argo I, Boyle P, Johnston DW, Sniehotta FF, et al. Do pedometers increase physical activity in sedentary older women? A randomized controlled trial. *J Am Geriatr Soc*. 2010;58:2099–106.
20. Partners E. Inside wearables: how the science of human behavior change offers the secret to long-term... Endeavour partners. 2017. <https://blog.endeavourpartners.com/inside-wearable-how-the-science-of-human-behavior-change-offers-the-secret-to-long-term-engagement-a15b3c7d4cf3>. Accessed 7 Feb 2018.
21. Higgins J, Green S. *Cochrane Handbook For Systematic Reviews Of Interventions*. 2011. <http://handbook.cochrane.org/>. Accessed 11 Apr 2016.
22. pubmeddev. MEDLINE database (pubmed). <https://www.ncbi.nlm.nih.gov/pubmed>. Accessed 3 Oct 2016.
23. Embase database. <http://www.embase.com.ez-jmk.statsbiblioteket.dk:2048/#search>. Accessed 3 Oct 2016.
24. Inc EIS. SPORTDiscus | Sports Medicine Database | EBSCO. EBSCO Information Services, Inc. | www.ebsco.com. page url. Accessed 21 Nov 2017.
25. EBSCO. CINAHL plus with full text[full text nursing journals | EBSCO | EBSCO health. 2016. <https://health.ebsco.com/products/cinahl-plus-with-full-text>. Accessed 23 Nov 2015.
26. CENTRAL database. <http://onlinelibrary.wiley.com/cochranelibrary/search>. Accessed 3 Oct 2016.
27. Lefebvre C, Manheimer E, Glanville J. Box 6.4.a: Cochrane highly sensitive search strategy for identifying randomized trials in MEDLINE: sensitivity-maximizing version (2008 revision); PubMed format. 2011.
28. Higgins J, Savovic J, Sterne JAC, Page M, Hróbjartsson A, Boutron A, et al. A revised tool to assess risk of bias in randomized trials (RoB 2.0). 2016. https://sites.google.com/site/riskofbiastool/20161020_RoB_2.0_template_parallel_assignment.docx?attredirects=0&d=1.
29. Higgins, Green. 16.5.4 How to include multiple groups from one study. 2011. http://handbook-5-1.cochrane.org/chapter_16/16_5_4_how_to_include_multiple_groups_from_one_study.htm. Accessed 4 Dec 2017.
30. Higgins JPT, Green S. 16.6.3 Multiple-treatments meta-analysis—Cochrane Handbook for Systematic Reviews Of Interventions. 2011. <http://handbook-5-1.cochrane.org/>. Accessed 19 Oct 2017.
31. REACH | HORIZON 2020. <http://reach2020.eu/>. Accessed 23 Nov 2017.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

