



EU-OPENSOURCE: A novel collaborative approach to facilitate chemical biology

Brennecke, Philip ; Rasina, Dace; Aubi, Oscar; Herzog, Katja; Lanskron, Johannes ; Cautain, Bastien; Vicente, Francisca; Quintana, Jordi ; Stechmann, Bahne; Brea, Jose

Total number of authors:
39

Published in:
SLAS DISCOVERY: Advancing Life Sciences R&D

Link to article, DOI:
[10.1177/2472555218816276](https://doi.org/10.1177/2472555218816276)

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Brennecke, P., Rasina, D., Aubi, O., Herzog, K., Lanskron, J., Cautain, B., Vicente, F., Quintana, J., Stechmann, B., Brea, J., Kolanowski, J., Pilarski, R., Orzaez, M., Pienda-Lucena, A., Laraia, L., Nami, F., Zielenkiewicz, P., Paruch, K., Hansen, E., ... Fecke, W. (2019). EU-OPENSOURCE: A novel collaborative approach to facilitate chemical biology. *SLAS DISCOVERY: Advancing Life Sciences R&D*, 24(3), 398-413. <https://doi.org/10.1177/2472555218816276>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



EU-OPENSREEN: A novel collaborative approach to facilitate chemical biology

Journal:	<i>SLAS Discovery</i>
Manuscript ID	DISC-18-0104.R1
Manuscript Type:	ASC Review
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Brennecke, Philip; Leibniz-Institut für Molekulare Pharmakologie, EU-OPENSREEN Rasina, Dace; Latvijas Organiskās sintēzes institūts, Organic Synthesis Methodology group Aubi, Oscar; University of Bergen, Department of Biomedicine Herzog, Katja; Leibniz-Institut für Molekulare Pharmakologie, EU-OPENSREEN Landskron, Johannes; University of Oslo, Centre for Molecular Medicine Norway, Nordic EMBL Partnership, Cautain, Bastien; Fundación MEDINA, Screening and target validation Vicente, Francisca; Fundación Medina / Centro de Excelencia en Investigación de Medicamentos Innovadores en Andalucía, Department of Screening Target and validation Quintana, Jordi; Universitat Pompeu Fabra, Department of Experimental and Health Sciences Mestres, Jordi; Universitat Pompeu Fabra, Department of Experimental and Health Sciences; IMIM Hospital del Mar Medical Research Institute, Research Program on Biomedical Informatics (GRIB) Stechmann, Bahne; Leibniz-Institut für Molekulare Pharmakologie, EU-OPENSREEN Brea, Jose; University of Santiago de Compostela, Pharmacology Kolanowski, Jacek; Institute of Bioorganic Chemistry - Polish Academy of Sciences, Department of Molecular Probes and Prodrugs Pilarski, Radosław; Institute of Bioorganic Chemistry - Polish Academy of Sciences, Department of Molecular Probes and Prodrugs Orzaez, Mar; Príncipe Felipe Research Center, Screening Platform Pineda-Lucena, Antonio; Health Research Institute Hospital La Fe, Drug Discovery Unit Laraia, Luca; Technical University of Denmark, Department of Chemistry; Technical University of Denmark, DK-OPENSREEN Nami, Faranak; Technical University of Denmark, Department of Chemistry; Technical University of Denmark, DK-OPENSREEN Zielenkiewicz, Piotr; Institute of Biochemistry and Biophysics - Polish Academy of Sciences, Department of Bioinformatics Paruch, Kamil; Masaryk University, Department of Chemistry - CZ-OPENSREEN Hansen, Espen; University of Tromsø, Marbio von Kries, Jens; FMP, Screening Unit Neuenschwander, Martin; Leibniz-Institut für Molekulare Pharmakologie, Screening Unit Specker, Edgar; Leibniz-Institut für Molekulare Pharmakologie, Medicinal Chemistry</p>

	<p>Bartunek, Petr; Institute of Molecular Genetics, Center for Chemical Genetics & CZ-OPENSREEN</p> <p>Simova, Sarka; Institute of Molecular Genetics, Center for Chemical Genetics & CZ-OPENSREEN</p> <p>Leśnikowski, Zbigniew ; Institute of Medical Biology - Polish Academy of Sciences, Laboratory of Molecular Virology and Biological Chemistry</p> <p>Krauss, Stefan ; Oslo universitetssykehus Ulleval, Department of Immunology and Transfusion Medicine; University of Oslo, Hybrid Technology Hub - Centre of Excellence - Institute of Basic Medical Sciences</p> <p>Lehtiö, Lari; University of Oulu, Faculty of Biochemistry and Molecular Medicine;</p> <p>Bilitewski, Ursula; Helmholtz-Zentrum für Infektionsforschung GmbH, Working Group Compound Profiling and Screening</p> <p>Broenstrup, Mark; Helmholtz-Zentrum für Infektionsforschung GmbH, Chemical Biology; German Center for Infection Research (DZIF), Partner Site Hannover-Brunswick</p> <p>Taskén, Kjetil; University of Oslo, Biotechnology Centre; University of Oslo and Oslo University Hospital, Centre for Molecular Medicine Norway, Nordic EMBL Partnership; Oslo universitetssykehus Ulleval, Department of Cancer Immunology - Institute for Cancer Research; University of Oslo, K.G. Jebsen Centre for Cancer Immunotherapy and K.G. Jebsen Centre for B cell malignancies- Institute of Clinical Medicine</p> <p>Jirgensons, Aigars ; Latvijas Organiskās sintezes institūts, Organic Synthesis Methodology group</p> <p>Lickert, Heiko; Helmholtz Centre Munich German Research Center for Environmental Health, Institute of Diabetes and Regeneration Research</p> <p>Clausen, Mads; Technical University of Denmark, Department of Chemistry; Technical University of Denmark , DK-OPENSREEN</p> <p>Andersen, Jeanette; University of Tromsø , Marbio</p> <p>Vicent, Maria; Principe Felipe Research Center, Screening Platform</p> <p>Genilloud, Olga; Fundación Medina / Centro de Excelencia en Investigación de Medicamentos Innovadores en Andalucía, Department of Screening Target and validation</p> <p>Martinez, Aurora; Universitetet i Bergen Det medisinsk-odontologiske fakultet, Biomedicine</p> <p>Nazaré, Marc; Leibniz-Institut für Molekulare Pharmakologie, Medicinal Chemistry</p> <p>Gibbon, Philip; Fraunhofer Institute for Molecular Biology and Environmental Ecology, ScreeningPort</p> <p>Fecke, Wolfgang; EU-OPENSREEN, Central Office</p>
Keywords:	Screening, Medicinal Chemistry, Chemical Biology, Open Access, Compound Library
Abstract:	<p>Compound screening in biological assays and subsequent optimization of hits is indispensable for the development of new molecular research tools and drug candidates. To facilitate such discoveries, the European Research Infrastructure EU-OPENSREEN was founded recently with support of its member countries and the European Commission. Its distributed character harnesses complementary knowledge, expertise and instrumentation in the discipline of chemical biology from 20 European partners, and its open working-model ensures that academia and industry can readily access EU-OPENSREEN's compound collection, equipment and generated data. To demonstrate the power of this collaborative approach, this review highlights recent projects from EU-OPENSREEN partner institutions. These studies yielded (i) 2-aminoquinazolin-4(3H)-ones as potential lead structures for new antimalarial drugs; (ii) a novel lipodepsipeptide specifically inducing apoptosis in cells deficient for the pVHL tumor suppressor; (iii) small molecule-based ROCK inhibitors that induce definitive endoderm formation and can potentially be used for regenerative medicine; (iv)</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	potential pharmacological chaperones for inborn errors of metabolism and a familiar form of acute myeloid leukemia (AML); and (v) novel tankyrase inhibitors which entered a lead-to-candidate program. Collectively, these findings highlight the benefits of small molecule screening, the plethora of assay designs, and the close connection between screening and medicinal chemistry within EU-OPENSREEN.



Color coding in revised manuscript file

Added/edited text to respond to Reviewers/Editors comments

References were deleted

References were added

Title

EU-OPENSREEN: A novel collaborative approach to facilitate chemical biology

Authors

Philip Brennecke¹, Dace Rasina², Oscar Aubi³, Katja Herzog¹, Johannes Landskron⁴, Bastien Cautain⁵, Francisca Vicente⁵, Jordi Quintana⁶, Jordi Mestres^{6,7}, Bahne Stechmann¹, Bernhard Ellinger⁸, Jose M. Brea⁹, Jacek L. Kolanowski¹⁰, Radosław Pilarski¹⁰, Mar Orzaez¹¹, Antonio Pineda-Lucena¹², Luca Laraia^{13,14}, Faranak Nami^{13,14}, Piotr Zielenkiewicz¹⁵, Kamil Paruch¹⁶, Espen Hansen¹⁷, Jens P. von Kries¹⁸, Martin Neuenschwander¹⁸, Edgar Specker¹⁹, Petr Bartunek²⁰, Sarka Simova²⁰, Zbigniew Leśnikowski²¹, Stefan Krauss^{22,23}, Lari Lehtiö²⁴, Ursula Bilitewski²⁵, Mark Brønstrup^{26,27}, Kjetil Taskén^{4,28,29,30}, Aigars Jirgensons², Heiko Lickert³¹, Mads H. Clausen^{13,14}, Jeanette H. Andersen¹⁷, Maria J. Vicent¹¹, Olga Genilloud⁵, Aurora Martinez³, Marc Nazaré¹⁹, Wolfgang Fecke^{32*}, Philip Gribbon^{8*}

Affiliations

¹Leibniz Research Institute for Molecular Pharmacology, EU-OPENSREEN, Berlin, Germany

²Latvian Institute of Organic Synthesis, Organic Synthesis Methodology group, Riga, Latvia

³University of Bergen, Department of Biomedicine, Bergen, Norway

⁴University of Oslo, Centre for Molecular Medicine Norway - Nordic EMBL Partnership, Oslo, Norway

1
2
3 ⁵Fundación MEDINA, Health Sciences Technology Park, Granada, Spain
4
5

6 ⁶Universitat Pompeu Fabra, Department of Experimental and Health Sciences, Barcelona, Spain
7
8

9 ⁷IMIM Hospital del Mar Medical Research Institute, Research Program on Biomedical Informatics
10
11 (GRIB), Barcelona, Spain
12
13

14 ⁸Fraunhofer Institute for Molecular Biology and Applied Ecology IME, Screening Port, Hamburg,
15
16 Germany
17
18

19 ⁹University of Santiago de Compostela, Institute for Research in Molecular Medicine and Chronic
20
21 Diseases - BioFarma Research Group, Santiago de Compostela, Spain
22
23

24 ¹⁰Institute of Bioorganic Chemistry - Polish Academy of Sciences, Department of Molecular
25
26 Probes and Prodrugs, Poznan, Poland
27
28
29

30 ¹¹Principe Felipe Research Center, Screening Platform, Valencia, Spain
31
32

33 ¹²Health Research Institute Hospital La Fe, Drug Discovery Unit, Valencia, Spain.
34
35

36 ¹³Center for Nanomedicine and Theranostics, Department of Chemistry, Technical University of
37
38 Denmark, Lyngby, Denmark
39
40

41 ¹⁴Technical University of Denmark, DK-OPENSREEN, Lyngby, Denmark
42
43

44 ¹⁵Institute of Biochemistry and Biophysics - Polish Academy of Sciences, Department of
45
46 Bioinformatics, Warsaw, Poland
47
48

49 ¹⁶Masaryk University, Department of Chemistry - CZ-OPENSREEN, Brno, Czech Republic
50
51

52 ¹⁷University of Tromsø – The Arctic University of Norway, Marbio, Tromsø, Norway
53
54

55 ¹⁸Leibniz Research Institute for Molecular Pharmacology, Screening Unit, Berlin, Germany
56
57
58
59
60

1
2
3 ¹⁹Leibniz Research Institute for Molecular Pharmacology, Medicinal Chemistry Research Group,
4
5 Berlin, Germany
6

7
8 ²⁰Institute of Molecular Genetics of the ASCR, CZ-OPENSREEN, Prague, Czech Republic
9

10
11 ²¹Institute of Medical Biology - Polish Academy of Sciences, Laboratory of Molecular Virology
12
13 and Biological Chemistry, Łódź, Poland
14
15

16
17 ²²Oslo University Hospital, Department of Immunology and Transfusion Medicine, Oslo, Norway
18

19
20 ²³University of Oslo, Hybrid Technology Hub - Centre of Excellence - Institute of Basic Medical
21
22 Sciences, Oslo, Norway
23

24
25 ²⁴University of Oulu, Faculty of Biochemistry and Molecular Medicine - Biocenter Oulu, Oulu,
26
27 Finland
28

29
30 ²⁵Helmholtz Centre for Infection Research, Working Group Compound Profiling and Screening,
31
32 Brunswick, Germany
33

34
35 ²⁶Helmholtz Centre for Infection Research, Department of Chemical Biology, Brunswick,
36
37 Germany
38

39
40 ²⁷German Center for Infection Research (DZIF), partner site Hannover-Brunswick, Brunswick,
41
42 Germany
43

44
45 ²⁸Oslo University Hospital, Department of Cancer Immunology - Institute for Cancer Research,
46
47 Oslo, Norway
48

49
50 ²⁹University of Oslo, K.G. Jebsen Centre for Cancer Immunotherapy - Institute of Clinical
51
52 Medicine, Oslo, Norway
53
54

1
2
3 ³⁰University of Oslo, K.G. Jebsen Centre for B cell malignancies - Institute of Clinical Medicine,
4
5 Oslo, Norway
6
7

8 ³¹Helmholtz Centre Munich German Research Center for Environmental Health, Institute of
9
10 Diabetes and Regeneration Research, Neuherberg, Germany
11
12

13 ³²EU-OPENSREEN, Central Office, Berlin, Germany
14
15

16 17 **Corresponding Authors:**

18
19 *Philip Gribbon, Fraunhofer Institute for Molecular Biology and Applied Ecology IME, Screening
20
21 Port, Schnackenburgallee 114, 22525 Hamburg, Germany, Email:
22
23 Philip.Gribbon@ime.fraunhofer.de
24
25

26
27 *Wolfgang Fecke, EU-OPENSREEN, Central Office, Robert-Roessle-Strasse 10, 13125 Berlin,
28
29 Germany, Email: wolfgang.fecke@eu-openscreen.eu
30
31

32 33 **Keywords**

34
35 Chemical biology, screening, medicinal chemistry, open access, compound library
36
37

38 39 **Abstract**

40
41 Compound screening in biological assays and subsequent optimization of hits is indispensable for
42
43 the development of new molecular research tools and drug candidates. To facilitate such
44
45 discoveries, the European Research Infrastructure EU-OPENSREEN was founded recently with
46
47 support of its member countries and the European Commission. Its distributed character harnesses
48
49 complementary knowledge, expertise and instrumentation in the discipline of chemical biology
50
51 from 20 European partners, and its open working-model ensures that academia and industry can
52
53 readily access EU-OPENSREEN's compound collection, equipment and generated data. To
54
55
56
57
58
59
60

1
2
3 demonstrate the power of this collaborative approach, this review highlights recent projects from
4 EU-OPENSSCREEN partner institutions. These studies yielded (i) 2-aminoquinazolin-4(3H)-ones
5 as potential lead structures for new antimalarial drugs; (ii) a novel lipodepsipeptide specifically
6 inducing apoptosis in cells deficient for the pVHL tumor suppressor; (iii) small molecule-based
7 ROCK inhibitors that induce definitive endoderm formation and can potentially be used for
8 regenerative medicine; (iv) potential pharmacological chaperones for inborn errors of metabolism
9 and a familiar form of acute myeloid leukemia (AML); and (v) novel tankyrase inhibitors which
10 entered a lead-to-candidate program. Collectively, these findings highlight the benefits of small
11 molecule screening, the plethora of assay designs, and the close connection between screening and
12 medicinal chemistry within EU-OPENSSCREEN.
13
14
15
16
17
18
19
20
21
22
23
24
25

26 **Scientific concept and operational model of EU-OPENSSCREEN**

27
28
29
30 Chemical biology is an interdisciplinary research field, which has emerged in the past decade from
31 classical pharmacology and cell biology, and it studies the effects of chemical compounds on
32 biological systems. In parallel, post-genome biology with its powerful technologies of genome
33 sequencing, transcriptomics, proteomics, metabolomics and genome editing by CRISPR/Cas9 as
34 well as new model systems such as, for example, microphysiological systems have provided a
35 rapidly expanding range of information on new cellular targets for basic research and early drug
36 discovery. However, the availability of selective ‘tools or probes’ for systematic biochemical
37 investigation of target function still remains a limiting factor in many investigations. Reviews on
38 the use of chemical compounds that elicit a well-defined biological effect, highlight the
39 opportunities (and also the limitations) of these tool compounds for the modulation of functions
40 of biological targets and for studying the underlying molecular mechanisms at the biochemical,
41 cellular, tissue and organismal levels¹. Chemical biology as a discipline is well represented in, but
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 not restricted to, academia, and increasingly, the drug discovery workflows in pharmaceutical
4 companies feature chemical biology studies. Discovery and optimization of probe compounds
5 form important parts of industrial tractability and “Go versus No-Go” decision-making on novel
6 targets as well as the study of target translatability as part of the early design of subsequent clinical
7 phases².
8
9

10
11
12
13
14
15 EU-OPENSREEN is a Research Infrastructure (RI) of screening and medicinal chemistry
16 platforms that was established by seven European member states and one observer state in 2018,
17 under the legal framework of a European Research Infrastructure Consortium (ERIC), in order to
18 enable chemical biology in an open-access setting (www.eu-openscreen.eu). The RI will provide
19 scientists access to a chemical library, assay development and screening facilities, medicinal
20 chemistry and informatics platforms as well as associated supporting facilities for protein
21 production, cell line generation (e.g., 2D and 3D models and patient-derived cells), computational
22 and structural biology as well as structure-based drug design (**Figure 1**). As it begins the
23 operational phase of its work, EU-OPENSREEN anticipates that a high demand exists from users
24 to access its platforms. In the year before initiation of EU-OPENSREEN, the partner sites which
25 came together to form the EU-OPENSREEN network participated in more than 135 screening
26 and medicinal chemistry projects, involving academic, **Small and Medium-sized Enterprise (SME)**
27 and large industry users coming from 22 European and non-European countries. Historically, the
28 partner sites of EU-OPENSREEN have been involved in multiple projects, which reached
29 clinical and/or pre-clinical stages for indications including leukemia, epilepsy, autism, myotonic
30 dystrophy, Parkinson’s disease and colon cancer.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51
52
53 The central office of the EU-OPENSREEN ERIC acts as a single point of contact for users,
54 assists in identifying the appropriate partner site and helps to formulate the project design. **Herein,**
55
56
57
58
59
60

1
2
3 the central office receives the project requests from users and contacts potential partners with
4 appropriate technologies and expertise. Together with the partner sites, the technical feasibility,
5 the scientific novelty of the proposed project and the most suitable partner site is evaluated, and,
6 if deemed feasible and novel, the project will be initiated in collaboration with the most suitable
7 EU-OPENSREEN partner. Depending on the individual needs of the users, projects take place
8 at one of 15 specialized and high capacity screening partner sites or 6 medicinal chemistry sites.
9
10 The assay development phase preceding the actual screening campaign emphasizes the quality,
11 integrity, robustness and pharmacological relevance of the readouts. Projects have access to the
12 EU-OPENSREEN compound collection³ containing 100,000 commercially available as well as
13 40,000 academic compounds crowdsourced through a network of national chemical biology
14 networks. A 5,000-compound subset—carefully selected to represent the diversity of the entire
15 collection and also containing 2,500 known bioactive compounds—is available for the process of
16 assay validation in pilot studies. Notably, during the selection process for commercial compounds
17 pan-assay interference compounds (PAINS)⁴ are partially selected against. Furthermore, access
18 to a wider range of chemical diversity will be provided through the isolation and characterization
19 of natural products by some of the EU-OPENSREEN partner sites, and, moreover, a fragment
20 library for fragment-based screening will be acquired. These two additions are currently the only
21 expansions envisioned for the EU-OPENSREEN compound collection, and future expansions
22 will depend on the funding situation. The compound collection is stored and managed at the
23 headquarters located in Berlin, and copies of the screening collection, as well as library updates,
24 are distributed to the EU-OPENSREEN partner sites. Results from the primary screens will be
25 made available globally in an open access database. To allow for publication of data or securing
26 intellectual property a grace period of up to three years is provided between completion of the

1
2
3 primary screen and data presentation in the EU-OPENSREEN database. After the official start
4 of EU-OPENSREEN in 2018, the database is currently still under construction. A link will be
5 provided on the EU-OPENSREEN website as soon as the database development has been
6 completed.
7
8
9
10
11

12 While EU-OPENSREEN just started its operational phase, its 20 partner institutions are already
13 operational as standalone institutes. Hence, in the following sections, recent successful chemical
14 biology projects at selected partner institutes are presented to give an overview of the capacities
15 and expertise present within the EU-OPENSREEN network as well as an outlook as to how EU-
16 OPENSREEN can facilitate academic and applied science in the future.
17
18
19
20
21
22
23
24

25 **Project A: 2-Aminoquinazolin-4(3H)-ones as potential new nonpeptidomimetic lead** 26 **structures for antimalarial agents** 27 28 29

30 Emerging drug resistance of *Plasmodium* parasites against approved antimalarial drugs represents
31 a major threat for preventing and treating malaria, one of the most detrimental infectious diseases,
32 in the future⁵. Hence, substantial efforts are currently being made to identify novel drug candidates,
33 which interrupt the life cycle of the parasite by yet unknown mechanisms of action. In this context,
34 the Latvian Institute of Organic Synthesis, a medicinal chemistry partner site of EU-
35 OPENSREEN, recently spearheaded a study that identified 2-aminoquinazolin-4(3H)-ones as
36 potent inhibitors of three digestive aspartic proteases (i.e., digestive plasmepsins), namely Plm I,
37 II, and IV⁶. Digestive plasmepsins (i.e., Plm I, II, IV and HAP), localized in the digestive vacuole
38 of the *Plasmodium* parasite, contribute towards the metabolism of hemoglobin into amino acids,
39 and represent a subgroup of the extensively studied family of aspartic plasmepsin proteases, which
40 have been considered as potential drug targets against malaria parasites for more than ten years⁷⁻⁹.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 Due to their high potential as drug targets, numerous inhibitors have been developed against
57
58
59
60

1
2
3 plasmepsins over the past years¹⁰⁻¹². Nonpeptidomimetic compounds show better selectivity
4
5 against human aspartic proteases, and tend to bind to the open-flap form of *Plasmodium*
6
7 plasmepsins, whereas peptidomimetic compounds preferably bind to the closed-flap form of the
8
9 proteases. Herein, opening of the flap results in accessibility of a protein subpocket, which can
10
11 subsequently be occupied by parts of the nonpeptidomimetic inhibitor. In close collaboration with
12
13 the Biomedical Research and Study Centre in Riga and the Francis Crick Institute in London, an
14
15 NMR-based fragment screening approach^{13, 14} was used to identify 2-aminoquinazolin-4(3H)-ones
16
17 as nonpeptidomimetic inhibitors of digestive plasmepsins (**Figure 2a**). To this end, a diversity
18
19 subset comprising 976 Astex “Rule of 3”-compliant compounds of the ChemBridge fragment
20
21 collection was screened against binding to the digestive plasmepsin Plm II. STD, T1 ρ and Water-
22
23 LOGSY NMR spectra were taken in both, the presence and absence of the Plm II target, and
24
25 specificity was assessed by competition experiments with the potent aspartic protease inhibitor
26
27 pepstatin A¹⁵. Binding to Plm II detected by at least two of the three NMR techniques as well as
28
29 competition with pepstatin A were set as hit criteria, and yielded a total of 49 fragment hits.
30
31 Subsequent FRET-based enzymatic assays identified fragment **1**, which is based on the 2-
32
33 aminoquinazolin-4(3H)-one scaffold (**Figure 2a**), as the most potent hit with one of the highest
34
35 ligand efficiencies ($IC_{50}(\text{Plm II}) = 24.3 \mu\text{M}$, $LE = 0.35$). Subsequent chemical optimization
36
37 experiments were thus performed using this scaffold as a starting point. An initial hint for the
38
39 medicinal chemistry optimization strategy came from NMR competition experiments using
40
41 fragment **1** in combination with fragment **2**: The two fragments did not bind to the same binding
42
43 site, and an observed interligand NOE between H-7 of fragment **1** and H-5 of fragment **2** indicated
44
45 close proximity of the two fragments – thus implying an additional binding pocket around H-7 of
46
47 fragment **1** (**Figure 2b**). Following this hint, the addition of a phenyl group at position 7 of
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 fragment **1** yielded more potent inhibitor *rac-3a* ($IC_{50}(\text{Plm II}) = 2.3 \mu\text{M}$, $LE = 0.32$) (**Figure 2b**).
4
5 Subsequent docking experiments with *rac-3a* using Schrödinger Glide software¹⁶ suggested high
6
7 quality binding to the open-flap conformation of Plm II with hydrogen bonding interactions
8
9 between the N1 and 2-amino group of fragment **1** and catalytic Asp34-Asp214 dyad and indicated
10
11 an extension of the added phenyl residue toward the deep flap pocket, while the tetrahydrofuran
12
13 (THF) moiety of fragment **1** resided in the S1' pocket of Plm II. Moreover, docking experiments
14
15 suggested additional space in S1', and indeed, adding a lipophilic cis-phenyl group to the 5-
16
17 position of the THF group yielded *rac-3b* as more potent inhibitor of Plm II ($IC_{50}(\text{Plm II}) = 0.57$
18
19 μM , $LE = 0.28$) with a more than 10-fold selectivity for Plm II when compared to human Cat D
20
21 (**Figure 2b**). A co-crystal of *rac-3b* and Plm II was obtained, and the X-ray structure (2.7 Å)
22
23 confirmed binding of the inhibitor to the open-flap conformation previously suggested by the
24
25 docking experiments and thus provided a better understanding of the binding of the inhibitor to
26
27 Plm II (**Figure 2c**). Importantly, the X-ray structure also showed that the flap pocket is for the
28
29 most part unoccupied by *rac-3b*, and thus suggested that adding hydrophobic groups to the 7-
30
31 phenyl substituent could help targeting the flap-pocket; and indeed, literature-informed^{10, 11, 17}
32
33 addition of *n*-pentyl group to the para position of the phenyl group yielded the even more potent
34
35 inhibitor *rac-4a* ($IC_{50}(\text{Plm II}) = 0.15 \mu\text{M}$, $LE = 0.27$) (**Figure 2d**). Notably, adding a 3-
36
37 phenylpropyl group to the para position of the phenyl group yielded the Plm IV specific inhibitor
38
39 *rac-4b* ($IC_{50}(\text{Plm IV}) = 0.13 \mu\text{M}$, $LE = 0.17$) (**Figure 2d**), whereas *rac-4a* inhibited the three
40
41 digestive plasmepsins Plm I, II and IV to similar extents. Selectivity over human Cat D was
42
43 comparable to previous reports on nonpeptidomimetic Plm inhibitors for both *rac-4a* and *rac-4b*¹⁰.
44
45 Interestingly, homology modeling of Plm IV using Plm II as a template suggested that the flap
46
47 pocket of Plm IV is slightly more spacious. Both structure models could accommodate the smaller
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *n*-pentyl group of *rac*-**4a** well, while the bulkier 3-phenylpropyl group of *rac*-**4b** could only fit
4 into the more spacious flap pocket of Plm IV. This suggests that specifically targeting the flap
5 pockets of digestive plasmepsins could yield additional subtype-specific inhibitors. Both *rac*-**4a**
6 and *rac*-**4b** were tested in an *in vitro* growth assay using the *P. falciparum* clone 3D7, and showed
7 growth inhibition at levels close to 1 μM (*rac*-**4a**: $1.1 \pm 0.2 \mu\text{M}$, *rac*-**4b**: $1.2 \pm 0.2 \mu\text{M}$), which is
8 considered as a criteria for promising leads for potential new antimalarial drugs¹⁰.
9

10
11
12
13
14
15
16
17
18 This study nicely demonstrates several aspects that are also relevant in the context of the EU-
19 OPENSREEN network. Firstly, the EU-OPENSREEN compound collection will include a
20 fragment-based subset of compounds, and the Latvian Institute of Organic Synthesis (LIOS) will
21 be involved in the design and setup of this library subset. Secondly, the expertise in fragment-
22 based drug discovery present at LIOS represents an important addition to the EU-OPENSREEN
23 capabilities, and future EU-OPENSREEN users designing similar experimental setups will find
24 LIOS to be a competent collaboration partner for running their experiments. Lastly, the study
25 shows that EU-OPENSREEN covers expertise regarding assay systems, screening as well as
26 subsequent chemical optimization of hit compounds. This will allow prospective users to identify
27 promising hits and develop them into lead structures for further drug discovery. Here, the
28 complementary expertise of currently 20 partner institutions ensures that prospective users will be
29 directed to the collaborator who best matches their individual experimental needs.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 **Project B: Discovery of the novel lipodepsiptide MDN-0066, a natural product that**
47 **specifically induces apoptosis in cells that do not express the functional pVHL tumor**
48 **suppressor.**
49
50
51

52
53
54 Inactivation of the Von Hippel Lindau tumor suppressor gene (pVHL) occurs in many renal cell
55 carcinomas (RCCs) as well as in other cancers. In clear cell renal cell carcinomas (CC-RCC), for
56
57
58
59
60

1
2
3 instance, either genetic or epigenetic inactivation of pVHL function is observed in 70% to 80% of
4 reported cases^{18, 19}. pVHL is ubiquitously expressed throughout human tissues, and its loss of
5 expression is thus restricted to the tumor tissue. Inactivation is usually accompanied by high levels
6 of tumor vascularization and poor prognosis for the patient. Moreover, it has been shown that
7 reintroduction of functional pVHL into RCC cells restores tumor suppression²⁰. Collectively, its
8 high rate of inactivation in RCCs, the local restriction of pVHL inactivation to the tumor tissue,
9 and its important role for malignancy make pVHL an ideal target for novel drug candidates, which
10 will possibly have only few side effects in unaffected tissues due to the differential expression of
11 functional pVHL in healthy and diseased cells. To identify novel chemical agents that exhibit cell
12 toxicity only in the context of pVHL inactivation, the EU-OPENSSCREEN partner site Fundación
13 MEDINA used a quantitative whole cell assay—previously developed²¹⁻²³ on site²⁴, to screen
14 pVHL-deficient human renal carcinoma cells (RCC4 cell line) transformed either with an empty
15 vector (RCC4-VA) or with a vector expressing a functional rescue copy of pVHL (RCC4-VHL)
16 against a collection of microbial extracts²⁵. Herein, the decision to use natural products for the
17 screen was informed by the fact that microorganisms are capable of producing a vast variety of
18 diverse chemical structures, which makes them highly interesting for drug discovery, and explains
19 why natural products keep showing desirable bioactivities in various therapeutic areas²⁶. A total
20 of 1,040 microbial acetone extracts were isolated from 117 unicellular microbial species grown in
21 a set of different growth conditions. Microbial extracts that induced cell death in RCC4-VA but
22 not in RCC4-VHL cells were subsequently fractionated, fractions were rescreened, and the ones
23 recapitulating the RCC4-VA-specific induction of cell death were further analyzed by LC/HRMS
24 and NMR to identify the active natural product and assess its novelty by chemical dereplication.
25 These dereplication experiments allowed rapid focusing on active fractions of the promising strain
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 F 278,770^T, which eventually turned out to be a previously not described *Pseudomonas* species,
4 now named *P. granadensis sp. nov.*²⁷. Subsequent large-scale fermentation, assay-guided
5 compound isolation and structural elucidation eventually yielded the natural product MDN-0066,
6 a lipodepsipeptide, which was not described previously in the literature and, due to its unique
7 structure, established a novel class of lipodepsipeptides (**Figure 3a**). Dose response analysis
8 showed that the pVHL-deficient cell line RCC4-VA was more sensitive to purified MDN-0066
9 (IC₅₀=24.4 μM) as compared to the pVHL rescue cell line RCC4-VHL (IC₅₀=69.23 μM), and
10 demonstrated that MDN-0066 already induced cell death in RCC4-VA cells at concentrations that
11 did not show an effect in the rescue cell line (**Figure 3b**). Importantly, it could be shown that
12 MDN-0066 is capable of inducing apoptosis specifically in pVHL deficient cells as judged by
13 PARP cleavage detection. This was confirmed by Annexin V / PI staining, which demonstrated
14 specific induction of apoptosis in pVHL deficient cells upon MDN-0066 treatment (**Figure 3c**).
15 This finding is promising with respect to drug development programs, as pVHL-deficient cells
16 have previously been described as being resistant to apoptosis²⁰. Moreover, flow cytometric DNA
17 content analysis of pVHL deficient RCC4-VA and RCC4-VHS cells upon 48 h exposure to MDN-
18 0066 revealed an S phase arrest unique to RCC4-VA cells (6.0 ± 2.1 to 18.5% ± 5.2; p = 0.005),
19 which was correlated with an G0/G1 decrease (75.5 ± 4.2 to 37.5% ± 4.6; p = 0.001) and a G2/M
20 increase (10 ± 0.5 to 26.2% ± 3.1; p<0.0005) (**see Supplementary Figure S1**). These results
21 suggest that MDN-0066 is critical for the S phase arrest in RCC4-VA cells, but can also be
22 involved in G2/M arrest in the absence of pVHL. Loss of HIF-1α causes an increased progression
23 into S phase, rather than a growth arrest and pVHL is associated with cell cycle arrest upon serum
24 withdrawal²⁸. Collectively, these results suggest that MDN-0066 has an effect on the VHL/HIF
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 pathway, which is involved in cell cycle regulation. Ongoing research tries to establish the link
4
5 between the cell cycle arrest and the induction of apoptosis.
6
7

8 In sum, this report represents the first potential lead compound from a natural source affecting
9
10 cellular growth specifically in the absence of the pVHL tumor suppressor, and thus could have an
11
12 important impact on the future treatment of renal cancers.
13
14
15

16 One third of top selling drugs are based on natural products, and they are used successfully in the
17
18 treatment of numerous diseases²⁶. To adequately cover this highly diverse class of potential drug
19
20 candidates, EU-OPENSREEN will include a set of natural products in its compound collection.
21
22 Hence, having the expertise to isolate and characterize natural products represented in the network
23
24 is of high importance to ensure that EU-OPENSREEN satisfactorily serves the users' demand
25
26 for structural diversity. Moreover, this study demonstrates the competency of the EU-
27
28 OPENSREEN network to develop novel assays that can be easily adapted for use in high
29
30 throughput screening campaigns. The screening of 1,040 natural products represented just a
31
32 starting point, and future screening campaigns using larger collections of natural products,
33
34 including both pure compounds and extract/fraction libraries, are envisioned. The initial assay
35
36 design and setup, which involves screening of small library subsets, is of utmost importance for
37
38 screening larger compound collections successfully later on; and EU-OPENSREEN users can
39
40 rely on competent collaborators, assistance and advice during this crucial first step of the
41
42 experimental workflow.
43
44
45
46
47
48

49 **Project C: High content screening to enable regenerative medicine strategies in Diabetes**
50
51 **Mellitus**
52
53
54
55
56
57
58
59
60

1
2
3 Human embryonic stem cells (hESCs) are pluripotent and hence are able to generate cell and tissue
4 types of the three germ layers endoderm, mesoderm and ectoderm. In this project, the EU-
5 OPENSREEN high capacity screening site Fraunhofer IME Screening Port Hamburg worked
6 closely with colleagues in the HMGU Munich to identify small molecules able to direct efficient
7 generation of definitive endoderm (DE) and serve as a platform to investigate the regulatory
8 mechanisms of endodermal differentiation²⁹. Access to such endodermal cell lineages in turn gives
9 hope to treat diseases such as Diabetes Mellitus by means of transplantation therapies. As the main
10 challenge in the pancreatic transplantation therapy is the shortage of donor islets, generation and
11 expansion of functional β -cells from hESCs and human induced pluripotent stem cells (iPSCs)
12 could represent a promising approach³⁰. Current differentiation protocols act by mimicking the
13 developmental steps *in vitro* and have facilitated generation and upscaling of pancreatic β -cells³¹.
14 However, a major drawback of these protocols is the generation of immature β -cells compared to
15 human β -cells found in islets³². One tactic to increase the efficiency of β -cell generation is to target
16 specific pathways that are directly linked to insulin production in cultured cells.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 Previous data indicate that the transcription factor FoxA2 is highly expressed in mesendoderm and
37 definitive endoderm at the early stages of differentiation and is also expressed in all the endoderm
38 derived organs³³. To facilitate high throughput screening of novel compounds that might induce
39 definitive endoderm, a mouse embryonic stem cell (mESC) line was used for primary screening.
40
41
42
43
44

45 The mESC cell line was selected because it was relatively straightforward to perform large scale
46 cell culture needed to support screening and the line was previously well characterized with respect
47 to signaling pathways involved in controlling endodermal differentiation³³. The endoderm
48 differentiation protocol used in the screen was based on previously published protocols with some
49 modifications to the basal medium to achieve higher levels of FoxA2 expression in the mESCs in
50
51
52
53
54
55
56
57
58
59
60

1
2
3 order to meet the needs of high throughput assays. Under the influence of TGF- β pathway factors
4 such as Activin A, the cells differentiate successfully into FoxA2-positive cells while the
5 undifferentiated stem cells show strong expression of the pluripotency marker Oct4²⁹.
6
7

8
9
10 The initial high throughput screen was performed in 384-well format with 23,406 small molecules
11 (Enamine, Monmouth Jct., NJ), which were selected on the basis of chemical diversity and which
12 excluded structural features associated with well described interference effects⁴. The compounds
13 were lead-like with a cLogP of < 4.2 and a molecular weight between 180 and 460 Da. In addition,
14 an FDA approved drug set from Enzo (Lörrach, Germany) was also screened both with the aim to
15 identify possible targets for active compounds and, in the longer term, for the possibility of drug
16 repurposing.
17
18

19
20 For the primary assay, Perkin Elmer Cell Carrier assay plates were coated with autoclaved 0.1%
21 gelatin, and compounds as well as controls were spotted by an echo liquid handling system
22 (Labcyte Echo 550) into coated 384-well plates (source compounds at 2 mM in 100 % DMSO).
23 Wells with the differentiation control contained Activin A and Wnt3a added (final concentrations
24 of 12.5 ng/ml (Activin A, R+D systems)) and 2 ng/ml (Wnt3a, R+D systems)). For the
25 pluripotency control, a GSK3 β inhibitor (CT 99021, Axon Medchem) was added to a final
26 concentration of 3 μ M. Negative controls were generated by adding only DMSO. Seeding of
27 mESCs involved addition of 50 μ l of mESC suspension to give between 11,000 – 14,000 cells per
28 well. Medium change with re-addition of fresh compound and control solutions occurred on day 2
29 post seeding, and cells were then fixed and prepared for imaging at day 5 after seeding. Cells were
30 permeabilized, blocked and primary [anti-FoxA2 goat polyclonal antibody (sc-6554, Santa Cruz
31 Biotechnology) and anti-Oct-3/4 mouse monoclonal antibody (sc-5279, Santa Cruz
32 Biotechnology)] and secondary [Alexa-Fluor 555 anti-mouse IgG and Alexa-Fluor 488 donkey
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 anti-goat] antibodies were added under automated screening conditions. The image acquisition
4 procedure used a Perkin Elmer Opera Imaging System with DNA-DAPI staining. Image analysis
5 used the features of the Columbus suite three Columbus Image Data Storage and Analysis System
6 (Perkin Elmer, Waltham, MA), which is a tool to access, store and explore images. A multiple step
7 image analysis sequence was developed (**Figure 4A**), and the DMSO control (**Figure 4B**),
8 pluripotency control (**Figure 4C**) and differentiation control (**Figure 4D**) were analyzed
9 automatically to determine the relative response to compounds in terms of capacity to induce both
10 proliferation and FoxA2 expression in mESCs.
11
12
13
14
15
16
17
18
19
20
21

22 For each phenotype, compounds which gave a minimum of 20% elevation in mean FoxA2 signal
23 intensity relative to solvent control treated cells were selected as “differentiation-inducing hits”.
24 A total of 400 primary differentiation-inducing hits from the primary screen were then tested in
25 triplicates and from these, 67 compounds were identified for follow-up in dose response
26 experiments. In addition, a further hit expansion was performed by selecting analogues of the 67
27 compounds from a set of 200,000 Enamine compounds, which were available as liquid stocks.
28 Around three members of each cluster were selected to give 351 compounds for analysis in a
29 seven-point dose response with serial compound dilutions and cell viability assessment to account
30 for cytotoxicity using the Cell-Titer-Glo Luminescent Assay (Promega). Based on the
31 differentiation controls as well as cell viability, the ROCK inhibitor Fasudil was selected for
32 further studies to induce definitive endoderm *in vitro* in an analogue-by-catalogue approach. Out
33 of the 16 analogues of Fasudil tested, two analogues gave rise to endoderm differentiation (with
34 27% and 31% of FoxA2+ cells respectively), which were comparable to the induction of FoxA2
35 by Fasudil (25.5%) and Wnt3a/Activin A (23%). **In further profiling studies using the human ESC**
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 line H9, the capacity of Fasudil to induce DE expression was confirmed by FACS and qPCR
4 analyses²⁹.
5
6

7
8 This project shows the potential of chemical biology approaches for elucidating signaling
9 pathways associated with cellular differentiation³⁴⁻³⁶. Small molecule-based-ROCK inhibition can
10 play an important role in early lineage formation of stem cells. Together with low batch-to-batch
11 variation compared to the recombinant proteins normally used for DE induction, the ROCK
12 inhibitor Fasudil could prove to be a very useful small molecule tool to promote differentiation of
13 ESCs towards definitive endoderm, and therefore in the longer term act as a contributor to
14 regenerative medicine approaches based upon pancreatic transplantation therapy. Interestingly, at
15 the CIPF EU-OPENSREEN partner site in Spain, it was recently also demonstrated that ROCK
16 inhibitors could have an impact not only on pancreatic cell transplantation therapies but also on
17 transplantation of ependymal stem/progenitor cells (epSPCs) of the spinal cord in acute as well as
18 chronic *in vivo* models of spinal cord injury (SCI)³⁷. In the context of EU-OPENSREEN, the
19 expertise in stem cell-based assays in conjunction with high throughput and high content
20 screening, present at the Fraunhofer IME Screening Port as well as the CIPF partner sites,
21 represents an important asset of EU-OPENSREEN to provide new tool compounds to support
22 the rapidly growing field of regenerative medicine.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 **Project D: Pharmacological chaperones to correct the instability of mutants associated with** 44 **inborn errors of metabolism and rare forms of cancer** 45

46 Many inherited disorders are associated with defective anabolic or catabolic pathways in cell
47 metabolism, affecting the synthesis, degradation and storage of biomolecules. These disorders,
48 collectively known as inborn errors of metabolism (IEM), are typically caused by mutations in
49 enzyme-coding genes and, although each of these disorders is rare, the combined incidence of IEM
50
51
52
53
54
55
56
57
58
59
60

1
2
3 is approximately 1 in 2,000 births worldwide, and new diseases continue to be recognized³⁸.
4
5 Nevertheless, only few IEMs can be treated effectively, and developing new therapies has proven
6
7 to be challenging. Increasing knowledge on the pathogenic mechanisms has revealed that many
8
9 IEM-associated mutations lead to destabilization and misfolding of the coded variants. This link
10
11 between mutation-associated destabilization and loss-of-function has been largely demonstrated
12
13 for phenylketonuria (PKU)³⁹, which is the most prevalent IEM, caused by mutations in
14
15 phenylalanine hydroxylase (PAH) that lead to neurotoxic levels of phenylalanine. Despite complex
16
17 quality control systems, the folding of unstable mutant proteins is not properly achieved, and the
18
19 consequent misfolding may cause increased protein degradation, aggregation and/or
20
21 mislocalization in the cell. This understanding points to pharmacological chaperones (PC) as
22
23 promising therapeutic strategies for IEM correction, earlier shown as a proof-of-concept for
24
25 PKU⁴⁰. PCs are small compounds that stabilize and rescue variant proteins by stimulating their
26
27 renaturation and preventing their misfolding *in vitro* and *in vivo*. The EU-OPENSREEN partner
28
29 site at the University of Bergen, Norway, is specialized in target-based biophysical screens, and
30
31 has developed concepts and a protocol for the discovery and development of PCs.
32
33
34
35
36
37

38 This protocol includes a primary screening of chemical libraries, searching for stabilizing binders
39
40 of target proteins by differential scanning fluorimetry (DSF)⁴¹. Customarily, DSF is performed in
41
42 a Real-Time PCR instrument with 384-well microplates and total assay volume of 25 μ L in each
43
44 well consisting of the protein sample at concentrations in the range 0.05-0.15 mg/mL in an optimal
45
46 buffer, and the extrinsic dye SYPRO Orange that emits fluorescence when interacting with
47
48 hydrophobic areas of denatured proteins (**Figure 5A**). The compounds are added to the assay
49
50 solution to a final concentration of 80 μ g/mL and 4% DMSO. Negative controls with only 4%
51
52 DMSO are routinely included on each plate. The unfolding curves are then registered from 20 °C
53
54
55
56
57
58
59
60

1
2
3 to 95 °C at a 2 °C/min scan rate and the midpoint melting temperature (T_m) and the corresponding
4 shift relative to the reference (ΔT_m) is calculated for each compound (**Figure 5A**). Thus,
5
6
7
8 compounds with significant increases in ΔT_m are selected using in-house software and those
9
10 showing adequate follow-up concentration-dependent DSF curves are considered primary
11 stabilizing hits. Further selection requires secondary assays to corroborate binding and protein-
12 target stabilization by several methods, preferentially surface plasmon resonance (SPR), which
13
14 also provides the dissociation constant (K_D) for the primary hits. Additional secondary assays
15
16 include enzymatic activity measurements to eliminate inhibitory compounds. Finally, the efficacy
17
18 of the best compounds is validated in tests measuring the increased activity and steady state levels
19
20 of the target protein in eukaryotic cells and animal models (Aubi et al. 2018; In press) (**Figure**
21
22
23
24
25
26
27 **5B**). The partner site at the University of Bergen has used this protocol in the early stage discovery
28 and validation of compounds for conformational stabilization of specific enzymes associated with
29
30 IEMs⁴²⁻⁴⁵. Methylmalonic aciduria cblB type (MMA cblB) is caused by the impairment of
31
32 ATP:cob(I)alamin adenosyltransferase (ATR), the enzyme that catalyzes the synthesis of
33
34 adenosylcobalamin. ATR was screened for the first time for small molecular weight stabilizing
35
36 ligands, and six potential PCs were discovered from the MyriaScreen Diversity Collection from
37
38 Sigma Aldrich/TimTec (Newark, DE), of which compound V (N-{{(4-
39
40 chlorophenyl)carbamothioyl]amino}-2-phenylacetamide) was very effective in increasing the
41
42 stability and activity of common MMA cblB-associated ATR-variants overexpressed in a cellular
43
44 model, as well as in mice^{43, 44}. The K_D -value for binding of compound V to ATR was 7.4 ± 0.4
45
46 μM , showing a relative high affinity for a primary hit. Molecular docking for compound V
47
48 identified a probable binding site surrounded by residues 228–240 and 165–175 at each subunit in
49
50 the ATR tetramer (**Figure 5C**). Recently, a similar DSF-based screening has been applied to
51
52
53
54
55
56
57
58
59
60

1
2
3 discover compounds that improve the conformational defect of mutant nucleophosmin (NPM).
4
5 This is a multifunctional nucleolar protein where deletion mutations at Trp288 and Trp290, located
6
7 in the C-terminal domain, lead to cytoplasmic localization and aggregation, features that are
8
9 associated with acute myeloid leukemia (AML)⁴⁶. Nucleophosmin (NPM) is an established target
10
11 in cancer, and as such has been the subject of several screening and medicinal chemistry
12
13 campaigns; most of this activity is reviewed in Di Matteo et al.⁴⁷ However, previous studies did
14
15 not consider NPM as a target for pharmacological chaperoning. By using the DSF-based screening,
16
17 the partner site in Bergen identified stabilizing hit-compounds that bind to a hydrophobic pocket
18
19 in NPM and show chaperoning potential. The phenyl ring of the best and novel NPM-hit N-(5-
20
21 bromopyridin-2-yl)-2-methyl-5-oxo-7-phenyl-4-pyridin-2-yl-4,6,7,8-tetrahydro-1H-quinoline-3-
22
23 carboxamide (compound 1), binds into the pocket where it is surrounded by a number of
24
25 hydrophobic residues adjacent to the AML-mutation sites Trp288 and 290. Compound 1
26
27 additionally establishes favorable polar interactions with Lys257, Ser260 (Figure 5D).
28
29 Furthermore, in a cellular system, this hit compound reduced the aggregation and increased nuclear
30
31 localization of NMP mutants. The identified hits appear promising for PCs-based therapies for
32
33 AML⁴⁶.

34
35 DSF constitutes a sensitive assay of ligand binding and an efficient biophysical readout for target-
36
37 based screening, contributing to the EU-OPENSOURCE portfolio. As exemplified above, this
38
39 primary screen is particularly appropriate for the early stage discovery of stabilizing ligands that
40
41 can be developed into chaperoning treatments. Nevertheless, DSF-based screening has also been
42
43 successfully applied for the discovery of effective inhibitors^{48, 49}. Most screening campaigns
44
45 performed so far at the Bergen partner site have used diversity libraries of up to 18,000 compounds.
46
47 However, recent updates in robotics allow the screening of libraries with a throughput of 50,000
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 compounds in about two weeks, paving the way for screening subsets of the EU-OPENSSCREEN
4
5 compound collection for selected targets in high throughput screening times.
6
7

8 **Project E: High quality chemical tools for the perturbation of aberrant Wnt/ β -Catenin**
9 **signaling - construction of tankyrase inhibitors by a hybridization approach**
10

11
12
13 WNT/ β -catenin signalling is altered in a variety of tumors including tumors emerging from
14 colorectal tissue, uterus, pancreas, skin, liver, thyroid, prostate, ovary, stomach, lung, lymphoid,
15
16 bladder, brain, breast and kidney. Increased β -catenin levels have been identified as a central
17
18 factor in T-cell infiltration in melanoma specimens⁵⁰ and a correlation between WNT/ β -catenin
19
20 pathway activation and immune exclusion has been observed across numerous human cancers⁵¹.
21
22
23 The key effector in the hippo pathway, YAP, has also been identified as an oncoprotein whose
24
25 expression is elevated in various human cancers⁵².
26
27
28

29
30 Tankyrase 1 and 2 (TNKS1/PARP-5a/ARTD5 and TNKS2/PARP-5b/ARTD6) are members of
31
32 the poly-ADP-ribose polymerase (PARP) subfamily of the human ADP-ribosyltransferase family
33
34 of enzymes with homology to diphtheria toxin. Tankyrases have been identified as regulators of
35
36 the WNT/ β -catenin signalling pathway via interactions with AXIN protein and a regulator of the
37
38 hippo-signalling pathway via interactions with members of the AMOT family of proteins⁵². The
39
40 inhibition of tankyrases produces elevated AXIN protein levels and reduced levels of cellular
41
42 β -catenin even in the absence of a dysfunctional and truncated form of APC protein. The inhibition
43
44 of tankyrases also stabilizes the AMOT family proteins, thereby suppressing YAP
45
46 oncogenic functions.
47
48
49

50
51
52 The work presented here is a close collaboration between the FMP medicinal chemistry research
53
54 group, which is an EU-OPENSSCREEN medicinal chemistry partner site, the Department of
55
56
57
58
59
60

1
2
3 Immunology and Transfusion Medicine, Oslo University Hospital and the Faculty of Biochemistry
4 and Molecular Medicine, Biocenter Oulu, University of Oulu⁵³. The project started off with an
5
6 initial screening campaign at the FMP Screening Unit — an EU-OPENSSCREEN screening partner
7
8 site — and yielded a 1,2,4-triazole based selective tankyrase inhibitor JW74 with reasonable
9
10 biochemical affinity (TNKS2 IC₅₀ = 0.46 μM) but only moderate cellular activity (HEK293 IC₅₀
11
12 = 1.01 μM)⁵⁴. While other reports on screening approaches in the context of tankyrase inhibitors
13
14 have been published previously as well⁵⁵⁻⁵⁸, a subsequent optimization by a classical analoguing
15
16 strategy led to a first lead structure **6** (G007-LK) (HEK293: IC₅₀ = 0.05μM; TNKS2: IC₅₀ =
17
18 0.025μM⁵⁹), which was highly potent and showed outstanding selectivity towards the other
19
20 members of the ARTD family⁶⁰. Crystal structures of tankyrase had revealed that the early
21
22 inhibitors compete with the substrate NAD⁺ through binding to the conserved nicotinamide⁶¹ or to
23
24 the adenosine subpocket⁶². Analysis of the co-crystal structure of **6** showed that it binds to the
25
26 adenosine subpocket explaining the high selectivity towards tankyrases over other ARTDs.
27
28 However, albeit showing an excellent oral bioavailability in mice further development of **6** was
29
30 hampered by poor pharmacokinetics in rats and concerns on solubility and photostability.
31
32 Structurally, this was attributed to the extended, highly conjugated, aromatic system incorporating
33
34 a vinylic bond in combination with the intrinsically high lipophilicity, low Fsp³ content and a
35
36 critically high molecular weight for some derivatives. Collectively, this suggested that incremental
37
38 structural changes by further classical optimization may not solve these issues. It was therefore
39
40 reasoned that a structure-guided hybridization approach aiming to partially deconstructing the
41
42 structure of **6** to a virtual fragment and merging/joining it with a new privileged tankyrase binding
43
44 motif would provide the necessary significant structural change while still preserving the binding
45
46 mode and distinct interaction pattern for affinity and selectivity⁵³. After analyzing several available
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 tankyrase inhibitor co-crystal structures, it was hypothesized based on the co-crystal structures of
4
5 TNKS2-**6** (PDB: 4HYF) and TNKS1-**7** (PDB: 4K4E)⁶³, that joining the diaryl substituted 1,2,4-
6
7 triazole of **6** and the benzimidazolone of **7** with an appropriate linker would yield a suitable hybrid
8
9 inhibitor (**Figure 6A, B**). Three different linker types, phenyl, cyclohexyl and cyclobutyl, were
10
11 chosen as replacements of the vinylic bond to provide the appropriate distance and conformational
12
13 adaptability within this new class of tankyrase inhibitors. Docking was used to check compatibility
14
15 with the binding pocket and synthesis of the three designed inhibitors was accomplished in a 9-
16
17 step synthesis. Initial testing for TNKS1 and TNKS2 affinity in a biochemical assay and activity
18
19 in human embryonic kidney HEK293 as well as in the human colon SW480 cell lines as functional
20
21 assays of the WNT/ β -catenin signaling pathway, revealed compound **8** as the most potent hybrid
22
23 tankyrase inhibitor for which an activity was observed (HEK293: IC_{50} = 19 nM; SW480: IC_{50} =
24
25 70 nM), accompanied by a favorable biochemical IC_{50} (TNKS1: IC_{50} = 29 nM; TNKS2: IC_{50} = 6.3
26
27 nM). Inspection of the co-crystal structure confirmed that the hybrid inhibitor recapitulated the
28
29 binding mode of both parent inhibitors (PDB: 5NOB, **Figure 6C**). All essential contacts of the
30
31 used fragments were preserved and addressed in the protein ligand complex TNKS-**8**. Selectivity
32
33 profiling showed an exceptional selectivity over the other ARTD enzymes and none of the other
34
35 tested ARTD enzymes were inhibited at 100 μ M or 10 μ M, respectively. This confirmed the
36
37 hypothesis that the selectivity characteristics of the parent inhibitors would also be inherited by
38
39 the newly designed hybrid compound **8**. In addition, compound **8** showed no relevant inhibition in
40
41 an extended kinase selectivity profiling (320 kinases with >50% inhibition at 10 μ M: 4/320; CLK2:
42
43 73%; MELK: 70%; PRKG1: 66% and TSF1: 52%). In line with the favorable calculated
44
45 physicochemical properties and Lipinski rule-of-5 compliance, **8** showed good *in vitro* ADME
46
47 properties, proved to be metabolically stable in human liver fractions across species, in human
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 hepatocytes, and also showed good predicted absorption properties acceptable for *in vivo* studies.
4
5 An overall good bioavailability in mouse (F = 47%), rat (F = 35%), and dog (F = 91%) upon oral
6
7 administration, including a surprisingly low compound excretion in urine and feces in rat,
8
9 underscored the suitability of **8** as a chemical tool for a subsequent pharmacological *in vivo*
10
11 evaluation.
12
13

14
15 Next, **8** was evaluated in mouse xenografts models using the human colorectal cancer cell line
16
17 COLO 320DM cells in male Balb/c nude mice and a *syngeneic* leukemic p388 mouse model.
18
19 Chronic treatment with **8** was well tolerated and resulted in 53%, 63% and 63% statistically
20
21 significant tumor size reductions after 21 days at 15 mg/kg, 30 mg/kg and 60 mg/kg once daily
22
23 oral administration in the COLO 320DM xenografts model. Similarly, in the leukemic p388 mouse
24
25 model statistically significant tumor size reductions of 32% and 57% for 15 mg/kg and 30 mg/kg,
26
27 respectively, were observed after 10 days (**Figure 6D**).
28
29

30
31 These results highlight the suitability of compound **8** as a new high quality chemical tool to
32
33 investigate the role of tankyrases in cellular but also in *in vivo* pharmacological models. This novel
34
35 lead structure is now further developed in an extensive medicinal chemistry lead-to-candidate
36
37 program in collaboration with Mercachem, Nijmegen, Netherlands (Patent application
38
39 WO2018118868A1).
40
41

42
43 This example shows how essential subsequent medicinal chemistry approaches are for optimizing
44
45 initial screening hits to obtain high quality probes with cellular and biochemical IC₅₀ values in the
46
47 nanomolar range. Moreover, it illustrates how non-classical medicinal chemistry approaches can
48
49 capitalize from available structural biology information and how a highly interdisciplinary project
50
51 environment will favorably impact future chemical tool compound development within the EU-
52
53 OPENSREEN network.
54
55

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Summary and outlook

The case studies described above represent only a small subset of the EU-OPENSSCREEN network capabilities. The infrastructure will provide the enabling framework and resources for fostering discoveries and multinational, multidisciplinary collaborations in research and development in the field of chemical biology. For example, a chemical compound discovered or synthesized in Spain will be included in the EU-OPENSSCREEN compound collection, will subsequently meet a biological target from Finland in a screening experiment conducted at a facility in Germany, and as a consequence of these collaborative efforts trigger a new research and development program. Moreover, the integration of diverse scientific disciplines is also demonstrated by the inclusion of the entire natural products drug discovery pipeline in EU-OPENSSCREEN, starting with the isolation of microorganisms from extreme environments, through fermentation, extraction and screening, to isolation and chemical characterization of the bioactive natural products. In a further extension of EU-OPENSSCREEN's capabilities, a fragment collection of 1,000 compounds will be established in cooperation with the structural biology infrastructure, INSTRUCT (<https://www.instruct-eric.eu>), and used to support structure based discovery efforts on isolated targets. EU-OPENSSCREEN will provide users access to the fragment collection for their screening campaigns and also to medicinal chemistry expertise to further optimize the identified fragment hits into leads and tool compounds. Other additions to the library such as, for example, DNA encoded libraries or large molecular weight compounds other than natural products are currently not envisioned, and further library expansions will depend on the funding situation.

Furthermore, EU-OPENSSCREEN is building synergies as it will cooperate with a number of other initiatives in Europe (ESFRI, JPI, ERA-Nets, IMI), which promote innovation in the health and

1
2
3 environment sectors. In this way, chemical biology research in Europe can reach a scale
4
5 competitive to other similar large communities in the USA, Asia and elsewhere. Common quality
6
7 and operating standards, which have been defined in the frame of EU-OPENSSCREEN by all
8
9 participating research groups, will for the first time allow screening results of different European
10
11 platforms to be collected and comparatively analyzed in a joint database. Such data (e.g., screening
12
13 results, assay protocols, chemical information about the substances and their biological activities),
14
15 which are generated in expensive research projects, are currently published only in a limited and
16
17 restricted manner (e.g., only selected positive data supporting a hypothesis). The EU-
18
19 OPENSSCREEN working model will link its database to other life sciences databases and
20
21 informatics resources, and actively promote interoperability of produced data.
22
23
24
25

26
27 In the example projects, the wide range of methodological approaches used illustrates the broad
28
29 expertise available within the EU-OPENSSCREEN partner site network and the ability to meet the
30
31 future demands from users. In the malarial target-based drug discovery project from the Latvian
32
33 Institute for Organic Synthesis the original fragment screen, NMR analyses and medicinal
34
35 chemistry design and synthesis allowed for a structure-informed optimization taking hits with
36
37 potencies in the range of 25 μ M to selective compounds with nanomolar potency and well defined
38
39 SAR for the target protein. In the renal cell carcinoma project from Fundación Medina, a natural
40
41 product workflow was described which identified a novel lipodepsipeptide, not previously
42
43 reported in the literature. The compound MDN-0066 was shown to have an effect on cell cycle
44
45 regulation via the VHL/HIF pathway, inducing apoptosis specifically in pVHL-deficient cells. In
46
47 the ESC differentiation project from Fraunhofer IME, a high content phenotypic screening
48
49 approach was implemented with assays extending up to 5 days with complex multi-step media
50
51 replacement, fixation, staining and imaging protocols. These analyses revealed pathway-specific
52
53
54
55
56
57
58
59
60

1
2
3 inhibitors, which were further elucidated using a cost-effective analogue-by-catalogue method,
4
5 which confirmed the role of ROCK inhibition in the initiation of embryonic stem cell
6
7 differentiation into definitive endoderm. The elaborate DSF-based biophysical assay system
8
9 developed at the University of Bergen yielded pharmacological chaperones with the potential to
10
11 correct inborn errors of metabolism and rare forms of cancer, based on conformational stabilization
12
13 of target proteins in a therapeutic setting; and the non-classical medicinal chemistry approaches
14
15 based on available structural biology information in conjunction with iterative screening at the
16
17 FMP medicinal chemistry and screening partner sites of EU-OPENSSCREEN led to the discovery
18
19 of novel tankyrase inhibitors, which recently entered a lead-to-candidate program.
20
21
22
23

24
25 EU-OPENSSCREEN will ensure to also meet the users' demand for exciting new technologies and
26
27 innovative assay designs in the future by regularly updating existing and/or adding new partner
28
29 institutions and member countries. Central to this will be the establishment of a new category of
30
31 EU-OPENSSCREEN partner site focused on offering access to chemical proteomic technologies
32
33 including affinity based and biophysical readouts such as cellular thermal shift assay platforms.
34
35 The newly planned sites will allow for identification of the underlying targets of compounds
36
37 identified in phenotypic screens, as well as off-target and liability associated activities. These
38
39 new platforms will be piloted in the next three years as part of the recently established Horizon2020
40
41 EU-OPENSSCREEN DRIVE project. Altogether, the result of the work of the EU-OPENSSCREEN
42
43 infrastructure and its associated community of screening, medicinal chemistry and disease biology
44
45 expert teams will be high quality chemical probes and bioactivity data sets for use by the wider
46
47 life science as well as the drug discovery communities^{64, 65}.
48
49
50
51
52

53 Acknowledgements

54
55
56
57
58
59
60

1
2
3 NOR-OPENSREEN, funded by the Research Council of Norway (RCN), and the K.G. Jebsen
4 Centre for Neuropsychiatric disorders, for financial support (to A.M.). Jarl Underhaug is
5 acknowledged for developing the in-house software for DSF screening. The Danish Research
6 Infrastructure for Chemical Biology, DK-OPENSREEN, acknowledges financial support from
7 the Ministry of Higher Education and Science (grant case no. 5072-00019B), the Technical
8 University of Denmark and the other contributing universities. The Latvian Institute of Organic
9 Synthesis acknowledges European Regional Development Fund (Agreement No.
10 1.1.1.1/16/A/290) for financial support. L.L. acknowledges the Academy of Finland (grant no.
11 287063 and 294085) and the Jane and Aatos Erkko foundation for funding. EU-OPENSREEN
12 acknowledges its member and observer states for funding and support. Petr Bartunek
13 acknowledges MEYS (LO1220, LM2015063) for funding.

29 Declaration of conflicts of interest

30
31 M.N., S.K., L.L. have filed a patent application on tankyrase inhibitors (WO2018118868). A.M.
32 is co-inventor of granted patent on pharmacological chaperones for MMA type cbIB
33 (ES2485540B1 21/10-2015).

40 References

- 41
42
43 1. Arrowsmith, C. H.; Audia, J. E.; Austin, C.; et al. The promise and peril of chemical probes. *Nat*
44 *Chem Biol* **2015**, *11*, 536-41.
- 45
46 2. Garbaccio, R. M.; Parmee, E. R. The Impact of Chemical Probes in Drug Discovery: A
47 Pharmaceutical Industry Perspective. *Cell Chem Biol* **2016**, *23*, 10-17.
- 48
49 3. Horvath, D.; Lisurek, M.; Rupp, B.; et al. Design of a general-purpose European compound
50 screening library for EU-OPENSREEN. *ChemMedChem* **2014**, *9*, 2309-26.
- 51
52
53 4. Baell, J. B.; Holloway, G. A. New substructure filters for removal of pan assay interference
54 compounds (PAINS) from screening libraries and for their exclusion in bioassays. *J Med Chem*
55 **2010**, *53*, 2719-40.
- 56
57
58
59
60

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
5. Wells, T. N.; Alonso, P. L.; Gutteridge, W. E. New medicines to improve control and contribute to the eradication of malaria. *Nat Rev Drug Discov* **2009**, *8*, 879-91.
6. Rasina, D.; Otikovs, M.; Leitans, J.; et al. Fragment-Based Discovery of 2-Aminoquinazolin-4(3H)-ones As Novel Class Nonpeptidomimetic Inhibitors of the Plasmepsins I, II, and IV. *J Med Chem* **2016**, *59*, 374-87.
7. Banerjee, R.; Liu, J.; Beatty, W.; et al. Four plasmepsins are active in the Plasmodium falciparum food vacuole, including a protease with an active-site histidine. *Proc Natl Acad Sci U S A* **2002**, *99*, 990-5.
8. Meyers, M. J.; Goldberg, D. E. Recent advances in plasmepsin medicinal chemistry and implications for future antimalarial drug discovery efforts. *Curr Top Med Chem* **2012**, *12*, 445-55.
9. Ersmark, K.; Feierberg, I.; Bjelic, S.; et al. Potent inhibitors of the Plasmodium falciparum enzymes plasmepsin I and II devoid of cathepsin D inhibitory activity. *J Med Chem* **2004**, *47*, 110-22.
10. Huizing, A. P.; Mondal, M.; Hirsch, A. K. Fighting malaria: structure-guided discovery of nonpeptidomimetic plasmepsin inhibitors. *J Med Chem* **2015**, *58*, 5151-63.
11. Boss, C.; Corminboeuf, O.; Grisostomi, C.; et al. Achiral, cheap, and potent inhibitors of Plasmepsins I, II, and IV. *ChemMedChem* **2006**, *1*, 1341-5.
12. Hof, F.; Schutz, A.; Fah, C.; et al. Starving the malaria parasite: inhibitors active against the aspartic proteases plasmepsins I, II, and IV. *Angew Chem Int Ed Engl* **2006**, *45*, 2138-41.
13. Harner, M. J.; Frank, A. O.; Fesik, S. W. Fragment-based drug discovery using NMR spectroscopy. *J Biomol Nmr* **2013**, *56*, 65-75.
14. Puchades-Carrasco, L.; Pineda-Lucena, A. Fragment-based Drug Design Using NMR Methods. *Emagres* **2015**, *4*, 241-253.
15. Ersmark, K.; Samuelsson, B.; Hallberg, A. Plasmepsins as potential targets for new antimalarial therapy. *Med Res Rev* **2006**, *26*, 626-66.
16. Schrödinger. Small-Molecule Drug Discovery Suite 2014-1: Glide, version 6.2; Schrödinger, LLC: New York, 2014. 2014.
17. Zurcher, M.; Diederich, F. Structure-based drug design: exploring the proper filling of apolar pockets at enzyme active sites. *J Org Chem* **2008**, *73*, 4345-61.
18. Gnarra, J. R.; Lerman, M. I.; Zbar, B.; et al. Genetics of renal-cell carcinoma and evidence for a critical role for von Hippel-Lindau in renal tumorigenesis. *Semin Oncol* **1995**, *22*, 3-8.
19. Gossage, L.; Eisen, T. Alterations in VHL as potential biomarkers in renal-cell carcinoma. *Nat Rev Clin Oncol* **2010**, *7*, 277-88.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
20. Iliopoulos, O.; Kibel, A.; Gray, S.; et al. Tumour suppression by the human von Hippel-Lindau gene product. *Nat Med* **1995**, *1*, 822-6.
21. Sutphin, P. D.; Chan, D. A.; Li, J. M.; et al. Targeting the loss of the von Hippel-Lindau tumor suppressor gene in renal cell carcinoma cells. *Cancer Res* **2007**, *67*, 5896-905.
22. Turcotte, S.; Chan, D. A.; Sutphin, P. D.; et al. A molecule targeting VHL-deficient renal cell carcinoma that induces autophagy. *Cancer Cell* **2008**, *14*, 90-102.
23. Turcotte, S.; Sutphin, P. D.; Giaccia, A. J. Targeted therapy for the loss of von Hippel-Lindau in renal cell carcinoma: a novel molecule that induces autophagic cell death. *Autophagy* **2008**, *4*, 944-6.
24. Cautain, B.; Pedro, N. d.; Escalona, M. M. d.; et al. HCS strategy targeting dysregulation of the VHL/HIF pathway for drug discovery. *Advances in Bioscience and Biotechnology* **2013**, *04*, 398-405.
25. Cautain, B.; de Pedro, N.; Schulz, C.; et al. Identification of the Lipodepsipeptide MDN-0066, a Novel Inhibitor of VHL/HIF Pathway Produced by a New Pseudomonas Species. *PLoS One* **2015**, *10*, e0125221.
26. Newman, D. J.; Cragg, G. M. Natural products as sources of new drugs over the 30 years from 1981 to 2010. *J Nat Prod* **2012**, *75*, 311-35.
27. Pascual, J.; Garcia-Lopez, M.; Bills, G. F.; et al. Pseudomonas granadensis sp. nov., a new bacterial species isolated from the Tejada, Almirajara and Alhama Natural Park, Granada, Spain. *Int J Syst Evol Microbiol* **2015**, *65*, 625-32.
28. Goda, N.; Ryan, H. E.; Khadivi, B.; et al. Hypoxia-inducible factor 1 alpha is essential for cell cycle arrest during hypoxia. *Mol Cell Biol* **2003**, *23*, 359-369.
29. Korostylev, A.; Mahaddalkar, P. U.; Keminer, O.; et al. A high-content small molecule screen identifies novel inducers of definitive endoderm. *Mol Metab* **2017**, *6*, 640-650.
30. Schiesser, J. V.; Wells, J. M. Generation of beta cells from human pluripotent stem cells: Are we there yet? *Ann Ny Acad Sci* **2014**, *1311*, 124-137.
31. Pagliuca, F. W.; Millman, J. R.; Gurtler, M.; et al. Generation of functional human pancreatic beta cells in vitro. *Cell* **2014**, *159*, 428-39.
32. Rezanian, A.; Bruin, J. E.; Arora, P.; et al. Reversal of diabetes with insulin-producing cells derived in vitro from human pluripotent stem cells. *Nat Biotechnol* **2014**, *32*, 1121-33.
33. Burtscher, I.; Lickert, H. Foxa2 regulates polarity and epithelialization in the endoderm germ layer of the mouse embryo. *Development* **2009**, *136*, 1029-38.
34. Lariosa-Willingham, K. D.; Rosler, E. S.; Tung, J. S.; et al. A high throughput drug screening assay to identify compounds that promote oligodendrocyte differentiation using acutely dissociated and purified oligodendrocyte precursor cells. *BMC Res Notes* **2016**, *9*, 419.

- 1
2
3 35. Mattiazzi Usaj, M.; Styles, E. B.; Verster, A. J.; et al. High-Content Screening for Quantitative Cell
4 Biology. *Trends Cell Biol* **2016**, *26*, 598-611.
5
6 36. Ryan, S. L.; Baird, A. M.; Vaz, G.; et al. Drug Discovery Approaches Utilizing Three-Dimensional
7 Cell Culture. *Assay Drug Dev Technol* **2016**, *14*, 19-28.
8
9 37. Requejo-Aguilar, R.; Alastrue-Agudo, A.; Cases-Villar, M.; et al. Combined polymer-curcumin
10 conjugate and ependymal progenitor/stem cell treatment enhances spinal cord injury functional
11 recovery. *Biomaterials* **2017**, *113*, 18-30.
12
13 38. Sharer, J. D. An Overview of Biochemical Genetics. *Curr Protoc Hum Genet* **2016**, *89*, 17 1 1-17 1
14 16.
15
16 39. Pey, A. L.; Stricher, F.; Serrano, L.; et al. Predicted effects of missense mutations on native-state
17 stability account for phenotypic outcome in phenylketonuria, a paradigm of misfolding diseases.
18 *Am. J. Hum. Genet.* **2007**, *81*, 1006-24.
19
20 40. Pey, A. L.; Ying, M.; Cremades, N.; et al. Identification of pharmacological chaperones as
21 potential therapeutic agents to treat phenylketonuria. *J. Clin. Invest.* **2008**, *118*, 2858-67.
22
23 41. Underhaug, J.; Aubi, O.; Martinez, A. Phenylalanine hydroxylase misfolding and pharmacological
24 chaperones. *Current topics in medicinal chemistry* **2012**, *12*, 2534-45.
25
26 42. Yuste-Checa, P.; Brasil, S.; Gamez, A.; et al. Pharmacological Chaperoning: A Potential Treatment
27 for PMM2-CDG. *Hum Mutat* **2017**, *38*, 160-168.
28
29 43. Jorge-Finnigan, A.; Brasil, S.; Underhaug, J.; et al. Pharmacological chaperones as a potential
30 therapeutic option in methylmalonic aciduria cblB type. *Human molecular genetics* **2013**, *22*,
31 3680-9.
32
33 44. Brasil, S.; Briso-Montiano, A.; Gamez, A.; et al. New perspectives for pharmacological
34 chaperoning treatment in methylmalonic aciduria cblB type. *Biochim Biophys Acta* **2018**, *1864*,
35 640-648.
36
37 45. Hole, M.; Underhaug, J.; Diez, H.; et al. Discovery of compounds that protect tyrosine
38 hydroxylase activity through different mechanisms. *Biochim Biophys Acta* **2015**, *1854*, 1078-89.
39
40 46. Urbaneja, M. A.; Skjaerven, L.; Aubi, O.; et al. Conformational stabilization as a strategy to
41 prevent nucleophosmin mislocalization in leukemia. *Sci Rep* **2017**, *7*, 13959.
42
43 47. Di Matteo, A.; Franceschini, M.; Chiarella, S.; et al. Molecules that target nucleophosmin for
44 cancer treatment: an update. *Oncotarget* **2016**, *7*, 44821-44840.
45
46 48. Aubi, O.; Flydal, M. I.; Zheng, H.; et al. Discovery of a Specific Inhibitor of Pyomelanin Synthesis
47 in *Legionella pneumophila*. *J Med Chem* **2015**, *58*, 8402-12.
48
49 49. Martin, I.; Underhaug, J.; Celaya, G.; et al. Screening and Evaluation of Small Organic Molecules
50 as ClpB Inhibitors and Potential Antimicrobials. *Journal of medicinal chemistry* **2013**, *56*, 7177-
51 89.
52
53
54
55
56
57
58
59
60

- 1
2
3 50. Spranger, S.; Bao, R.; Gajewski, T. F. Melanoma-intrinsic beta-catenin signalling prevents anti-
4 tumour immunity. *Nature* **2015**, *523*, 231-5.
5
6 51. Luke, J. J.; Bao, R. Y.; Spranger, S.; et al. Correlation of WNT/beta-catenin pathway activation
7 with immune exclusion across most human cancers. *J Clin Oncol* **2016**, *34*.
8
9 52. Wang, W.; Li, N.; Li, X.; et al. Tankyrase Inhibitors Target YAP by Stabilizing Angiomin Family
10 Proteins. *Cell Rep* **2015**, *13*, 524-532.
11
12 53. Anumala, U. R.; Waaler, J.; Nkizinkiko, Y.; et al. Discovery of a Novel Series of Tankyrase
13 Inhibitors by a Hybridization Approach. *J Med Chem* **2017**, *60*, 10013-10025.
14
15 54. Waaler, J.; Machon, O.; Tumova, L.; et al. A novel tankyrase inhibitor decreases canonical Wnt
16 signaling in colon carcinoma cells and reduces tumor growth in conditional APC mutant mice.
17 *Cancer Res* **2012**, *72*, 2822-32.
18
19 55. Shultz, M. D.; Kirby, C. A.; Stams, T.; et al. [1,2,4]triazol-3-ylsulfanylmethyl-3-phenyl-
20 [1,2,4]oxadiazoles: antagonists of the Wnt pathway that inhibit tankyrases 1 and 2 via novel
21 adenosine pocket binding. *J Med Chem* **2012**, *55*, 1127-36.
22
23 56. Narwal, M.; Haikarainen, T.; Fallarero, A.; et al. Screening and structural analysis of flavones
24 inhibiting tankyrases. *J Med Chem* **2013**, *56*, 3507-17.
25
26 57. Narwal, M.; Fallarero, A.; Vuorela, P.; et al. Homogeneous screening assay for human tankyrase.
27 *J Biomol Screen* **2012**, *17*, 593-604.
28
29 58. Samain, F.; Ekblad, T.; Mikutis, G.; et al. Tankyrase 1 Inhibitors with Drug-like Properties
30 Identified by Screening a DNA-Encoded Chemical Library. *J Med Chem* **2015**, *58*, 5143-9.
31
32 59. Voronkov, A.; Holsworth, D. D.; Waaler, J.; et al. Structural basis and SAR for G007-LK, a lead
33 stage 1,2,4-triazole based specific tankyrase 1/2 inhibitor. *J Med Chem* **2013**, *56*, 3012-23.
34
35 60. Lau, T.; Chan, E.; Callow, M.; et al. A novel tankyrase small-molecule inhibitor suppresses APC
36 mutation-driven colorectal tumor growth. *Cancer Res* **2013**, *73*, 3132-44.
37
38 61. Karlberg, T.; Markova, N.; Johansson, I.; et al. Structural basis for the interaction between
39 tankyrase-2 and a potent Wnt-signaling inhibitor. *J Med Chem* **2010**, *53*, 5352-5.
40
41 62. Narwal, M.; Venkannagari, H.; Lehtio, L. Structural basis of selective inhibition of human
42 tankyrases. *J Med Chem* **2012**, *55*, 1360-7.
43
44 63. Bregman, H.; Chakka, N.; Guzman-Perez, A.; et al. Discovery of novel, induced-pocket binding
45 oxazolidinones as potent, selective, and orally bioavailable tankyrase inhibitors. *J Med Chem*
46 **2013**, *56*, 4320-42.
47
48 64. Antolin, A. A.; Jalencas, X.; Yelamos, J.; et al. Identification of Pim Kinases as Novel Targets for
49 PJ34 with Confounding Effects in PARP Biology. *Acs Chem Biol* **2012**, *7*, 1962-1967.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 65. Antolin, A. A.; Mestres, J. Distant Polypharmacology among MLP Chemical Probes. *Acs Chem Biol*
4 **2015**, *10*, 395-400.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Figure Legends

Figure 1 | Working model of the EU-OPENSSCREEN ERIC. Three categories of users are envisioned (assay provider, compound provider, database user). The compound collection comprising academic and commercial compounds is stored centrally at the headquarters. Bioprofiling of academic and commercial compounds occurs at the bioprofiling site. Compound screening and medicinal chemistry is carried out at the individual EU-OPENSSCREEN partner sites. Data is stored at the database site and can be accessed via the EU-OPENSSCREEN database.

Figure 2 | 2-Aminoquinazolin-4(3H)-ones as potential new non-peptidomimetic lead structures for antimalarial agents. A) Experimental pathway that led to the discovery of the 2-aminoquinazolin-4(3H)-one scaffold. B) Optimized inhibitor hit *rac-3a* from NMR studies of fragment hits **1** and **2**, and inhibitor *rac-3b* from molecular modeling studies. C) X-ray structure of *rac-3b* bound to Plm II (PDB ID 4Z22). D) Chemical structures of the two Plm inhibitors *rac-4a* and *rac-4b*. Figure adapted with permission from Rasina et al. (J. Med. Chem. 59, 1, 374-387)⁶. Copyright (2016) American Chemical Society.


Figure 3 | MDN-0066 as a potential new lead structure inducing apoptosis specifically in cells deficient for the tumor suppressor pVHL. A) Structure of the newly discovered lipodepsipeptide MDN-0066. B) Concentration response curve of MDN-0066 showing increased sensitivity to MDN-0066 in pVHL deficient cells compared to the pVHL rescue cell line. C) Annexin V/propidium iodide (PI) staining confirming induction of apoptosis specifically in RCC4-VA cells in response to MDN-0066 exposure. Lower left quadrant = live cells; lower right quadrant = early apoptotic cells; upper right quadrant = apoptotic cells; upper left quadrant = necrotic cells. Figure adapted from Cautain et al. (PLoS One, 2015, 10, e0125221)²⁵.

1
2
3 **Figure 4** | ROCK inhibitors as inducers of definitive endoderm formation. A) General analysis
4 workflow for primary hit selection. B) DMSO solvent control. C) Pluripotency control. D)
5 Differentiation control (blue: DAPI = nuclei stain, red: Alexa-Fluor 555 = Oct-3/4 detection, green:
6 Alexa-Fluor 488 = FoxA2 detection). Figure adapted from Korostylev et al. (2017)²⁹.
7
8
9

10
11
12 **Figure 5** | DSF-based screening and discovery of pharmacological chaperones. A) In DSF-
13 screening the protein is mixed with SYPRO Orange and the compounds transferred from the
14 chemical libraries. Fluorescence is recorded at increasing temperatures and stabilizing compounds
15 (i.e., significant positive ΔT_m) with respect to controls are selected. B) Schematic funnel for the
16 discovery of pharmacological chaperones, including the primary DSF screening, secondary assays
17 by SPR and enzymatic activity and validation by efficacy tests in cells and *in vivo*. C) The binding
18 site of hit compound V in one subunit of tetrameric ATR (PDB ID 2IDX), predicted by molecular
19 docking, showing interacting residues and the location of the substrate ATP. See also Jorge-
20 Finnigan et al. (2013)⁴³. D) The binding mode of hit compound 1 to the C-terminal domain of
21 NPM (PDB ID 2LLH), predicted by docking, wherein its phenyl ring is placed in a hydrophobic
22 pocket establishing polar interactions with solvent accessible residues. See also Urbaneja et al.
23 (2017)⁴⁶. The proteins in C) and D) are shown as solvent-accessible surfaces with projected
24 interpolated positive (blue) and negative (red) charges.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42


43 **Figure 6** | Structure guided chemical hybridization approach towards a new specific tankyrase
44 inhibitor as a high quality chemical tool. A) Structures of **6** – **8**. Hybridization logic to yield **8** from
45 the deconstructed parent inhibitors **6** and **7**. B) Superposition of **6** (blue) and **7** (green) co-crystal
46 structures (PDB: 4HYF and 4K4E). Only TNKS2 protein is shown for clarity. C) Co-crystal
47 structure of **8** (magenta) with TNKS2. (PDB: 5NOB). D) Anti-tumor activity of **8** in xenograft
48 models. Left panel: COLO320DM colon cancer xenograft; right panel: isogenic p388 leukemia
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 mouse model. Reduction of tumor volume (mm³) versus vehicle treated controls (blue) after once
4 daily oral dosing of **8** at various depicted doses. Statistical significance is indicated: ANOVA on
5 Ranks/Dunn's method, $p < 0.05$ (*), One Way ANOVA/Holm-Sidak method, $p < 0.001$ (**) and
6 One-tailed p -value < 0.05 (***). Figure compiled and modified with permission based on the
7 original publication Anumala et al. (J. Med. Chem. 60, 24, 10013-10025)⁵³. Copyright (2017)
8 American Chemical Society.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Working model of the EU-OPENSREEN ERIC

302x152mm (150 x 150 DPI)




2-Aminoquinazolin-4(3H)-ones as potential new non-peptidomimetic lead structures for antimalarial agents

292x187mm (150 x 150 DPI)




MDN-0066 as a potential new lead structure inducing apoptosis specifically in cells deficient for the tumor suppressor pVHL.

318x189mm (150 x 150 DPI)




ROCK inhibitors as inducers of definitive endoderm formation

155x141mm (150 x 150 DPI)



DSF-based screening and discovery of pharmacological chaperones

200x152mm (300 x 300 DPI)



25 Structure guided chemical hybridization approach towards a new specific tankyrase inhibitor as a high
26 quality chemical tool

27 338x190mm (112 x 112 DPI)

Supplemental Material for EU-OPENSREEN: A novel collaborative approach to facilitate chemical biology.

Corresponding Authors:


Philip Gribbon, Fraunhofer Institute for Molecular Biology and Applied Ecology IME, Screening Port, Schnackenburgallee 114, 22525 Hamburg, Germany, Email:

Philip.Gribbon@ime.fraunhofer.de

Wolfgang Fecke, EU-OPENSREEN, Central Office, Robert-Roessle-Strasse 10, 13125 Berlin, Germany, Email: wolfgang.fecke@eu-openscreen.eu

Cell Cycle arrest (48 h)

	RCC4VA -	RCC4VA +	RCC4VHL-	RCC4VHL+
SubG0	2.5 ± 0.8	2.9 ± 0.6	3.3 ± 0.3	3 ± 1.1
G0/G1	75.5 ± 4.2	37.5 ± 4.6	60 ± 1.4	53 ± 3.1
S	6 ± 2.1	18.5 ± 5.2	11 ± 0.8	7 ± 2.3
G2/M	10 ± 0.5	26.2 ± 3.1	17 ± 3	24 ± 1.2



Supplementary Figure 1 | Cell cycle arrest of RCC4-VA/VHL cells upon MDN-0066 treatment.

Flow cytometry analysis using a PI Flow Kit upon MDN-0066 treatment was measured. Percent of cell cycle phase is shown in a table form with the SubG0, G0/G1, S and G2/M phases for the two cell lines in the absence or following a 48 h incubation with MDN-0066 at 24 μ M. Represent results of one of 3 independent experiments. Figure adapted from Cautain et al. (PLoS One, 2015, 10, e0125221).

References

Cautain, B.; de Pedro, N.; Schulz, C.; et al. Identification of the Lipodepsipeptide MDN-0066, a Novel Inhibitor of VHL/HIF Pathway Produced by a New Pseudomonas Species. *PLoS One* **2015**, *10*, e0125221.