



Predicting and mapping human risk of exposure to *Ixodes ricinus* nymphs in northern Europe using climatic and environmental data

Kjær, Lene Jung; Soleng, Arnulf ; Skarsfjord Edgar, Kristin ; Lindstedt, Heidi Elisabeth H.; Mørk Paulsen, Katrine; Andreassen, Åshild Kristine; Korslund, Lars; Kjelland, Vivian; Slettan, Audun; Stuen, Snorre

Total number of authors:

16

Publication date:

2018

Document Version

Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Kjær, L. J., Soleng, A., Skarsfjord Edgar, K., Lindstedt, H. E. H., Mørk Paulsen, K., Andreassen, Å. K., Korslund, L., Kjelland, V., Slettan, A., Stuen, S., Kjellander, P., Christensson, M., Teräväinen, M., Baum, A., Schou, K. K., & Bødker, R. (2018). *Predicting and mapping human risk of exposure to Ixodes ricinus nymphs in northern Europe using climatic and environmental data*. Abstract from Workshop on Arthropod-Borne Diseases, Isle of Riems, Mecklenburg-Vorpommern, Germany.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

In recent years, focus on tick-borne diseases has increased as diseases such as Lyme disease and tick-borne encephalitis have become more common and represent a health problem in many parts of Scandinavia. More effective prevention of infections requires a better understanding of the factors affecting the vector abundance as well as human exposure to the vectors. Hence, there is a great need for analyses and models that can predict how vectors and their associated diseases are distributed now and possibly in the future.

As a part of the ScandTick Innovation project, we surveyed tick nymphs at 159 sites (forests and meadows) in Denmark, southern Norway and south-eastern Sweden. At each site we measured presence/absence, and used the data obtained along with environmental data from satellite images to run Boosted Regression Tree machine learning algorithms to predict overall distribution in southern Scandinavia. Together with the predicted distribution maps, we used human density maps to identify and plot areas with high risk of exposure to ticks.

The predicted distribution and the spatial variation found corresponded well with known distributions of ticks in Scandinavia (sensitivity: 91%, specificity: 60%), and we found that the model was predominantly temperature-driven. Because presence was strongly correlated with forested habitats the risk areas were much larger in Sweden and Norway compared to Denmark. When combining these distribution maps with human population density maps, we were able to quantify the proportion of people living in areas with tick presence in Scandinavia. We found that although tick nymphs were restricted to a small proportion of the modelled area, high proportions of the human populations (67-79%) lived within these same areas. The model suggests that a potential future range expansion of *I. ricinus* in Scandinavia is likely but may only affect a relatively small additional proportion of the human population.