



Antioxidant composition and activity of seaweed *Saccharina latissima*: a seasonal perspective

S. Marinho, Gonçalo ; Sørensen, Ann-Dorit Moltke; Safafar, Hamed; Pedersen, Anja H.; Holdt, Susan L.

Publication date:
2017

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

S. Marinho, G., Sørensen, A-D. M., Safafar, H., Pedersen, A. H., & Holdt, S. L. (2017). *Antioxidant composition and activity of seaweed Saccharina latissima: a seasonal perspective*. Abstract from 6th congress of international society for applied phycology, Nantes, France.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Antioxidant composition and activity of seaweed

Saccharina latissima: a seasonal perspective

Gonçalo S. Marinho, Ann-Dorit Moltke Sørensen, Hamed Safafar, Anja H. Pedersen,
Susan L. Holdt

DTU Food, The National Food Institute, Technical University of Denmark, DK-2800 Kgs.
Lyngby, Denmark
gosm@food.dtu.dk

Keywords: Sugar kelp, TPC, Polyphenols, Flavonoids, HPLC, DPPH

Safety concerns regarding reported toxicity of artificial antioxidants lead the search for novel natural antioxidants. In this context, seaweeds have been receiving increasing attention as a promising source of antioxidants such as phenolic compounds (e.g. phenolic acids and flavonoids), carotenoids (e.g. fucoxanthin and β -carotene), and phycobiliproteins. Nevertheless, seaweed composition generally presents marked seasonal variations. The present study aimed at evaluating seasonal variations in the antioxidant composition and activity of sugar kelp, *Saccharina latissima*, cultivated at two different sites; in close proximity to a blue mussel and rainbow trout farm (IMTA), and at a reference/control site (REF), outside Horsens fjord, Denmark. Rope cultivated seaweed was sampled bi-monthly from three droppers ($n=3$; ≥ 10 specimens per replicate) from May 2013 to May 2014; the biomass was freeze-dried and stored frozen until further analysis. Total phenolic compounds (TPC) were determined both spectrophotometrically (Folin-Ciocalteu) and by HPLC coupled to UV detection (255 nm). Concentration of TPC determined spectrophotometrically was higher in November and January (1,225-2,406 $\mu\text{g GAE}\cdot\text{g}^{-1}$ DW) compared to September. Noteworthy, concentration of TPC determined by HPLC was around 10 times lower compared to the previous method. Gallic acid, rutin and quercitrin were identified out of 18 polyphenol standards. Moreover, TPC concentration in November (203 $\mu\text{g}\cdot\text{g}^{-1}$ DW) was similar to that found in September and January, but significantly higher compared to May 2014. The highest concentration of flavonoids was found in September (4,558-4,830 $\mu\text{g RE}\cdot\text{g}^{-1}$ DW; $p<0.05$). The pigment profile (i.e. type of pigments) did not change seasonally, whereas fucoxanthin and chlorophyll a were the most abundant. Fucoxanthin concentration was lower in September compared with all the other months (261-665 $\mu\text{g}\cdot\text{g}^{-1}$ DW; $p<0.05$). Total antioxidant capacity (TAC) was significantly higher in November (3,835-4,050 $\mu\text{g GAE}\cdot\text{g}^{-1}$ DW). Regarding the antioxidant activity (DPPH), there was no significant difference in the IC₅₀ between seasons, although it was tendentially lower in September and November (0.42-0.49 $\text{mg}\cdot\text{mL}^{-1}$). Statistical analysis reveals a strong positive correlation between TAC and TPC, whereas flavonoids and fucoxanthin were poorly correlated to TAC. There was no significant difference for all the analyzed variables between cultivation sites. This study reveals marked seasonal variations, and high biological variability (difference among replicates), in the content of antioxidants which needs to be taken into account when considering seaweed as a source of antioxidants. Moreover, our results suggest that spectrophotometric methods may result in overestimation of TPC and that HPLC is more selective, also identifying some phenolic acids and flavonoids.