



Optimization of individual fish intake in Denmark

Persson, Inez Maria; Fagt, Sisse; Poulsen, Morten; Vieux, Florent; Nauta, Maarten

Publication date:
2017

Document Version
Version created as part of publication process; publisher's layout; not normally made publicly available

[Link back to DTU Orbit](#)

Citation (APA):
Persson, I. M., Fagt, S., Poulsen, M., Vieux, F., & Nauta, M. (2017). *Optimization of individual fish intake in Denmark*. Abstract from GoFood 2017, Lund, Sweden.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Only 15% of the Danes meet the fish intakes recommended by the official Danish dietary guidelines. This study therefore proposes a method to find fish intakes that fulfill several dietary criteria, while simultaneously deviating as little as possible from the current intakes. For an individual consumer this is more realistic and achievable than a general recommendation.

A mathematical optimization model was developed, using quadratic programming. Individual recommended fish intakes were identified, meeting recommendations on fatty acids and vitamin D, without violating tolerable intake levels on methyl mercury and dioxins. Individual intakes for eight fish species were proposed for 3016 Danes, whose intakes were known from a Dietary survey. For women, the average change in fish intake was + 14.2 g of lean fish and + 62.6 g of fatty fish per week and for men the respective numbers were + 12.2 and + 55.1.

This study shows how mathematical optimization models can be used to advise individual consumers on the optimization of their diet, based on their current intake. The model can be modified to address other foods, food component and contaminants and the method can also address sustainability (e.g. by using constraints on greenhouse gas emission) or costs.