



## Sentinel-1 SAR for wind energy roughness maps over land

Dellwik, Ebba; Karagali, Ioanna; Svensson, Elin; Skriver, Henning; Badger, Merete

*Published in:*  
Geophysical Research Abstracts

*Publication date:*  
2018

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Dellwik, E., Karagali, I., Svensson, E., Skriver, H., & Badger, M. (2018). Sentinel-1 SAR for wind energy roughness maps over land. *Geophysical Research Abstracts*, 20.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## **Sentinel-1 SAR for wind energy roughness maps over land**

Ebba Dellwik (1), Ioanna Karagali (1), Elin Svensson (1), Henning Skriver (2), and Merete Badger (1)

(1) Technical University of Denmark, Wind Energy Department, Roskilde, Denmark (ebde@dtu.dk), (2) Technical University of Denmark, DTU Space, Lyngby Denmark

For the wind energy application, updated information on aerodynamic surface roughness is important for an accurate prediction of the land surface effect on the atmosphere. Tall wind turbines are commonly sited in forested areas, and since the forest both increase turbulence levels and decrease the wind resource, the parametrization of forest roughness in wind models is of high relevance. Here, we investigate whether the Sentinel-1 SAR images can be used to identify high-roughness forested areas. The backscatter properties of the SAR images are compared to digital surface models and vegetation density maps derived from near-concurrent aerial lidar scans (ALS). These ALS products have previously shown good results in wind models for the wind energy application, but the scans are costly to perform and therefore typically only represent a snapshot in time, whereas the Sentinel mission SAR images provide frequent updated information. We investigate how the SAR images vary with season over both deciduous and needle-leaf forests and in addition test whether nearby meteorological observations can explain image to image differences in the backscatter level. In order to understand the backscatter level, new products from the ALS point cloud are derived and compared with the SAR images. This part of the work is focused on whether we also can quantify the roughness based on the SAR backscatter. Since SAR images are affected by speckle noise, they are averaged over monthly and bimonthly intervals after careful inspection of each image. The work is focused on the Østerild test site for large wind turbines in Northern Denmark, where extensive wind experiments have been performed.