



Toxoplasma gondii seroprevalence in feral and farmed Danish mink

Petersen, Heidi Huus; Sengupta, Mita Eva; Pagh, Stine; Chriél, Mariann

Publication date:
2018

Document Version
Version created as part of publication process; publisher's layout; not normally made publicly available

[Link back to DTU Orbit](#)

Citation (APA):
Petersen, H. H., Sengupta, M. E., Pagh, S., & Chriél, M. (2018). *Toxoplasma gondii seroprevalence in feral and farmed Danish mink*. Abstract from 13th Workshop of National Reference Laboratories for Parasites, Rome, Italy.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Thirteenth Workshop of National Reference Laboratories for Parasites

Time: 23 – 24 May 2019

Venue: Istituto Superiore de Sanità, Rome, Italy

Toxoplasma gondii seroprevalence in feral and farmed Danish mink

Petersen, H.H.^{1*}, Sengupta, M.E.², Pagh, S.³, Chriel, M.⁴

¹National Veterinary Institute, DTU, Lyngby, Denmark.

²Center for Macroecology, Evolution and Climate, University of Copenhagen, Copenhagen, Denmark

³ Department of Chemistry and Bioscience, University of Aalborg, Aalborg

*presenting author email: hhet@vet.dtu.dk

Abstract

In Denmark, American mink (*Neovison vison*) are bred for their fur, and Denmark is currently one of the largest producers of mink skins. Moreover, feral mink populations are common in Denmark, but they are regarded as an invasive species originating from farm escapes. The zoonotic parasite *Toxoplasma gondii* has a wide host range, including mink. A Danish study from 1994 observed 3% sero-positive farmed mink by latex agglutination test. However, the literature on *T. gondii* in mink is generally scarce, and to our knowledge, no studies on *T. gondii* in feral, Danish mink have been carried out. *Toxoplasma gondii* infections in mink might be of public health importance, when dead mink are handled. This study examined the antibodies against *T. gondii* in feral- and farmed mink. In total, 112 farmed mink submitted for diagnostic examination, and 228 feral mink sampled by Danish hunters were included in the study. At necropsy, the heart was removed and stored at -20°C until analysis. Meat juice was extracted from the hearts when thawed and analyzed for anti-*T. gondii* antibodies using a commercial indirect ELISA. None of the farmed mink were sero-positive, while 50.5% of the feral mink were sero-positive. Of the positive feral mink, 13.4% (15/112) had fur colors other than brown, indicating a recent escape from farms. Significantly ($p=0.0002$) more feral male (63.5%) than female (36.2%) were sero-positive. In consideration of the results, precautions should be taken when handling feral mink. In contrast, handling farmed mink, pose a neglectable risk of acquiring *T. gondii* infections.