



Effect of removing phenolic compounds on interfacial behavior of protein isolated from de-oiled sunflower cake

Gültekin Subaşı, Büşra; Casanova, Federico; Çapanoğlu Güven, Esra; Mohammadifar, Mohammad Amin

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Gültekin Subaşı, B., Casanova, F., Çapanoğlu Güven, E., & Mohammadifar, M. A. (2019). *Effect of removing phenolic compounds on interfacial behavior of protein isolated from de-oiled sunflower cake*. Poster session presented at 8th International Symposium on Food Rheology and Structure, Zürich, Switzerland.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Effect of removing phenolic compounds on interfacial behavior of protein isolated from de-oiled sunflower cake

Büşra Gültekin Subaşı^{abc}, Federico Casanova^{b*}, Esra Çapanoğlu Güven^c,
Mohammad Amin Mohammadifar^{b*}

^a Cumhuriyet University, Hafik Kamer Ornek MYO, Sivas, Turkey, ^b Food Production Engineering, DTU Food, Technical University of Denmark, Søtofts Plads 227, Dk-2800 Lyngby, Denmark, ^c Food Engineering Department, Chemical and Metallurgical Engineering Faculty, İstanbul Technical University, İstanbul, Turkey.




Table 1. Chemical composition of natural (N) and de-phenolized (DP) protein isolates

Sample	Moisture(%)	Ash	Protein (%)	Carbohydrate (%)	Total phenolics (†)
N	3.16 ± 0.11	1.52 ± 0.45	88.18 ± 1.21	1.82 ± 0.21	5.05 ± 0.60
DP	5.12 ± 0.18	0.64 ± 0.14	95.46 ± 0.42	0.42 ± 0.07	0.09 ± 0.04

Total phenolic content is defined as mg gallic acid acid equivalent (GAE) / 100g dry weight

Table 2. Amino acid composition of natural (N) and de-phenolized (DP) protein isolates

Amino acid	Natural SPI (g / 100 g)	De-phenolized SPI (g / 100 g)
Arg	12.43	12.56
Ser	4.65	4.81
Hyp	0	0.05
Gly	4.48	4.52
Thr	3.78	4.06
Ala	3.46	3.68
Pro	3.93	4.38
Met	1.05	1.79
Asp	10.27	11.65
Val	10.87	12.03
His	2.53	2.86
Lys	2.92	3.14
Glu	31.54	30.25
Leu	6.16	6.81
Phe	5.44	5.83
Ile	6.23	7.53
C-C	0.80	0.93
Tyr	2.23	2.65




Figure 1. Storage modulus at the air / water interface of sunflower protein isolates





Figure 2. Foaming capacity and stability of sunflower protein isolates

Results

Table 1. Indicates that removing phenol content results in significant change in ash, protein, carbohydrate content and amino acid. Low influence in amino acid profile was observed (Table 2). Stronger viscoelastic structure, interfacial layer was observed for dephenolized samples (Figure 1) which was associated with less foam capacity (Figure 2)

Conclusion

Presence of phenolic compounds negatively affects foaming and interfacial properties.

Future directions

For more clear statements and better understanding about the foam ability of DP sunflower protein, other analytical approaches are planned: dynamic interfacial tension, ellipsometry as well as film pressure balance.

* Contact:

Dr. Federico Casanova, feca@food.dtu.dk, Building 227, room 162

Dr. Mohammad Amin Mohammadifar moamo@food.dtu.dk, Building 227, room 160