



Terminal Pleistocene Alaskan genome reveals first founding population of Native Americans

Moreno-Mayar, J Víctor; Potter, Ben A; Vinner, Lasse; Steinrücken, Matthias; Rasmussen, Simon; Terhorst, Jonathan; Kamm, John A; Albrechtsen, Anders; Malaspinas, Anna-Sapfo; Sikora, Martin

Total number of authors:

18

Published in:

Nature

Link to article, DOI:

[10.1038/nature25173](https://doi.org/10.1038/nature25173)

Publication date:

2017

Document Version

Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Moreno-Mayar, J. V., Potter, B. A., Vinner, L., Steinrücken, M., Rasmussen, S., Terhorst, J., Kamm, J. A., Albrechtsen, A., Malaspinas, A-S., Sikora, M., Reuther, J. D., Irish, J. D., Malhi, R. S., Orlando, L., Song, Y. S., Nielsen, R., Meltzer, D. J., & Willerslev, E. (2017). Terminal Pleistocene Alaskan genome reveals first founding population of Native Americans. *Nature*, 553(7687), 203-207. <https://doi.org/10.1038/nature25173>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



LJMU Research Online

Moreno-Mayar, JV, Potter, BA, Vinner, L, Steinrücken, M, Rasmussen, S, Terhorst, J, Kamm, JA, Albrechtsen, A, Malaspinas, A-S, Sikora, M, Reuther, JD, Irish, JD, Malhi, RS, Orlando, L, Song, YA, Nielsen, R, Meltzer, DJ and Willerslev, E

Terminal Pleistocene Alaskan genome reveals first founding population of Native Americans

<http://researchonline.ljmu.ac.uk/7887/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Moreno-Mayar, JV, Potter, BA, Vinner, L, Steinrücken, M, Rasmussen, S, Terhorst, J, Kamm, JA, Albrechtsen, A, Malaspinas, A-S, Sikora, M, Reuther, JD, Irish, JD, Malhi, RS, Orlando, L, Song, YA, Nielsen, R, Meltzer, DJ and Willerslev, E (2018) Terminal Pleistocene Alaskan genome reveals first

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

1 **Terminal Pleistocene Alaskan genome reveals first founding population of Native**
2 **Americans**

3
4 **J. Víctor Moreno-Mayar**^{1,*}, **Ben A. Potter**^{2,*}, **Lasse Vinner**^{1,*} **Matthias Steinrücken**
5 ^{3,4,5}, **Simon Rasmussen**⁶, **Jonathan Terhorst**⁴, **John A. Kamm**^{4,7}, **Anders**
6 **Albrechtsen**⁸, **Anna-Sapfo Malaspinas**^{1,9,10}, **Martin Sikora**¹, **Joshua D. Reuther**²,
7 **Joel D. Irish**¹¹, **Ripan S. Malhi**^{12,13}, **Ludovic Orlando**¹, **Yun S. Song**^{3,4,14,15}, **Rasmus**
8 **Nielsen**^{1,4,14}, **David J. Meltzer**^{1,16} and **Eske Willerslev**^{1,7,17,**}.

9
10 1. Centre for GeoGenetics, Natural History Museum of Denmark, University of
11 Copenhagen, 1350 Copenhagen, Denmark.

12 2. Department of Anthropology, University of Alaska, Fairbanks, AK 99775.

13 3. Computer Science Division, University of California, Berkeley, CA 94720, USA.

14 4. Department of Statistics, University of California, Berkeley, CA 94720, USA.

15 5. Department of Biostatistics and Epidemiology, University of Massachusetts, Amherst,
16 MA 01003, USA.

17 6. Center for Biological Sequence Analysis, Department of Systems Biology, Technical
18 University of Denmark, 2800 Kongens Lyngby, Denmark.

19 7. Wellcome Trust Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge,
20 CB10 1SA, UK.

21 8. The Bioinformatics Centre, Department of Biology, University of Copenhagen, 2200
22 Copenhagen, Denmark.

23 9. Institute of Ecology and Evolution, University of Bern, CH-3012 Bern, Switzerland.

24 10. Swiss Institute of Bioinformatics, 1015 Lausanne, Switzerland.

25 11. Research Centre in Evolutionary Anthropology and Palaeoecology, Liverpool John
26 Moores University, Liverpool L3 3AF, UK.

27 12. Department of Anthropology, University of Illinois at Urbana-Champaign, Urbana,
28 IL 61801, USA.

29 13. Carle R. Woese Institute for Genomic Biology, University of Illinois at Urbana-
30 Champaign, Urbana, IL 61801, USA.

31 14. Department of Integrative Biology, University of California, Berkeley, CA 94720,
32 USA.

33 15. Department of Mathematics and Department of Biology, University of Pennsylvania,
34 PA 19104.

35 16. Department of Anthropology, Southern Methodist University, Dallas, TX 75275,
36 USA.

37 17. Department of Zoology, University of Cambridge, Downing St, Cambridge CB2 3EJ,
38 UK

39
40 * These authors contributed equally to this work.

41 ** Corresponding author: ewillerslev@snm.ku.dk

42
43 **Despite broad agreement that the Americas were initially populated via Beringia,**
44 **when and how this happened is debated**¹⁻⁵. **Key to this debate are human remains**
45 **from Late Pleistocene Alaska. The first and only such remains were recovered at**
46 **Upward Sun River (USR), and date to ~11.5 kya**^{6,7}. **We sequenced the USR1 genome**
47 **to an average coverage of ~17X. We find USR1 is most closely related to Native**
48 **Americans, but falls basal to all previously sequenced contemporary and ancient**
49 **Native Americans**^{1,8,9}. **As such, USR1 represents a distinct Ancient Beringian (AB)**

50 **population. Using demographic modelling we infer the AB population and ancestors**
51 **of other Native Americans descend from a single founding population that initially**
52 **split from East Asians $\sim 36 \pm 1.5$ kya, with gene flow persisting until $\sim 25 \pm 1.1$ kya.**
53 **Gene flow from ancient north Eurasians into all Native Americans took place 25-20**
54 **kya, with AB branching off $\sim 22-18.1$ kya. Our findings support long-term genetic**
55 **structure in ancestral Native Americans, consistent with the Beringian Standstill**
56 **Model¹⁰. We find that the basal Northern (NNA) and Southern (SNA) branches, to**
57 **which all other Native Americans belong, diverged $\sim 17.5-14.6$ kya, likely south of**
58 **the North American ice sheets. After 11.5 kya, some NNA populations received gene**
59 **flow from a Siberian population most closely related to Koryaks, but not**
60 **Paleoeskimos¹, Inuit or Kets¹¹, and that Native American gene flow into Inuit was**
61 **via NNA and not SNA groups¹. Our findings further suggest the far northern North**
62 **American presence of NNA is from a back migration that replaced or absorbed the**
63 **initial AB founding population.**

64
65 The peopling of the Americas, and particularly the population history of Beringia, the
66 land bridge that connected far northeast Asia to northwestern North America during the
67 Pleistocene, remains unresolved^{2,3}. Humans were present in the Americas south of the
68 continental ice sheets by ~ 14.6 kya¹², indicating they traversed Beringia earlier, possibly
69 around the Last Glacial Maximum (LGM). Then, the region was marked by harsh
70 climates and glacial barriers⁵, which may have led to the isolation of populations for
71 extended periods, and at times complicated dispersal across the region¹³. Still
72 controversial are questions of whether and how long Native American ancestors were
73 isolated from Asian groups in Beringia prior to entering the Americas^{2,10,14}; if one or
74 more early migrations gave rise to the founding population of Native Americans^{1-4,8,15}
75 (it is commonly agreed Paleoeskimos and Inuit represent separate and later migrations
76^{1,16,17}); and, when and where the basal split between SNA and NNA occurred. Unresolved
77 too is whether the genetic affinity between some SNA groups and indigenous
78 Australasians^{2,3}, reflects migration by non-Native Americans^{3,4,15}, early population
79 structure within the first Americans³, or later gene flow². Key to resolving these
80 uncertainties is a better understanding of the population history of Beringia, the entryway
81 for the Pleistocene peopling of the Americas.

82
83 Genomic insight into that population history has now become available with the recently
84 recovered infant remains (USR1 and USR2) from the Upward Sun River site, Alaska
85 (eastern Beringia), dated to ~ 11.5 kya^{7,18}. Mitochondrial DNA sequences (haplogroups
86 C1 and B2, respectively) were previously acquired from these individuals^{7,18} (SI 1,4,5).
87 We have since obtained whole-genome sequence data, which provides a broader
88 opportunity to investigate the number, source(s) and structure of the initial founding
89 population(s), and the timing and location of their subsequent divergence. We sequenced
90 the genome of USR1 to an average depth of $\sim 17X$, based on eight sequencing libraries
91 from USER-treated extracts previously confirmed to contain DNA fragments with
92 characteristic ancient DNA misincorporation patterns (SI 2-4). We estimated modern
93 human contamination at $\sim 0.14\%$ based on the nuclear genome and $\sim 0.15\%$ based on

94 mtDNA (SI 4). As expected, the error rate in the USER-treated sequencing data was low
95 (0.09% errors per-base), and comparable to other high-coverage contemporary genomes,
96 based on called genotypes (SI 4). While USR2⁷ did not show sufficient endogenous DNA
97 for high-coverage genome sequencing, we found both individuals were close relatives (SI
98 5), equally related to worldwide present-day populations (Figure S4g).

99
100 We assessed the genetic relationship between USR1, a set of ancient genomes^{2,8,9,15,17},
101 and a panel of 167 worldwide populations genotyped for 199,285 SNPs^{1,2,19} (SI 6), using
102 outgroup f_3 statistics²⁰, model-based clustering^{21,22} and multidimensional scaling (MDS)
103²³ (SI 7-9). Outgroup f_3 statistics of the form $f_3(\text{Yoruba}; X, \text{USR1})$ revealed that USR1 is
104 more closely related to present-day Native Americans than to any other tested population,
105 followed by Siberian and East Asian populations^{1,2} (Figure 1a). Pairwise comparisons of
106 the f_3 -statistics for USR1 and a set of ancient and contemporary Native American
107 genomes^{2,8,15} (SI 6) showed that all are similarly related to Old World populations,
108 though other Native American genomes (Aymara², Athabascan1¹⁶, 939², Anzick1⁸ and
109 Kennewick¹⁵) have a higher affinity for contemporary Native Americans than USR1 does
110 (SI 9). MDS and ADMIXTURE analysis showed that the USR1 genome did not cluster
111 with any specific Native American group (Figures 1d, S3b). These results imply that
112 USR1 belonged to a previously unknown Native American population not represented in
113 the reference dataset, herein identified as Ancient Beringians (SI 8.3).

114
115 To investigate if USR1 derived from the same source population that gave rise to
116 contemporary Native Americans, we computed 11,322 allele frequency based- D -
117 statistics^{1,20} of the form $D(\text{Native American}, \text{USR1}; \text{Siberian1}/\text{Han}, \text{Siberian2}/\text{Han})$ (SI
118 10.4). The resulting Z -score distribution corresponds qualitatively to the expected normal
119 distribution under the null hypothesis that USR1 forms a clade with Native Americans to
120 the exclusion of Siberians and East Asians – except for a set of Eskimo-Aleut, Athabascan
121 and Northern Amerind-speaking populations for which recent Asian gene flow has been
122 previously documented (Figures 1c, S5a, S6)^{1,2,15,19}. Additionally, we found that present-
123 day Native Americans and USR1 yield similar results for $D(\text{Native American}/\text{USR1},$
124 $\text{Han}; \text{Mal'Ta}, \text{Yoruba})$, suggesting they are equally related to the ancient north Eurasian
125 population represented by the 24 kya Mal'ta individual⁹ (SI 10.5). These results confirm
126 that USR1 and present-day Native Americans derived from the same ancestral source,
127 which carried a mixture of East Asian and Mal'ta-related ancestry. We infer that
128 descendants of this source represent the basal group that first migrated into the Americas.

129
130 To explore the relationship between USR1 and present-day Native Americans, we
131 computed allele frequency-based and genome-wide D -statistics of the form $D(\text{Native}$
132 $\text{American}, \text{Aymara}; \text{USR1}, \text{Yoruba})$. We could not reject the null hypothesis that USR1
133 is an outgroup to any pair of Native Americans, with the exception of a set of populations
134 bearing recent Asian gene flow^{1,2,15,19} (Figures 1b, S7). We confirmed the phylogenetic
135 placement of USR1 at a basal position in the Native American clade using TreeMix²⁴
136 and two methods to estimate average genomic divergence and genetic drift, respectively
137 (SI 14-16). These results support the branching of USR1 within the Native American

138 clade, but being equidistant to NNA and SNA. Below we discuss the potential geographic
139 locations of the USR1-NNA+SNA and the NNA-SNA splits (Figure 2) based on the
140 genetic results, the glacial geography of terminal Pleistocene North America^{25,26} and the
141 extant archaeological evidence (also SI 20).

142

143 Recent detection of an Australasian-derived genetic signature in some Native American
144 groups^{2,3} led us to explore whether USR1 bears that signal (SI 10.7, 11-13). Using
145 frequency-based and 'enhanced' D-statistics, we found no support for USR1 being closer
146 to Papuans (a proxy for Australasians) than other Native Americans.

147

148 We leveraged the position of USR1 on the Native American branch prior to the NNA-
149 SNA split to re-assess the origins of Athabascan and Eskimo populations by fitting
150 admixture graphs. We considered a whole-genome dataset including Siberian, East Asian,
151 Native American and Eskimo groups, as well as Mal'ta (SI 17). The heuristic approach in
152 TreeMix²⁴ showed that the best proxies for the Asian component in Athabascans and
153 Greenlandic Inuit are Koryaks and the Saqqaq individual, respectively. We then followed
154 an incremental approach for fitting an *f*-statistic-based admixture graph²⁰, including the
155 Kets, previously suggested to share a linguistic and perhaps a genetic link with
156 Athabascans^{11,27}. This approach recapitulated the TreeMix results, and yielded a model
157 in which both Athabascans and Greenlandic Inuit derive from the NNA branch. However,
158 the Asian ancestry in Athabascans is most closely related to the Asian component in
159 Koryaks, while the Saqqaq genome is the best proxy for the Siberian component in the
160 Greenlandic Inuit (Figure 3). We infer the latter is a consequence of Palaeo- and Neo-
161 Eskimos having been derived from a similar Siberian population^{1,16}. This model appears
162 to be a good fit to the data, as the observed *f*-statistic that deviated the most from the
163 model prediction yielded $Z=3.27$. In SI 17.3 we tested the robustness of this model and
164 predictions by computing individual *D* statistics, and re-fitting the model using alternative
165 datasets.

166

167 Lastly, we inferred the demographic history of USR1 with respect to Native Americans,
168 Siberians and East Asians, using two independent methods: *diCal2*²⁸ and *mom2*²⁹ (SI
169 18-19). *diCal2* results indicate that the founding population of USR1, Native Americans,
170 and Siberians had a very weak structure from ~36 kya up to ~24.5 kya (Table S7), when
171 the ancestors of USR1 and Native Americans began to diverge substantially from
172 Siberians. USR1 diverged from other Native Americans around 20.9 kya, with a period
173 of ensuing moderate gene flow between them (Table S6 and S7), as indicated by a
174 simulation study that showed a significant increase in likelihood when comparing a 'clean
175 split' model to an 'isolation with migration' model (SI 18.4). Using *mom2* and *SMC++*
176 we estimated a backbone demography where Karitiana and Athabascans split at ~15.7
177 kya, while their ancestral population split from Koryaks ~23.3 kya (Figure 4). With
178 *mom2*, we inferred the most likely branch (the population immediately ancestral to
179 NNA+SNA) and time (~21 kya) for the USR1 population to join the backbone
180 demography, while allowing for possible gene flow between USR and other populations
181 (SI 19, Figure 4b), results consistent with¹⁴ and the *diCal2* inference.

182

183 These new findings, along with existing data, allow us to place Ancient Beringians (AB)
184 within the broader context of the Pleistocene peopling of the Americas. The Native
185 American founding population (comprised of both AB and NNA+SNA) began to diverge
186 from ancestral Asians as early as ~36 kya, likely in northeast Asia, as there is no evidence
187 of people in Beringia or northwest North America at this period. A high level of gene
188 flow was maintained between them and other Asians until as late as ~25 kya^{2,14}. The
189 subsequent isolation of the Native American founding population ~24 kya roughly
190 corresponds with a decline in archaeological evidence for a human presence in Siberia³⁰.
191 Both changes may result from the same underlying cause: the onset of harsh LGM
192 climatic conditions². These findings, coupled with a divergence date of ~20.9 kya
193 between USR1 and Native Americans, are in agreement with the Beringian Standstill
194 Model¹⁰ (SI 21). The common ancestor of NNA+SNA and AB began to diverge ~20.9
195 kya, after which gene flow ensued, although whether it was with NNA+SNA, or the
196 already differentiated NNA and SNA branches, cannot be determined owing to shallow
197 divergence times among the groups.

198

199 These findings allow us to consider possible scenarios regarding where ancient Native
200 American populations diverged (SI 20-21, Figure 2). Scenarios C-E require extended
201 periods of strong population structure marking AB, NNA, and SNA as separate groups,
202 for which we do not see compelling genetic evidence; hence these can be rejected.
203 Scenarios A and B are compatible with our evidence of continuous gene flow among
204 these groups, but differ as to the location of the AB versus NNA+SNA split at 20.9 kya,
205 whether in northeast Asia (Scenario A) or eastern Beringia (Scenario B). Each has
206 strengths and weaknesses relative to genetic and archaeological evidence: Scenario A best
207 fits the archaeological and paleoecological evidence, as the earliest securely dated sites
208 in Beringia are no older than ~15-14 kya, and the LGM cold period is unlikely to be
209 associated with northward expanding populations³⁰. Scenario B is genetically most
210 parsimonious, given evidence of continuous gene flow between the AB and NNA+SNA,
211 suggesting their geographical proximity 20.9-11.5 kya, and that all three were isolated
212 from Asian/Siberian groups after ~24 kya and form a clade.

213

214 Scenarios A and B are both consistent with the NNA-SNA split at ~15 kya² having
215 occurred in a region south of eastern Beringia. The ice sheets were then still a significant
216 barrier to movement that would have helped maintain separation from the AB population.
217 While members of the SNA branch have not been documented in regions that were once
218 north of the glacial ice^{1,19}, NNA groups (including Athabascan-speakers) are present in
219 Alaska today; thus, the latter are likely descendants of a population that moved north
220 sometime after 11.5 kya²⁶.

221

222 The USR1 results provide the first direct genomic evidence that all Native Americans can
223 be traced back to the same source population from a single Late Pleistocene founding
224 event. Descendants of that population were present in eastern Beringia until at least 11.5
225 kya. By then, however, a separate branch of Native Americans had already established

226 itself in unglaciated North America, and diverged into the two basal groups that ultimately
227 became the ancestors of most of the indigenous populations of the Americas.

228

229 **Data availability**

230

231 Sequence data was deposited in the ENA under accession: PRJEB20398.

232

233 **Acknowledgements**

234

235 The Upward Sun River excavations and analysis were conducted under a Memorandum
236 of Agreement (MOA) signed by the State of Alaska, the National Science Foundation,
237 the Healy Lake Tribal Council, and the Tanana Chiefs Conference. We appreciate the
238 cooperation of all parties. We thank Morten Allentoft, Shyam Gopalakrishnan, Thorfinn
239 Korneliussen, Pablo Librado, Jazmín Ramos-Madrigal, Gabriel Renaud and Filipe Vieira
240 for discussions. We thank the Danish National High-throughput Sequencing Centre for
241 assistance in data generation. GeoGenetics members were supported by the Lundbeck
242 Foundation and the Danish National Research Foundation (DNRF94) and KU2016.
243 J.V.M.-M. was supported by Conacyt (Mexico). Samples were recovered during
244 excavations by B.A.P. supported by NSF Grants 1138811 and 1223119. Research
245 supported in part by NIH grant R01-GM094402 (M.S., J.T., J.A.K., and Y.S.S.) and a
246 Packard Fellowship for Science and Engineering (Y.S.S.). D.J.M. is supported by the
247 Quest Archaeological Research Fund. A.-S.M. is supported by the Swiss National
248 Science Foundation and the ERC.

249

250 **Author Contributions**

251 Project conceived by E.W. and B.A.P., and headed by E.W. and J.V.M.-M. L.V.
252 processed ancient DNA. J.V.M.-M. and S.R. assembled datasets. J.V.M.-M., M.S., J.T.,
253 J.A.K. and A.A. analysed genetic data. B.A.P. led the USR field investigation, and B.A.P.
254 and D.J.M. provided anthropological contextualization. B.A.P., J.D.R., and J.D.I.
255 conducted archaeological and bioanthropological work. R.N., Y.S.S., M.Si., A.-S.M., and
256 L.O. supervised bioinformatic and statistical analyses. B.A.P. engaged with indigenous
257 communities. J.V.M.-M., B.A.P., D.J.M and E.W. wrote the manuscript with input from
258 L.V., A.-S.M., M.Si., R.S.M., L.O., Y.S.S, R.N. and remaining authors.

259

260 **References**

261

- 262 1. Reich, D. *et al.* Reconstructing Native American population history. *Nature* **488**,
263 370–374 (2012).
- 264 2. Raghavan, M. *et al.* Genomic evidence for the Pleistocene and recent population
265 history of Native Americans. *Science* **349**, aab3884–aab3884 (2015).
- 266 3. Skoglund, P. *et al.* Genetic evidence for two founding populations of the
267 Americas. *Nature* (2015). doi:10.1038/nature14895
- 268 4. von Cramon-Taubadel, N., Strauss, A. & Hubbe, M. Evolutionary population
269 history of early Paleoamerican cranial morphology. *Sci. Adv.* **3**, e1602289 (2017).
- 270 5. Hoffecker, J. F., Elias, S. A., O'Rourke, D. H., Scott, G. R. & Bigelow, N. H.

- 271 Beringia and the global dispersal of modern humans: Beringia and the Global Dispersal
272 of Modern Humans. *Evol. Anthropol. Issues News Rev.* **25**, 64–78 (2016).
- 273 6. Potter, B. A., Irish, J. D., Reuther, J. D., Gelvin-Reymiller, C. & Holliday, V. T.
274 A Terminal Pleistocene Child Cremation and Residential Structure from Eastern
275 Beringia. *Science* **331**, 1058–1062 (2011).
- 276 7. Potter, B. A., Irish, J. D., Reuther, J. D. & McKinney, H. J. New insights into
277 Eastern Beringian mortuary behavior: A terminal Pleistocene double infant burial at
278 Upward Sun River. *Proc. Natl. Acad. Sci.* **111**, 17060–17065 (2014).
- 279 8. Rasmussen, M. *et al.* The genome of a Late Pleistocene human from a Clovis
280 burial site in western Montana. *Nature* **506**, 225–229 (2014).
- 281 9. Raghavan, M. *et al.* Upper Palaeolithic Siberian genome reveals dual ancestry of
282 Native Americans. *Nature* **505**, 87–91 (2013).
- 283 10. Tamm, E. *et al.* Beringian Standstill and Spread of Native American Founders.
284 *PLoS ONE* **2**, e829 (2007).
- 285 11. Flegontov, P. *et al.* Na-Dene populations descend from the Paleo-Eskimo
286 migration into America. (2016).
- 287 12. Dillehay, T. D. *et al.* Monte Verde: seaweed, food, medicine, and the peopling
288 of South America. *Science* **320**, 784–786 (2008).
- 289 13. Goebel, T. & Potter, B. A. First Traces: Late Pleistocene Human Settlement of
290 the Arctic. in *The Oxford handbook of the prehistoric Arctic* 223–252 (Oxford
291 University Press, 2016).
- 292 14. Llamas, B. *et al.* Ancient mitochondrial DNA provides high-resolution time
293 scale of the peopling of the Americas. *Sci. Adv.* **2**, e1501385–e1501385 (2016).
- 294 15. Rasmussen, M. *et al.* The ancestry and affiliations of Kennewick Man. *Nature*
295 (2015). doi:10.1038/nature14625
- 296 16. Raghavan, M. *et al.* The genetic prehistory of the New World Arctic. *Science*
297 **345**, 1255832–1255832 (2014).
- 298 17. Rasmussen, M. *et al.* Ancient human genome sequence of an extinct Palaeo-
299 Eskimo. *Nature* **463**, 757–762 (2010).
- 300 18. Tackney, J. C. *et al.* Two contemporaneous mitogenomes from terminal
301 Pleistocene burials in eastern Beringia. *Proc. Natl. Acad. Sci.* 201511903 (2015).
302 doi:10.1073/pnas.1511903112
- 303 19. Verdu, P. *et al.* Patterns of Admixture and Population Structure in Native
304 Populations of Northwest North America. *PLoS Genet.* **10**, e1004530 (2014).
- 305 20. Patterson, N. *et al.* Ancient Admixture in Human History. *Genetics* **192**, 1065–
306 1093 (2012).
- 307 21. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of
308 ancestry in unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).
- 309 22. Skotte, L., Korneliusen, T. S. & Albrechtsen, A. Estimating Individual
310 Admixture Proportions from Next Generation Sequencing Data. *Genetics* **195**, 693–702
311 (2013).
- 312 23. Malaspina, A.-S. *et al.* bammds: a tool for assessing the ancestry of low-depth
313 whole-genome data using multidimensional scaling (MDS). *Bioinformatics* **30**, 2962–
314 2964 (2014).
- 315 24. Pickrell, J. K. & Pritchard, J. K. Inference of Population Splits and Mixtures
316 from Genome-Wide Allele Frequency Data. *PLoS Genet.* **8**, e1002967 (2012).
- 317 25. Dyke, A. S., Moore, A. & Robertson, L. Deglaciation of North America. (2003).
- 318 26. Pedersen, M. W. *et al.* Postglacial viability and colonization in North America’s
319 ice-free corridor. *Nature* (2016). doi:10.1038/nature19085
- 320 27. Kari, J. M. & Potter, B. A. *The Dene-Yeniseian connection.* (University of

321 Alaska Department of Anthropology/Alaska Native Language Center, 2011).
322 28. Steinrücken, M., Kamm, J. A. & Song, Y. S. Inference of complex population
323 histories using whole-genome sequences from multiple populations. *bioRxiv* (2015).
324 doi:10.1101/093468
325 29. Kamm, J. A., Terhorst, J. & Song, Y. S. Efficient computation of the joint
326 sample frequency spectra for multiple populations. *J. Comput. Graph. Stat.* **26**, 182–194
327 (2016).
328 30. Goebel, T. The ‘microblade adaptation’ and recolonization of Siberia during the
329 late Upper Pleistocene. *Archeol. Pap. Am. Anthropol. Assoc.* **12**, 117–131 (2002).

330
331

332 **Figure 1. Genetic affinities between USR1, present-day Native Americans, and**
333 **world-wide populations. a.** f_3 statistics of the form $f_3(\text{San}; X, \text{USR1})$, for each
334 population in the genotype panel. Warmer colors represent greater shared drift between a
335 population and USR1. **b.** D -statistics of the form $D(\text{Native American}, \text{Aymara}; \text{USR1},$
336 $\text{Yoruba})$ (points). The Andean Aymara were used to represent SNA. *: Native American
337 populations with Asian admixture ($|Z|$ for $D(\text{HI}, \text{Aymara}; \text{Han}, \text{Yoruba}) > 3.3$) (Figure
338 S5a). Error bars represent 1 and ~ 3.3 standard errors (p -value ~ 0.001). Native American
339 populations were grouped by language family ¹. **c.** Quantile-quantile plot comparing
340 observed Z -scores to the expected normal distribution under the null hypothesis (H_0), for
341 all possible $D(\text{Nat. Am.}, \text{USR1}; \text{Siberian1}, \text{Siberian2})$. Colors correspond to the Z -score
342 obtained for $D(\text{HI}, \text{Aymara}; \text{Han}, \text{Yoruba})$. The expected normal distribution under the
343 null hypothesis was computed for all groups jointly (SI Section 10.4). Thick and thin lines
344 represent a Z -score of ~ 3.3 (p -val ~ 0.001) and a Z -score of ~ 4.91 (p -val ~ 0.01 after
345 applying a Bonferroni correction for 11,322 tests). The bottom-right panel shows the
346 expected tree under the null hypothesis. **d.** Admixture proportions estimated by
347 *ADMIXTURE* ³⁷ assuming $K=20$ ancestral populations. Bars represent individuals, and
348 colors represent admixture proportions from each ancestral component. Admixture
349 proportions in ancient genomes (wider bars) were estimated using a genotype likelihood-
350 based approach ³⁸.

351

352 **Figure 2. Possible geographic locations for the USR1 and NNA-SNA splits.** We
353 propose two possible locations for the split between USR1 and other Native Americans:
354 the Old World (A, C, E) and Beringia (B, D); and three possible locations for the
355 NNA_SNA split: the Old World (E), Beringia (C, D), and North America south of
356 Beringia (A, B). Schematics show estimated glacial extent ~ 14.8 kya. Dashed lines
357 represent the Native American migration south of eastern Beringia, but they do not
358 correspond to a specific migration route Model discussion (SI 20) is based on extant
359 archaeological evidence and inferred demographic parameters: a USR1-NNA+SNA split
360 ~ 20 kya with ensuing moderate gene flow and a NNA-SNA split ~ 15 kya (SI 18-19)..

361

362 **Figure 3. A model for the formation of the different Native American populations.**
363 We fitted an admixture graph by sequentially adding admixed leaves to a 'seed' graph
364 including the Yoruba, Han, Mal'ta, Ket, USR1, Anzick1 and Aymara genomes. For each
365 'non-seed' admixed group, we found the pair of edges that produced the best-fitting graph,

366 based on the fitting and maximum $|Z|$ scores (3.27 for this graph). Ellipse-shaped nodes:
367 sampled populations; box-shaped nodes: metapopulations; *: single high-depth ancient
368 genome. **: single low-depth genome. †: subgraphs whose structure we were unable to
369 resolve due to sequencing and genotyping error in the Saqqaq genome (SI 17). Sample
370 sizes and locations are shown at the top.

371

372 **Figure 4. USR1 demographic history in the context of East Asians, Siberians and**
373 **other Native Americans. a.** *SMC++* inferred effective population sizes with respect to
374 time for Athabascans (NNA), Karitiana (SNA), Han, Koryaks and USR1 (SI 19.1). We
375 used these demographic histories as a basis for fitting a joint model for these populations.
376 **b.** A ‘backbone demography’ was fitted excluding USR1 using *mom2*, an SFS-based
377 maximum likelihood approach (Figure S27), along with the most likely join-on point for
378 USR1 onto the backbone demography (SI 19). We show the likelihood heatmap for the
379 latter; warmer colors correspond to a higher likelihood of USR1 joining at a given point.
380 These estimates agree with those obtained through *diCal2*, a method based on haplotype
381 data (SI 18).

382

383