



Optimizing healthy and safe fish intake recommendations: a trade-off between personal preference and cost

Persson, Maria; Fagt, Sisse; Nauta, Maarten J

Published in:
British Journal of Nutrition

Link to article, DOI:
[10.1017/S0007114519000989](https://doi.org/10.1017/S0007114519000989)

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Persson, M., Fagt, S., & Nauta, M. J. (2019). Optimizing healthy and safe fish intake recommendations: a trade-off between personal preference and cost. *British Journal of Nutrition*, 122(2), 206-219.
<https://doi.org/10.1017/S0007114519000989>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Optimising healthy and safe fish intake recommendations: a trade-off between personal preference and cost

Maria Persson*, Sisse Fagt and Maarten J. Nauta

Division of Diet, Disease Prevention and Toxicology, National Food Institute, Technical University of Denmark, Kgs. Lyngby, Denmark

(Submitted 8 November 2018 – Final revision received 8 April 2019 – Accepted 10 April 2019; First published online 24 June 2019)

Abstract

Individuals may perceive personalised dietary advice as more relevant and motivational than national guidelines. Personal preference and food cost are factors that can affect consumer decisions. The objective of this study was to present a method for modelling and analysing the trade-off between deviation from preference and food cost for optimised personalised dietary recommendations. Quadratic programming was applied to minimise deviation from fish preference and cost simultaneously with different weights on the cost for 3016 Danish adults (whose dietary intake and body weight were recorded in a national dietary survey). Model constraints included recommendations for EPA, DHA and vitamin D and tolerable levels for methyl mercury, dioxins and dioxin-like polychlorinated biphenyls. When only minimising deviation from preference, 50 % of the study population should be recommended to increase fish intake, 48 % should be suggested to maintain current consumption and 2 % should be suggested to decrease fish consumption. When only minimising cost, the vast majority (99 %) should be recommended to only consume herring, which is the least-expensive fish species. By minimising deviation from preference and cost simultaneously with different weights on the cost, personalised optimal trade-off curves between deviation from fish intake preference and fish cost could be generated for each individual in our study population, except for twenty-two individuals (0.7 %) whose contaminant background exposure was too high. In the future, the method of this paper could be applied in the personal communication of healthy and safe food recommendations that fit the preferences of individual consumers.

Key words: Personalised intake recommendations: Quadratic programming: Optimal trade-off curves: Personal food preference: Cost

National food-based dietary guidelines (FBDG) are developed to help consumers make healthy food choices and to improve public health⁽¹⁾. For example, the official Danish FBDG directed at the healthy population over 3 years of age recommend an intake of 350 g of fish per week, of which 200 g should be fatty fish⁽²⁾. If the population would follow these FBDG, a considerable public health gain is expected⁽³⁾. However, according to national dietary survey data (*n* 3016), only about 15 % of the Danish adult population follow the national recommendation for fish⁽⁴⁾. Furthermore, the variation in fish intake between individuals is large, partly due to the fact that 11 % did not report consumption of fish. The compliance to other Danish FBDG is low as well; for example, reducing saturated fat, it is as low as 3 %, and for increasing dietary fibre, 16 % of the adult Danes comply with the recommendation⁽⁵⁾.

The proportion of the Danes that are aware of the FBDG is unknown, and among the aware individuals, there is likely a range of factors limiting their adherence. The relationship

between awareness of dietary recommendations and behaviour change is complicated by several components⁽¹⁾. Food choice is a complex behaviour that is influenced by many interacting factors, and the understanding of the determinants of food choice is limited^(6,7).

Food cost is known to influence diet quality^(8–11). Groups with lower incomes are less likely to make food purchasing choices consistent with dietary guideline recommendations^(12,13). Hence, fish cost may make the Danish national recommendation for fish unattractive and maybe not feasible for some individuals in the population. Another possible, more general reason for the low adherence (15 %) may be that the recommendation deviates too much from the current fish consumption for a large part of the population. Consumers may not be motivated to follow national FBDG if they differ a great deal from individual food preferences.

Since national guidelines have limited effect^(1,5,14), alternative and/or complimentary ways to improve public health by

Abbreviations: DKK, Danish krone; dl-PCB, dioxin-like polychlorinated biphenyls; FBDG, food-based dietary guidelines; LP, linear programming; QP, quadratic programming; TEQ, toxic equivalency.

* **Corresponding author:** M. Persson, email marper@food.dtu.dk

modifying consumption patterns are relevant. Previous studies suggests that consumers may perceive personalised dietary advice as more relevant and motivational than 'one size fits all' generic guidelines since they can take personal characteristics such as preference and needs into account⁽¹⁵⁻¹⁷⁾.

Mathematical optimisation of food intakes ('diet optimisation') have been applied to analyse several aspects of diet sustainability⁽¹⁸⁾. Mathematical optimisation methods are suitable for this purpose, thanks to their ability to deal with several factors simultaneously; they provide a multi-dimensional approach. Dimensions of diet sustainability include environmental impact, nutritional adequacy, affordability, cultural acceptability and accessibility, for example. The goal of a diet optimisation problem is to optimise a combination of food items subject to some predefined criteria. Diet modelling has been applied at individual⁽¹⁹⁻²²⁾ and population level⁽²³⁻²⁵⁾. The majority of the previous diet optimisation studies have considered nutritional recommendations as criteria, and few studies also included contaminant exposure limits⁽¹⁸⁾. Several diet optimisation studies have assessed the cultural acceptability dimension of diet sustainability by modelled combinations of foods that deviate as little as possible from observed intake⁽¹⁹⁻²⁴⁾. The advantage of such food combinations is that they minimise the behavioural changes necessary. Smaller deviations from observed intakes may be more culturally acceptable than larger deviations, but no formal definition of minimum deviation from observed intake exists⁽¹⁸⁾. Hence, different methods to minimise this behavioural change can be applied. Linear programming (LP) and quadratic programming (QP) are two alternative approaches for minimising behavioural change subject to a set of constraints, giving different outputs. Diet affordability has been analysed in previous diet optimisation studies, and it has been concluded that a cost constraint can impact diet quality negatively^(9,26).

This study presents a novel model for generating personalised fish intake recommendations by accounting for two factors that are likely to affect consumption: personal preference and fish cost. These two factors are minimised simultaneously subject to constraints on nutrients and contaminants. Hence, the modelled individual combinations of fish species may be perceived as more sustainable than national FBDG, since the model accounts for cultural acceptability and affordability while considering nutritional adequacy and contaminant limits. Fish consumption provides health benefits mainly due to its content of the nutrients such as EPA, DHA and vitamin D, but fish consumption also constitutes health risks, mainly due to the presence of the contaminants such as methyl mercury, dioxins and dioxin-like polychlorinated biphenyls (dl-PCB) in fish⁽²⁷⁾. In the previous studies, we modelled personalised fish intake recommendations by applying a QP model that minimises the deviation from personal preference (defined as observed individual intake), while ensuring that constraints on these nutrients and contaminants are fulfilled^(21,22). Hence, personalised recommendations for fish intake that differed as little as possible from individual preference were generated. The results suggest that an intake of 350 g of fish per week, of which 200 g should be fatty fish, as recommended by the Danish food authorities, is not necessary^(21,22). These studies concluded that meeting the official recommendation is healthy and safe, but does require

larger behaviour changes in the population than necessary, which may lead to lack of compliance. This suggests that personalised recommendations for fish can work as compliments to the FBDG in favour for improving healthier consumption patterns. In this study, the food choice determinant cost was also modelled and analysed.

The objective of this study was to present a method for modelling and analysing the trade-off between deviation from preference (defined as observed individual intake) and cost for optimised personalised intake recommendations. As this is a novel method, we wish to point out that this study is first and foremost methodological. Fish intake in Denmark demonstrates the use of the model.

Methods

Data

Observed intake and body weights. The observed intake (7-d estimated records) and self-reported body weights were obtained from the Danish National Survey of Diet and Physical Activity (DANSDA)⁽⁴⁾ (n 3016, aged 18–75 years). We excluded the species garfish and saithe from the optimisation variable, which were less than 2 % of the individuals reported. Garfish and saithe were instead defined as background exposure foods. The remaining nine reported species were included, denoting the nine elements of the optimisation variable in the model. The observed intake data are shown in [Table 1](#). The observed intakes of five representative individuals in the study population with different personal preference are shown in [Table 2](#). Other than to some extent having a clear preference (further explained in the 'Results' section), the individuals were arbitrarily chosen. Two of these individuals already fulfilled their nutrient and contaminant constraints, that is, their observed intakes were within the feasible region of the model.

Nutrient compositions, contaminant concentrations and limit values. The average nutrient compositions (EPA + DHA and vitamin D) and the average contaminant concentrations (methyl mercury and dioxins + dl-PCB) for the fish species were calculated as in our previous study⁽²²⁾ (see [Table 3](#)). Nutrient compositions for species subcategories such as raw, smoked, canned and marinated were extracted from a Danish food composition database⁽²⁸⁾. Nutrient compositions for the fish species were calculated as weighted averages of the extracted data with weights equal to the reported intake amounts of the different subcategories. Contaminant concentrations for fish subcategories were extracted from two Danish chemical contaminant reports^(29,30). The same procedure as for the nutrients was applied to calculate the contaminant concentrations, but since there were two contaminant reports, the extra step of calculating the weighted averages of the two contaminant reports with the number of samples per report serving as weights was performed. The conservative approach regarding methyl mercury as used by the European Food Safety Authority (EFSA)⁽³¹⁾ was applied: 100 % of the total mercury in fish was considered as methyl mercury, and 80 % of the total mercury in seafood other than fish was considered as methyl mercury.

Table 1. Observed fish intake⁽⁴⁾ and prices of fish (*n* 3016, aged 18–75 years) (Numbers of individuals with reported intake (*n_r*), mean values and standard deviations; medians and interquartile ranges (IQR))

	Women (<i>n</i> 1552)					Men (<i>n</i> 1464)					Mean price (DKK/kg)
	<i>n_r</i>	Intake (g/week)				<i>n_r</i>	Intake (g/week)				
		Mean	SD	Median	IQR		Mean	SD	Median	IQR	
Total fish intake	1397	188	186	144	228	1272	235	252	165	311	176
Lean fish (≤5 % fat)	1108	80	107	36	120	1039	102	150	45	159	169
Cod (raw)	591	25	56	0.0	22	545	30	69	0.0	26	210
European plaice (raw)	408	25	66	0.0	9.7	387	34	101	0.0	9.7	173
Tuna (canned)	753	21	49	0.0	15	698	25	64	0.0	19	126
European flounder (raw)	233	7.6	24	0.0	0.0	242	11	30	0.0	0.0	142
Fatty fish (>5 % fat)	1231	108	138	58	161	1089	134	191	50	197	180
Salmon (raw, smo)	924	41	68	8.6	54	728	42	77	0.0	45	274
Herring (mar, raw, smo)	860	31	63	1.4	38	783	49	103	0.72	54	90
Mackerel (can, smo, raw)	947	23	40	9.2	33	832	31	57	9.2	37	139
Trout (raw)	355	11	24	0.0	0.0	270	11	29	0.0	0.0	229
Greenland halibut (raw, smo)	487	1.4	5.7	0.0	1.5	374	1.8	12	0.0	0.63	379

DKK, Danish krone; smo, smoked; mar, marinated.

Table 2. Observed fish intake⁽⁴⁾ and observed total background nutrient intake/contaminant exposure for five selected individuals in the study population

	Observed intake inside feasible region		Observed intake outside feasible region		
	Individual 1	Individual 2	Individual 3	Individual 4	Individual 5
	Total fish intake (g/week)	355	185	1754	0.0
Lean fish (≤5 % fat) (g/week)	220	0.0	1112	0.0	218
Cod (raw) (g/week)	14	0.0	0.0	0.0	70
European plaice (raw) (g/week)	20	0.0	1049	0.0	20
Tuna (canned) (g/week)	145	0.0	64	0.0	88
European flounder (raw) (g/week)	42	0.0	0.0	0.0	42
Fatty fish (>5 % fat) (g/week)	135	185	642	0.0	0.0
Salmon (raw, smo) (g/week)	85	118	0.0	0.0	0.0
Herring (mar, raw, smo) (g/week)	3.5	2.9	418	0.0	0.0
Mackerel (can, smo, raw) (g/week)	17	23	224	0.0	0.0
Trout (raw) (g/week)	28	38	0.0	0.0	0.0
Greenland halibut (raw, smo) (g/week)	2.1	2.9	0.0	0.0	0.0
Total background intake/exposure					
EPA + DHA (g/week)	0.49	0.62	0.73	0.15	1.0
Vitamin D (µg/week)	214	63	76	65	79
Methyl mercury (µg/week)	1.2	2.3	2.6	0.0	2.9
Dioxins + dl-PCB (pg TEQ/week)	289	299	393	229	303

smo, Smoked; mar, marinated; dl-PCB, dioxin-like polychlorinated biphenyls; TEQ, toxic equivalency.

Recommended daily intakes for the nutrients and tolerable weekly intakes for the contaminants (see Table 4) were used for calculating limit values for the constraints in the model. The recommendations and tolerable weekly intakes are regarding total exposure, and therefore background exposure amounts are subtracted. Daily values are converted to weekly values in the model by multiplying daily recommendations by seven, and per-body-weight values are converted to per-individual values in the model by multiplication with individual body weight, as available from the data.

Prices of fish. The average price of each fish species product (raw, smoked, canned or marinated) was calculated from prices collected from three Danish online supermarkets (nemlig.com, mad.coop.dk and osuma.dk) and two Danish online fish stores (fiskehuset.dk and fish4you.dk) in April 2018. The prices of the fish species in the model (see Table 1) were calculated as weighted

averages of the fish products, with the mean reported intake of the different products serving as weights. This means that each fish product was assigned a weight determining its relative importance for the total price of the corresponding fish species. The prices for lean and fatty fish, as well as all fish, were calculated as weighted averages of the fish species, with the mean intake of the study population for the different species serving as weights.

Background exposure

Fish is not the only source of the nutrients and contaminants included in the model. Background exposure can be due to consumption of several other foods than fish, dietary supplements and potentially also by other environmental sources. Background exposure amounts were subtracted from the recommendations for EPA + DHA, vitamin D, methyl mercury and dioxins + dl-PCB, as in our previous study⁽²²⁾. The background

Table 3. Nutrient compositions and contaminant concentrations for fish^(28–30)

	EPA+DHA (mg/g)	Vitamin D (µg/g)	Methyl mercury (µg/g)	Dioxins dl-PCB (pg TEQ/g)
Lean fish (≤5 % fat)				
Cod (raw)	2.2	0.010	0.045	0.13
European plaice (raw)	6.0	0.011	0.035	0.31
Tuna (can)	2.0	0.027	0.151	0.05
European flounder (raw)	4.2	0.0080	0.035*	0.65
Fatty fish (>5 % fat)				
Salmon (raw, smo)	16	0.079	0.011	0.81
Herring (mar, raw, smo)	18	0.095	0.037	1.2
Mackerel (can, smo, raw)	26	0.044	0.28	1.0
Trout (raw)	14	0.16	0.023	0.38
Greenland halibut (raw, smo)	8.0	0.048	0.057	0.56

dl-PCB, dioxin-like polychlorinated biphenyls; TEQ, toxic equivalency; smo, smoked; mar, marinated.

* Plaice data.

Table 4. Recommendations for nutrients and contaminants

	Value	Reference
Recommended daily intake		
EPA + DHA (mg/g)	250	(43)
Vitamin D (µg/g)	10	(44)
Tolerable weekly intake		
Methyl mercury (µg/kg BW per week)	1.3	(31)
Dioxins + dl-PCB (pg TEQ/kg BW per week)	14	(45)

BW, body weight; dl-PCB, dioxin-like polychlorinated biphenyls; TEQ, toxic equivalency.

exposure sources included were foods other than fish, vitamin D supplements, sun and air. For foods other than fish and vitamin D supplements, individual background exposure was calculated from observed intake data⁽⁴⁾ combined with composition and concentration data^(28–30) (see Table 5). No EPA + DHA supplementation data were available, and we decided to not include EPA + DHA supplementation in the model.

For the environmental background exposure, no individual data were available. Therefore, estimated average values calculated in our previous study⁽²²⁾ were used for all individuals in the study population (see Table 5). For EPA + DHA, there is no environmental background exposure. For vitamin D, UVB radiation from the sun (that gets synthesised in the skin) is a significant source in Denmark^(32,33). The seasonal variation of the radiation is large; the highest level is in summer, and the lowest in winter. In this study, a so-called ‘mid-season’ scenario is used, which is the average of a summer scenario and a winter scenario⁽²²⁾. For methyl mercury exposure, fish and seafood consumption is the major source^(31,34). We assumed food to be the only source of exposure which was considered appropriate since we used conservative assumptions for the methyl mercury concentrations in fish. For dioxins, food consumption contributes to more than 90 % of the total exposure⁽³⁵⁾. We assumed that the remaining 10 % are from air and calculated an estimated average airborne dioxin exposure using this 9:1 relationship and inserting the average exposure due to food consumption (376 pg toxic equivalency (TEQ) per week). The total background exposure for five representative individuals in the study population is shown in Table 2.

Model overview

Quadratic programming model. The QP model of the study is expressed as

$$\underset{x}{\text{minimize}} \quad x - x_{\text{obs}}2 + \lambda Cx, \quad (1)$$

$$\text{subject to } Bx \geq b, \quad (2a)$$

$$Rx \leq r, \quad (2b)$$

$$x \geq 0, \quad (2c)$$

where the vector x ($d \times 1$) is the optimisation variable describing weekly intake amounts of d number of different fish species (g/week) for one individual, the vector x_{obs} ($d \times 1$) describes the individual observed intake amounts of the corresponding fish species (g/week), the parameter λ (≥ 0) is a constant that determines the relative importance of the cost of fish intake, the vector C ($1 \times d$) describes the prices of the different fish species (Danish krone; DKK/g), the matrices B ($m \times d$) and R ($k \times d$) denote the average of the nutrient compositions and contaminant concentrations in fish (mg/g, µg/g or pg TEQ/g), respectively, and the vectors b ($m \times 1$) and r ($k \times 1$) describe the weekly lower and upper intake amounts of the nutrients and contaminants (mg/week, µg/week, µg/kg body weight per week or pg TEQ/kg body weight per week).

The solution of this optimisation problem, which is a fish intake recommendation for the individual, is the vector x that fulfils the constraints (equations 2a–2c) and minimises the objective function (equation 1). The model is nine-dimensional, since nine fish species are considered ($d = 9$). A QP model punishes large deviations and may make small changes in all nine elements in the optimisation variable. Our assumption is that many smaller are realistic and achievable for consumers, which corresponds to the minimum total deviation possible (the shortest distance between the model recommendation and the observed intake, as explained below).

Objective function. The objective function of an optimisation problem chooses the optimal solution for x among the feasible solutions. In our model, the vector x_{obs} ($d \times 1$) of the objective

Table 5. Observed nutrient intake and contaminant exposure* (Mean values and standard deviations; medians and interquartile ranges (IQR))

	Women (n 1552)				Men (n 1464)			
	Mean	SD	Median	IQR	Mean	SD	Median	IQR
Intake/exposure from all foods								
EPA + DHA (g/week)	2.8	3.2	1.8	3.5	3.4	4.1	1.9	4.3
Vitamin D (µg/week)	28	20	23	19	35	24	29	23
Methyl mercury (µg/week)	11	13	8.2	13	15	18	8.9	17
Dioxins + dl-PCB (pg TEQ/week)	326	306	265	220	428	303	346	275
Intake/exposure from foods other than fish								
EPA + DHA (g/week)	0.40	0.92	0.23	0.35	0.44	0.87	0.25	0.42
Vitamin D (µg/week)	18	14	15	10	22	16	19	13
Methyl mercury (µg/week)	1.0	2.2	0.11	1.1	0.98	2.6	0.055	0.89
Dioxins + dl-PCB (pg TEQ/week)	211	251	178	98	278	199	247	135
Exposure from supplements								
Vitamin D (µg/week)	65	96	33	93	39	66	0.0	70
Exposure from environment								
Sun: vitamin D (µg/week)	51				51			
Air: dioxins + dl-PCB (pg TEQ/week)	42				42			

dl-PCB, dioxin-like polychlorinated biphenyls; TEQ, toxic equivalency.

* Observed intake⁽⁴⁾ multiplied with nutrient composition and contaminant concentration data^(28–30), supplement intake⁽⁴⁾ and estimated average environmental exposure (n 3016, aged 18–75 years).

function (equation 1) is a constant vector with the observed fish intake amounts (g/week) of an individual, which is defined as individual preference. The vector C ($1 \times d$) is a constant vector with the prices of the different fish species (DKK/g). The L_2 -norm of the difference between the optimisation variable and the observed intake, $x - x_{\text{obs}2}$, describes the deviation from the observed intake. This is a straight-line distance, which is the shortest distance between the modelled recommendation and the observed intake, a distance in a d -dimensional space ($d = 9$ in our case):

$$x - x_{\text{obs}2} = \sqrt{|x_1 - x_{\text{obs},1}|^2 + |x_2 - x_{\text{obs},2}|^2 + \dots + |x_n - x_{\text{obs},d}|^2} \quad (3)$$

The product Cx is the total weekly cost of fish intake (DKK/week) for the individual when the amount of fish described by vector x is consumed. Hence, the objective function (equation 1) of the model minimises the deviation from the individual observed fish intake (the preference) and the corresponding cost of fish intake simultaneously. The parameter λ (≥ 0) is a constant that determines the relative importance of the cost for the optimisation. With $\lambda = 0$, the model is identical with the model from our previous studies in which the cost was not considered, but only deviation from personal preference was minimised^(21,22). The model of this study can be modified to minimise only cost of fish intake by eliminating the L_2 -norm. Then, the following alternative objective function is used:

$$\underset{x}{\text{minimize}} Cx. \quad (4)$$

With this cost objective function, the model becomes an LP instead of a QP. When $\lambda \rightarrow \infty$ in the original model, its objective function becomes equal to the cost objective function.

Feasible region. The feasible region of an optimisation problem is the set of all possible (feasible) solutions to the problem

from which the objective function chooses the optimal solution. It is the set of vectors that fulfil the constraints. The constraints of our model ensure that the modelled recommendation for an individual meets the lower limits on the nutrients EPA + DHA and vitamin D (equation 2a) without violating upper limits on the contaminants methyl mercury and dioxins + dl-PCB (equation 2b), and makes sure that no modelled recommendations are negative (equation 2c). The vector b ($m \times 1$) describes the weekly lower limits for the nutrient intake amounts due to fish intake ($m = 2$: EPA + DHA and vitamin D), and r ($k \times 1$) describes the weekly upper limits for the contaminant intake amounts ($k = 2$: methyl mercury and dioxins + dl-PCB). The elements of limit vector b are recommended intakes for the nutrients subtracted with the individual background intake. For vector r , the elements are tolerable intakes for the contaminants minus the individual amount due to background exposure. The matrices B ($m \times d$) and R ($k \times d$) denote the nutrient compositions and contaminant concentrations in fish, respectively. The feasible region of an individual, made up by her/his constraints, is body weight dependent (due to the contaminant constraint (equation 2b) and also dependent on her/his background exposure (due to the nutrient and contaminant constraints (equations 2a and 2c).

Optimal trade-off curve. In general, there is a trade-off between deviation from personal preference $x - x_{\text{obs}2}$ and cost Cx , when an individual's preference is not the combination of fish species that minimises the cost while fulfilling the model constraints. Analysing the trade-off between these two implies analysing how much an individual has to deviate from her/his preference if a lower cost of consumption is desired. We achieved an optimal trade-off curve between deviation from preference and cost by defining different parameter values λ and running the QP model with these, whereby λ parametrises the optimal trade-off curve. We

Table 6. Observed fish intakes⁽⁴⁾ and modelled fish intake recommendations with different objective functions and corresponding costs (Mean values and standard deviations; medians and interquartile ranges (IQR); minimum (Min) and maximum (Max) values)

Observed fish intake	Mean	SD	Median	IQR	Min	Max	Mean	SD	Median	IQR	Min	Max
	Women (n 1552)						Men (n 1464)					
Observed lean fatty fish intake (g/week)	78 108	106 138	35 58	114 161	0.0 0.0	763 1015	100 134	147 191	43 50	157 197	0.0 0.0	1574 1706
Cost of observed lean fatty fish intake (DKK/week)	13 20	18 28	5.7 10	19 29	0.0 0.0	155 179	17 23	26 34	6.3 8.5	25 34	0.0 0.0	289 263
Modelled fish intake	Mean	SD	Median	IQR	Min	Max	Mean	SD	Median	IQR	Min	Max
	Minimise deviation from personal reference ($\lambda = 0$)											
All women with feasible recommendation (n 1539)						All men with feasible recommendation (n 1455)						
Modelled recommended lean fatty fish intake (g/week)	82 134	102 109	41 89	106 90	0.0 0.0	763 781	104 156	142 153	45 88	146 128	0.0 0.0	1574 1048
Ratio of lean fish/(lean fish + fatty fish)	0.33	0.27	0.27	0.42	0	1	0.35	0.29	0.27	0.43	0	1
Cost of modelled lean fatty fish intake (DKK/week)	14 25	18 23	6.3 17	18 17	0.0 0.0	155 179	17 28	25 29	6.8 17	24 21	0.0 0.0	289 213
Women with recommended increase (n 788)						Men with recommended increase (n 723)						
Modelled recommended increase* in lean fatty fish intake (g/week)	10 58	4.6 29	10 62	7.6 45	0.0 0.2	20 153	10 58	4.4 27	11 62	7.3 42	0.0 0.1	19 148
Minimise cost, alternative cost objective function												
All women with feasible recommendation (n 1539)						All men with feasible recommendation (n 1455)						
Modelled recommended lean fatty fish intake (g/week)	0.0 82	0.0 22	0.0 86	0.0 19	0.0 0.0	0.0 181	0.0 79	0.0 22.6	0.0 85	0.0 22.7	0.0 0.0	0.0 188
Cost of modelled lean fatty fish intake (DKK/week)	0.0 7.4	0.0 2.0	0.0 7.7	0.0 1.7	0.0 0.0	0.0 19	0.0 7.1	0.0 2.0	0.0 7.6	0.0 2.0	0.0 0.0	0.0 17

DKK, Danish krone.

* Modelled recommendation minus observed intake.

Trade-off between personal preference and cost




Fig. 1. Deviation from personal preference is minimised: empirical cumulative distribution functions for the suggested changes in fish intake (modelled recommendation minus observed intake) for the study population with $\lambda = 0$. Total lean fish and total fatty fish (a) and all nine species (b). The value on the y-axis corresponding to a specified value of change in fish intake on the x-axis is the fraction of individuals in the study population that should be suggested to make a change less than or equal to the specified value on the x-axis. Original to this manuscript.




Fig. 2. Cost is minimised: empirical cumulative distribution functions for the suggested changes in fish intake (modelled recommendation minus observed intake) for the study population with the alternative cost objective function (equation 4). Total lean fish and total fatty fish (a) and all nine species (b). The value on the y-axis corresponding to a specified value of change in fish intake on the x-axis is the fraction of individuals in the study population that should be suggested to make a change less than or equal to the specified value on the x-axis. Original to this manuscript.

used the arbitrary numbers $\lambda = 0, 0.2, 0.4, 0.6, 0.8, \dots$ up to a number that minimises the cost for each individual. For comparison, the model was also run with the cost objective function (equation 4) that only minimises the cost of fish intake.

Software

To solve the QP problem, we used CVX, a Matlab (R2015b, version 8.6) package for specifying and solving convex programs^(36,37).

Results

National food-based dietary guidelines and current consumption

The cost of consuming 350 g of fish per week, of which 200 g is fatty fish, as recommended in the official Danish FBDG, is 61 DKK per week. Currently, the average woman in Denmark consumes 186 (78 + 108) fish per week and the average Danish man consumes 234 g (100 + 134) fish per week, which cost 33 (13 + 20) and 40 (17+23) DKK per week, respectively (see Table 6).

Table 7. Modelled fish intake recommendations with different values for λ for five selected individuals in the study population

	Individual 1				Individual 2				Individual 3				Individual 4				Individual 5			
λ	0.0	2.0	4.4	71.4	0.0	2.6	6.2	14.0	0.0	4.8	5.0	6.0	0.0	1.4	3.0	174.6	0.0	3.4	6.0	57.0
Cod (g/week)	14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.2	0.0	0.0	0.0	70	56	0.0	0.0
European plaice (g/week)	20	6.6	0.0	0.0	0.0	0.0	0.0	0.0	993	504	230	0.0	6.1	0.0	0.0	0.0	20	11	0.0	0.0
Tuna (g/week)	145	135	86	0.0	0.0	0.0	0.0	0.0	55	0.0	0.0	0.0	2.0	0.0	0.0	0.0	88	81	7	0.0
European flounder (g/week)	42	31	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4.2	0.0	0.0	0.0	42	35	0.0	0.0
Salmon (g/week)	85	65	0.0	0.0	118	88	0.0	0.0	0.0	0.0	0.0	0.0	16	9.5	0.0	0.0	1.3	0.0	0.0	0.0
Herring (g/week)	3.5	0.0	16	70	2.9	0.0	46	71	203	134	0.0	57	18	24	31	89	1.5	4.8	16	42
Mackerel (g/week)	17	6.3	30	0.0	23	8.6	4.4	0.0	45	0.0	0.0	0.0	26	34	39	0.0	2.2	6.4	17	0.0
Trout (g/week)	28	11	0.0	0.0	38	14	14	0.0	0.0	0.0	0.0	0.0	14	9.0	1.0	0.0	1.1	0.0	0.0	0.0
Greenland halibut (g/week)	2.1	0.0	0.0	0.0	2.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	8.1	0.0	0.0	0.0	0.66	0.0	0.0	0.0




Fig. 3. Optimal trade-off curve between deviation from preference and change in cost (modelled cost minus observed cost) (a) and modelled fish intake recommendation for different parameter values λ (b) for a woman consuming all nine fish species (individual 1). Original to this manuscript. DKK, Danish krone.

Personal preference

When only the deviation from individual preference is minimised ($\lambda = 0$ in equation 1), the average values for the modelled recommendations of the study population were 82 g of lean fish and 134 g of fatty fish per week for women and 104 g of lean fish and 156 g of fatty fish per week for men (Table 6). The suggested changes in fish intake (modelled recommendation minus observed intake) with $\lambda = 0$ are visualised with an empirical cumulative distribution function in Fig. 1. From this figure, we can see that the species that should be recommended to be increased the most are mackerel, herring, salmon and trout, in that given order. A total of 48 % of the 3016 individuals (741 women and 703 men) already fulfilled their model constraints (for EPA + DHA, vitamin D, methyl mercury and dioxins + dl-PCB) with their observed consumption, that is, had current consumption within her/his feasible region. These individuals should be recommended to retain their fish consumption pattern. Half of the study population, 50 % (788 women and 723 men), should be suggested to increase their fish intake with up to 173 g of fish per week in order to meet the lower constraints of the nutrients. A small fraction of the study population, the remaining 2 % (twenty-three women and thirty-eight men), should be suggested to decrease their fish intake due to too high

exposure to contaminants with their current consumption. There were twenty-two individuals (thirteen women and nine men) who had too high of a background exposure to dioxins + dl-PCB to obtain a feasible solution. For these participants, no personalised fish intake recommendation could be generated. Instead, these individuals should be suggested to modify their background exposure, typically by consuming less animal products⁽²²⁾. From now on, we exclude these individuals from the discussion of the results. The cost for the average modelled fish intake recommendations are 39 (14 + 25) and 45 (17 + 28) DKK per week for women and men, respectively (see Table 6).

Cost

When only cost is minimised (the alternative cost objective function (equation 4) is used), the cost of the average modelled recommendations are 7.4 (0.0 + 7.4) and 7.1 (0.0 + 7.1) DKK per week for women and men, respectively (Table 6). When preference is not considered, but the aim is to recommend the cheapest fish consumption that fulfils the nutrient requirements, 99 % of the study population should be recommended to only eat various amounts of herring. The exceptions are two women who should be recommended to consume mackerel and trout, and twelve women and twenty-two men who should be




Fig. 4. Optimal trade-off curve between deviation from preference and change in cost (modelled cost minus observed cost) (a) and modelled fish intake recommendation for different parameter values λ (b) for a woman only consuming fatty fish species (individual 2). Original to this manuscript. DKK, Danish krone.




Fig. 5. Optimal trade-off curve between deviation from preference and change in cost (modelled cost minus observed cost) (a) and modelled fish intake recommendation for different parameter values λ (b) for a man with the highest fish consumption in the study population (individual 3). Original to this manuscript. DKK, Danish krone.

recommended to not consume fish at all, since their background exposure to the nutrients and contaminants fulfilled the constraints of the model. The average values for the modelled recommendations of the study population were 82 g of fatty fish (81.8 g herring, 0.034 g mackerel and 0.089 g trout) per week for women and 79 g of herring per week for men. See the empirical distribution function for the suggested changes (modelled recommendation minus observed intake) when only cost of fish intake is minimised in Fig. 2.

Personal preference and cost

The optimal trade-off curve for an individual can be visualised by plotting, with different parameter values for λ , the minimised

change in cost (modelled cost minus observed cost) $Cx - Cx_{obs}$ (DKK/week) on the x-axis and the corresponding minimised deviation from her/his preference $x - x_{obs2}$ (g/week) on the y-axis. For all individuals in our study population, there is a trade-off between deviation from preference and cost. This is illustrated for five representative individuals with different personal preference. The aim when choosing the individuals was to illustrate the variation in trade-off curves. Two individuals who fulfilled their nutrient and contaminant constraints were chosen, that is, their observed intakes were within the feasible region of the model. Three who did not meet their model constraints were chosen. Other than these criteria, the individuals were arbitrarily chosen. Below is a list of these five individuals (see their recorded fish intakes in Table 2).




Fig. 6. Optimal trade-off curve between deviation from preference and change in cost (modelled cost minus observed cost) (a) and modelled fish intake recommendation for different parameter values λ (b) for a woman not consuming fish (individual 4). Original to this manuscript. DKK, Danish krone.




Fig. 7. Optimal trade-off curve between deviation from preference and change in cost (modelled cost minus observed cost) (a) and modelled fish intake recommendation for different parameter values λ (b) for a man only consuming lean fish species (individual 5). Original to this manuscript. DKK, Danish krone.

Observed intake within feasible region:

- (1) a woman consuming all fish species
- (2) a woman only consuming fatty fish.

Observed intake outside feasible region:

- (1) a man with the highest fish consumption in the study population
- (2) a woman not consuming fish
- (3) a man only consuming lean fish.

In Figs. 3–7, the optimal trade-off curves for the five individuals along with the modelled recommended fish intake (g/week) for different parameter values λ are shown. In Table 7, the modelled recommendations with a few different λ are given. As shown in Figs. 3(b)–7(b) and Table 7, with large enough λ , all

individuals should be recommended to consume only herring. The rightmost point on any trade-off curve equals $\lambda = 0$, which corresponds to the recommended fish intake that minimise the deviation from the individual's preference (Figs. 3(a)–7(a)). With this parameter value, the cost of the fish intake is not considered. When $\lambda > 0$, the cost is included in the objective function. When λ is increased (moving left on the trade-off curve), the cost of fish consumption is decreased. Simultaneously, the deviation from the individual's preference is increased. The deviation from the preference is always a positive number, but the change in cost can be both positive and negative. When the cost of the modelled fish intake recommendation is larger than the cost of the observed intake, the change is positive; otherwise, it is negative. The leftmost point on a trade-off curve corresponds to the parameter value λ that minimises the cost for the individual. As mentioned, when λ becomes

large, the objective function of the model (equation 1) approaches the alternative objective function (equation 4) that only minimises cost.

For individual 1 consuming all species of fish and individual 2 consuming only fatty fish (Figs. 3 and 4), both the change in cost and the deviation from preference are 0 when the cost part of objective function is removed and only the deviation from personal preference is minimised, that is, $\lambda = 0$. This is because the observed intake of these individuals already fulfils the nutrient and contaminant constraints, and they would therefore be recommended to retain their fish consumption pattern when cost is not considered. As mentioned, this is the case for 43 % of the study population (Fig. 1(a)). An increased λ (>0) corresponds to a decreased (change in) cost (which, as mentioned, applies to all individuals) and a larger deviation from the preference. Individual 3 (Fig. 5) has an observed intake that is larger than the constraints allow. Hence, the change in cost is negative and the deviation from the preference is larger than 0 for all parameter values λ . This individual should be suggested to decrease fish intake for any λ , and the larger the λ , the larger the decrease in both fish intake and cost. For individual 4 who does not consume any fish (Fig. 6), the change in cost is positive and the deviation from the preference is larger than 0 for all parameter values λ . This individual should be suggested to increase fish intake for all λ , and the larger the λ , the larger the increase in fish intake and decrease in cost. When only minimising the deviation from preference ($\lambda = 0$), this non-fish consuming individual will be recommended the combination of species that gives the minimum total fish intake (g/week). Lastly, for individual 5 only consuming lean fish (Fig. 7), the change in cost is positive and the deviation from the preference is larger than 0 when $\lambda = 0$. Between $\lambda = 2.2$ and $\lambda = 2.4$, the change in cost goes from positive to negative. Hence, for $\lambda < 2.2$, the individual would save money by following the generated recommendation as compared with consuming the observed intake; and for $\lambda > 2.4$, the modelled recommendation would be the less expensive option. A recommendation that costs as much as the observed consumption can be given to this individual.

Discussion

To our knowledge, this is the first intake optimisation study that simultaneously minimises deviation from preference, which is defined as observed intake, and food cost. This allows for analysing the optimal trade-off curve between these two factors that may affect consumer choice. Fish consumption in Denmark was used for demonstrating the method and the model. As food choice is complex, it is not known with certainty to what extent and in which direction personal preference and cost affect individual's food choices. With this complexity, the model developed enables analysing the combination of these two important food choice determinants on individual level. We concluded that an optimal trade-off curve between deviation from fish intake preference and cost of fish could be generated for each individual in our study population of 3016 adult Danes, aside from twenty-two individuals who had a too high background exposure to the contaminants we included in the study (methyl

mercury and dioxins + dl-PCB). In order to obtain a trade-off curve, these individuals would need to modify intake other than fish.

When only minimising the deviation from preference for the individuals in the study population, 50 % should be suggested to increase their fish intake. This was also concluded in our previous fish intake optimisation study⁽²²⁾. The species that should be recommended to be increased the most are mackerel, herring, salmon and trout, in that given order. These are all fatty fish species, which are highest in EPA + DHA and vitamin D. The costs for the 'average recommendations' are 39 and 45 DKK per week for women and men, respectively. As compared with the cost when consuming 150 g of lean fish and 200 g of fatty fish, which is 61 DKK per week, the modelled recommendations that minimise the deviation from preference would be 20 and 14 DKK less expensive per week for the average woman and man, respectively.

When only minimising the cost, the vast majority of the study population (99 %) should be recommended to only consume herring. This is due to the fact that herring is the least expensive fish, while its nutrient compositions and contaminant concentrations allow for a feasible solution for all individuals except two women, who should be recommended to include mackerel and trout as well. However, it is not realistic to recommend almost a whole population to consume only herring. First, the recommendations generated by minimising cost do not account for preferences that probably make them unrealistic in the first place. As the constant λ determines the relative importance of the preferred diversity of an individual, in this 'extreme case' when $\lambda \rightarrow \infty$, a potential preference for diversity in fish species consumed is not considered separately in the model. Second, overfishing would most likely be a problem if the proportion of individuals that should be recommended to only consume herring as derived in this study is extrapolated to the general population. Third, the price of herring would eventually increase if the demand suddenly increased dramatically. Fourth, the recommendation would likely lead to discontent among the fish industry.

We argue that it is appropriate to minimise the deviation from the preference and cost simultaneously when generating personalised fish intake recommendations, since both factors may affect consumer choice and the aim is to obtain as realistic and achievable fish intake recommendations as possible. For example, salmon and trout, which are two of the species that should be increased the most when only minimising deviation from preference, are relatively expensive fish species. Hence, recommending these species to some individuals may result in lower adherence as compared with a recommendation that deviates more from their current preference, but to a lower cost. However, the relative importance of the cost is hard to draw conclusions on; we cannot define the best value for λ as it is a specific and subjective individual choice. Data on correlations between cost and consumer choice are needed in order to draw conclusions on the optimal trade-off between cost and personal preference.

Communication of the personalised recommendations generated with this model is a future research topic. A prospective utilisation of the model could be to apply it in a user-friendly

software tool (e.g. an application (app) or a computer program) that the population could access. This would allow individuals to generate personalised recommendations themselves and analyse her/his personal trade-off curve between deviation from preference and cost. If an individual is not interested in, for example, minimising the cost, the parameter λ can be set to 0. The model can be modified to not allow undesired species in the recommendation, which allows individuals to personalise their trade-off curves further. Data on preference regarding undesired fish species would have benefitted the analysis in this study, but no such data were available. However, if applied in an app, such data could easily be inserted by the user.

Previous studies have applied LP to model diets that deviate as little as possible from individual observed diets^(19,20). In general, QP may make small changes to all elements of the optimisation variable, whereas LP typically makes large changes in a few elements and leaves the others unchanged. What is perceived as realistic changes in intake is probably individual specific. Some individuals may prefer to make several small changes, and others may prefer fewer larger changes. We had no data on this preference, but we made the assumption that QP modelling generates more acceptable fish intake recommendations for our study population. This assumption is based on the fact that a recommendation would be ineffective if an individual was suggested to only consume a lot (more) of a species that she/he did not want to consume (more of). We argue that several small changes may be less likely to be deemed unacceptable, especially when new species are recommended to be introduced. As mentioned, it would be best if data regarding individual undesired species were available. A future research topic could be to perform a sensitivity analysis on the difference between QP and LP when minimising the optimal trade-off curve between personal fish intake preference and cost.

Recently, more and more countries have stated to incorporate sustainability considerations into their FBDG, for example, the environmental dimension⁽³⁸⁾. Fish consumption raises a significant trade-off between health (e.g. nutritional goals) and environmental impact. Therefore, consumption of fish from sustainable stocks is increasingly incorporated in recommendations. This study on fish intake assessed the nutrition/health, the economic/affordability and the cultural acceptability dimensions of diet sustainability. Several diet optimisation studies have analysed the environmental dimension when generating sustainable diets^(18,39–42), by including one or more environmental metrics such as greenhouse gas emissions, land use and fossil use. Our model could in a future study, in line with the developments of the FBDG and the focus on sustainable fish species, be modified to include some environmental metric.

The prices of the fish species used in this study were collected in the month of April. However, there may well be a seasonal variation in the prices. Also, average values were used, which may not reflect the behaviour of the population; some individuals may consequently prefer to buy low-price products, whereas others may choose the more expensive options. The nutrient compositions and contaminant concentrations for different fish species were described by average values in the model. This can be argued to reflect a realistic long-term consumption and exposure. Furthermore, there may be variation in nutrient

compositions and contaminant concentrations depending on region and season of capture, whether the fish is wild-caught or farmed-raised, etc.⁽²⁷⁾, but the intake data did not provide such information, and the model could not be personalised considering this.

Only EPA + DHA, vitamin D, methyl mercury, dioxins and dl-PCB were included in the modelling in this study. To make the results more precise, additional nutrients and/or contaminants present in fish could be included in the model. We considered it suitable to only include these nutrients and contaminants that present the major benefits and risks, since this paper is mainly methodological. Also, substitution with other foods was not considered in the model. It is likely that an individual would modify some other food intake if increasing or decreasing the intake of fish, which would change her/his background exposure⁽³⁾. The approach of this paper could be extended to optimise the intake of several foods or whole diets. With whole diet optimisation^(19,20,23), the substitution between foods is naturally covered in the model. The whole diet approach can also cover all dietary recommendations, and not only those associated with fish, as in this study. This, however, requires a lot of additional data, in terms of compositions, concentrations, prices, limit values, etc.

Furthermore, the results of this method should be applied in a real-life setting in order to validate their impact. In this paper, we assumed that the personalised dietary recommendations deviating as little as possible, to possibly at a lower cost, will have higher adherence than general FBDG, but it is not certain and we found no literature verifying this. Exploring this would require knowledge from other research fields.

Acknowledgements

We thank Sara M. Pires and Morten Poulsen for the valuable discussions and for critically reading the manuscript.

The preparation of this manuscript was funded through the Metrix project by the Ministry for Environment and Food in Denmark. The Ministry for Environment and Food in Denmark had no role in the design, analysis or writing of this article.

The authors' contributions are as follows: M. P. and M. J. N. formulated the research question, designed the study and drafted the manuscript; S. F. provided essential material; M. P. carried out the study and analysed the data; all the authors approved the final version of the manuscript.

None of the authors has any conflicts of interest to declare.

References

1. Brown KA, Timotijevic L, Barnett J, *et al.* (2011) A review of consumer awareness, understanding and use of food-based dietary guidelines. *Br J Nutr* **106**, 15–26.
2. Tetens I, Andersen LB, Astrup A, *et al.* (2013) *The Evidence-Base for the Danish Guidelines for Diet and Physical Activity*. Søborg, Denmark: National Food Institute, Technical University of Denmark.
3. Thomsen ST, Pires SM, Devleeschauwer B, *et al.* (2018) Investigating the risk-benefit balance of substituting red and processed meat with fish in a Danish diet. *Food Chem Toxicol* **120**, 50–63.
4. Pedersen AN, Christensen T, Matthiessen J, *et al.* *Unpublished Data from the Danish National Survey of Diet and Physical*

- Activity (DANSDA) April 2011 – August/September 2013.* Søborg, Denmark: National Food Institute, Technical University of Denmark.
5. Pedersen AN, Christensen T, Matthiessen J, *et al.* (2015) *Dietary Habits in Denmark 2011–2013*. Søborg, Denmark: National Food Institute, Technical University of Denmark.
 6. Leng G, Adan RAH, Belot M, *et al.* (2017) The determinants of food choice. *Proc Nutr Soc* **76**, 316–327.
 7. Kö EP (2007) Diversity in the determinants of food choice: a psychological perspective. *Food Qual Prefer* **20**, 70–82.
 8. James WP, Nelson M, Ralph A, *et al.* (1997) Socioeconomic determinants of health: the contribution of nutrition to inequalities in health. *BMJ* **314**, 1545–1549.
 9. Darmon N & Drewnowski A (2015) Contribution of food prices and diet cost to socioeconomic disparities in diet quality and health: a systematic review and analysis. *Nutr Rev* **73**, 643–660.
 10. Lo Y-T, Chang Y-H, Drph M-SL, *et al.* (2009) Health and nutrition economics: diet costs are associated with diet quality. *Asia Pac J Clin Nutr* **18**, 598–604.
 11. Aggarwal A, Monsivais P, Cook AJ, *et al.* (2011) Does diet cost mediate the relation between socioeconomic position and diet quality? *Eur J Clin Nutr* **65**, 1059–1066.
 12. Turrell G, Hewitt B, Patterson C, *et al.* (2002) Socioeconomic differences in food purchasing behaviour and suggested implications for diet-related health promotion. *J Hum Nutr Diet* **15**, 355–364.
 13. Giskes K, Van Lenthe FJ, Brug J, *et al.* (2007) Socioeconomic inequalities in food purchasing: the contribution of respondent-perceived and actual (objectively measured) price and availability of foods. *Prev Med (Baltim)* **45**, 41–48.
 14. King JC (2007) An evidence-based approach for establishing dietary guidelines. *J Nutr* **137**, 480–483.
 15. Brug J, Campbell M & van Assema P (1999) The application and impact of computer-generated personalized nutrition education: a review of the literature. *Patient Educ Couns* **36**, 145–156.
 16. Brug J, Oenema A & Campbell M (2003) Past, present, and future of computer-tailored nutrition education. *Am J Clin Nutr* **77**, 1028S–1034S.
 17. Celis-Morales C, Livingstone KM, Marsaux CFM, *et al.* (2016) Effect of personalized nutrition on health-related behaviour change: evidence from the Food4me European randomized controlled trial. *Int J Epidemiol* **46**, 578–588.
 18. Gazan R, Brouzes CMC, Vieux F, *et al.* (2018) Mathematical optimization to explore tomorrow's sustainable diets: a narrative review. *Adv Nutr* **9**, 602–616.
 19. Maillot M, Vieux F, Amiot MJ, *et al.* (2010) Individual diet modeling translates nutrient recommendations into realistic and individual-specific food choices. *Am J Clin Nutr* **91**, 421–430.
 20. Maillot M, Vieux F, Ferguson E, *et al.* (2009) To meet nutrient recommendations, most French adults need to expand their habitual food repertoire. *J Nutr* **139**, 1721–1727.
 21. Persson M, Fagt S, Pires SM, *et al.* (2018) Use of mathematical optimization models to derive healthy and safe fish intake. *J Nutr* **148**, 275–284.
 22. Persson M, Fagt S & Nauta MJ (2018) Personalised fish intake recommendations: the effect of background exposure on optimisation. *Br J Nutr* **120**, 946–957.
 23. Barré T, Vieux F, Perignon M, *et al.* (2016) Reaching nutritional adequacy does not necessarily increase exposure to food contaminants: evidence from a whole-diet modeling approach. *J Nutr* **146**, 2149–2157.
 24. Barré T, Perignon M, Gazan R, *et al.* (2018) Integrating nutrient bioavailability and co-production links when identifying sustainable diets: how low should we reduce meat consumption? *PLOS ONE* **13**, e0191767.
 25. Sirot V, Leblanc J-C & Margaritis I (2012) A risk–benefit analysis approach to seafood intake to determine optimal consumption. *Br J Nutr* **107**, 1812–1822.
 26. Darmon N, Ferguson EL & Briend A (2002) A cost constraint alone has adverse effects on food selection and nutrient density: an analysis of human diets by linear programming. *J Nutr* **132**, 3764–3771.
 27. Norwegian Scientific Committee for Food Safety (VKM) (2014) *Benefit-risk Assessment of Fish and Fish Products in the Norwegian Diet – An Update*. Oslo, Norway: Norwegian Scientific Committee for Food Safety.
 28. National Food Institute at Technical University of Denmark (DTU) (2017) *Frida Version 2 Udgave 2017-06-06*. Søborg, Denmark: National Food Institute, Technical University of Denmark.
 29. National Food Institute at Technical University of Denmark (DTU) (2011) *Chemical Contaminants 2004–2011*. Søborg, Denmark: National Food Institute, Technical University of Denmark.
 30. National Food Institute at Technical University of Denmark (DTU) (2013) *Chemical Contaminants 2012–2013*. Søborg, Denmark: National Food Institute, Technical University of Denmark.
 31. EFSA Panel on Contaminants in the Food Chain (CONTAM) (2012) Scientific opinion on the risk for public health related to the presence of mercury and methylmercury in food. *EFSA J* **10**, 2985.
 32. Andersen R, Brot C, Jakobsen J, *et al.* (2013) Seasonal changes in vitamin D status among Danish adolescent girls and elderly women: the influence of sun exposure and vitamin D intake. *Eur J Clin Nutr* **673**, 270–274.
 33. Hansen L, Tjønneland A, Køster B, *et al.* (2016) Sun exposure guidelines and serum vitamin D status in Denmark: the StatusD study. *Nutrients* **8**, E266.
 34. Hong Y-S, Kim Y-M & Lee K-E (2012) Methylmercury exposure and health effects. *J Prev Med Public Health* **45**, 353–363.
 35. European Commission (2000) *Dioxin Contamination of Feeding Stuffs and their Contribution to the Contamination of Food of Animal Origin*. Brussels, Belgium: European Commission.
 36. Grant M & Boyd S (2013) CVX: Matlab software for disciplined convex programming, version 2.0 beta.
 37. Grant MC & Boyd SP (2008) Graph implementations for nonsmooth convex programs. In *Recent Advances in Learning and Control. Lecture Notes in Control and Information Sciences*, vol. 371, pp. 95–110 [VD Blondel, SP Boyd and H Kimura, editors]. London: Springer.
 38. Gonzalez Fischer C & Garnett T (2018) Plates, pyramids and planets developments in national healthy and sustainable dietary guidelines: a state of play assessment. <http://www.fao.org/3/a-i5640e.pdf>
 39. Perignon M, Masset G, Ferrari G, *et al.* (2016) How low can dietary greenhouse gas emissions be reduced without impairing nutritional adequacy, affordability and acceptability of the diet? A modelling study to guide sustainable food choices. *Public Health Nutr* **19**, 2662–2674.
 40. Tyszler M, Kramer G & Blonk H (2016) Just eating healthier is not enough: studying the environmental impact of different diet scenarios for Dutch women (31–50 years old) by linear programming. *Int J Life Cycle Assess* **21**, 701–709.
 41. Horgan GW, Perrin A, Whybrow S, *et al.* (2016) Achieving dietary recommendations and reducing greenhouse gas emissions: modelling diets to minimise the change from current intakes. *Int J Behav Nutr Phys Act* **13**, 46.
 42. Kramer GF, Tyszler M, Veer PV, *et al.* (2017) Decreasing the overall environmental impact of the Dutch diet: how to find healthy and sustainable diets with limited changes. *Public Health Nutr* **20**, 1699–1709.



43. EFSA Panel on Dietetic Products, Nutrition, and Allergies (NDA) (2010) Scientific opinion on dietary reference values for fats, including saturated fatty acids, polyunsaturated fatty acids, monounsaturated fatty acids, *trans* fatty acids, and cholesterol. *EFSA J* **8**, 1461.
44. Nordic Council of Ministers (2014) *Nordic Nutrition Recommendations 2012 Integrating Nutrition and Physical Activity*. Copenhagen, Denmark: Nordic Council of Ministers.
45. EU Scientific Committee on Food (2001) *Fact SHEET on Dioxin in Feed and Food*. Brussels, Belgium: European Commission.