



The interplanetary dust population as measured by the MAG investigation on the Juno mission

Jørgensen, John; Benn, Mathias; Connerney, Jack; Andersen, Anja; Jørgensen, Peter; Denver, Troelz; Bolton, Scott; Levin, Steve

Published in:
Geophysical Research Letters

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Jørgensen, J., Benn, M., Connerney, J., Andersen, A., Jørgensen, P., Denver, T., Bolton, S., & Levin, S. (2019). The interplanetary dust population as measured by the MAG investigation on the Juno mission. *Geophysical Research Letters*, 21, Article EGU2019-7448.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



The interplanetary dust population as measured by the MAG investigation on the Juno mission

John Jørgensen (1), Mathias Benn (1), Jack Connerney (2,3), Anja Andersen (4), Peter Jørgensen (1), Troelz Denver (1), Scott Bolton (5), and Steve Levin (6)

(1) Technical University of Denmark, DTU-Space, Space, Denmark (jlj@space.dtu.dk), (2) Space Research Corporation, Annapolis, Maryland, USA, (3) Goddard SpaceFlight Center, Maryland, USA, (4) NBI, Copenhagen University, Denmark, (5) Southwest Research Institute, San Antonio, Tx, USA, (6) Jet Propulsion Laboratory, Caltech, Pasadena, CA, USA

NASA's Jupiter mission Juno was launched from Earth August 5, 2011, and arrived at Jupiter nearly 5 years later, on July 5, 2016. During cruise from Earth to Jupiter, the solar powered Juno spacecraft followed a trajectory which took advantage of a gravity assist maneuver, passing close by Earth October 9, 2013. Juno's ~600 W of electrical power is provided by 60m² of solar panels arranged on three solar panel wings radiating from the main body of the spinning spacecraft. One of these wings accommodates two MAG investigation sensor suites, separated by two meters on a non-magnetic boom at the outer extremity. Each sensor suite consists of a fluxgate vector magnetometer and two μ ASC star cameras. The primary purpose of the μ ASC cameras is to provide accurate attitude for the MAG sensors, but they may also be used for general low-light imaging, asteroid detection and tracking, and energetic particle detection. Here we describe how the μ ASC is used to detect hypervelocity dust impacts on the spacecraft, characterizing the flux of relatively large ($> \mu\text{m}$) interplanetary dust (IPD) particles that have heretofore escaped direct detection by dust instrumentation flown for IDP detection. Direct optical detection of such IDPs is prohibited by their relative velocity (order 10's km/s), but these particles impact the spacecraft and generate secondary ejecta which are readily observed and tracked by the μ ASC. Thus we employ Juno's 60 m² of solar array as a very large aperture dust detector, with orders of magnitude more collecting area than any IDP instrument yet flown. This copious collecting area provides us with sufficient fluxes to characterize the distribution of interplanetary dust particles that are responsible for the Zodiacal light. We present this novel IPD detection technique, its detection sensitivity, and discuss the distribution of IDP particles between Earth and Jupiter.