



Synthesis of Arabinoxylan Oligosaccharides by Pre-Activation-Based Iterative Glycosylations

Underlin, Emilie Nørmølle; Böhm, Maximilian Felix; Madsen, Robert

Published in:
Journal of organic chemistry

Link to article, DOI:
[10.1021/acs.joc.9b02529](https://doi.org/10.1021/acs.joc.9b02529)

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Underlin, E. N., Böhm, M. F., & Madsen, R. (2019). Synthesis of Arabinoxylan Oligosaccharides by Pre-Activation-Based Iterative Glycosylations. *Journal of organic chemistry*, 84(24), 16036-16054.
<https://doi.org/10.1021/acs.joc.9b02529>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Synthesis of Arabinoxylan Oligosaccharides by Pre-Activation-Based Iterative Glycosylations

Emilie N. Underlin, Maximilian Böhm, and Robert Madsen

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.9b02529 • Publication Date (Web): 25 Nov 2019

Downloaded from pubs.acs.org on December 3, 2019

Just Accepted


“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Synthesis of Arabinoxylan Oligosaccharides by Pre-Activation-Based Iterative Glycosylations

Emilie N. Underlin, Maximilian Böhm, and Robert Madsen*

Department of Chemistry, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark

E-mail: rm@kemi.dtu.dk



Abstract

A concise synthetic strategy has been developed for assembling densely substituted arabinoxylan oligosaccharides, which are valuable substrates for characterizing hemicellulose-degrading enzymes. The xylan backbone has been prepared by an iterative pre-activation-based glycosylation approach with phenyl thioglycosides. The pre-activation has been performed with in situ generated *p*-nitrobenzenesulfonyl triflate prior to addition of the acceptor. The glycosylation temperature was shown to have an important impact on the yield of the coupling. The arabinose substituents have been introduced in one high-yielding glycosylation with a *N*-phenyl trifluoroacetimidate donor. The strategy has been successfully employed for the

1
2
3 synthesis of three heptasaccharides in seven steps and overall yields of 24 – 36% from the
4
5 corresponding monosaccharide building blocks.
6
7

11 INTRODUCTION

12
13
14
15 The hemicellulose polysaccharides comprise 15 – 35% by weight of the plant cell
16
17 wall which is the main source of lignocellulosic biomass.¹ The largest group of
18
19 polysaccharides in hemicellulose are the xylans which are characterized by a common
20
21 backbone of $\beta(1\rightarrow4)$ -linked xylopyranosides.¹ The xylan structures are highly complex
22
23 and heterogeneous due to the differences in the substitution patterns. One of the most
24
25 abundant xylan subclasses are the arabinoxylans where some of the xylose units are
26
27 substituted with L-arabinofuranosyl residues through either single substitution by
28
29 $\alpha(1\rightarrow2)$ or $\alpha(1\rightarrow3)$ linkages or double substitution by both linkages.^{1,2}
30
31
32
33
34
35
36
37
38
39
40
41
42

43 The arabinoxylans are dietary fibers that are not hydrolyzed by the digestive
44
45 enzymes.³ Instead they are fermented by microbes in the gut and in this way influences
46
47 the composition and abundance of the colonic microbiota.³ The result is a prebiotic
48
49 effect where the health of the host is benefitted. Arabinoxylans have been shown to
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 strengthen the immune system and to lower the risk for obesity, type 2 diabetes, colon
5
6
7 cancer and cardiovascular diseases.³ The physiological functions, however, are
8
9
10 strongly connected to the structure of the arabinoxylans, i.e. the length of the xylan
11
12
13 backbone and the distribution of the arabinose units.³ Besides food applications
14
15
16 arabinoxylans are also used as a renewable source to prepare biodegradable materials
17
18
19
20
21 such as films and hydrogels.^{1,4}
22
23

24 Arabinoxylans are cleaved by glycosyl hydrolases (GH) to form smaller saccharides.
25
26
27
28 The hydrolysis of the xylose linkages can be achieved with GH10 and GH11 xylanases,
29
30
31 which prefer several unsubstituted xylose units, and with GH5 xylanases where an
32
33
34 arabinose substituent is allowed close to the cleavage site.⁵ Hydrolysis of the arabinose
35
36
37 residues can be performed with GH43, GH51 and GH62 arabinofuranosidases where
38
39
40
41 some of the enzymes exclusively cleave arabinoses linked to the 2 or the 3 position of
42
43
44 single substituted xylose residues, whereas others are able to remove the pentose from
45
46
47 disubstituted xylose moieties.^{5,6} All the enzyme classes, however, suffer from poor
48
49
50
51
52 access in the densely substituted regions of arabinoxylans and the difficulties are
53
54
55
56
57
58
59
60

1
2
3
4 further complicated by the lack of well-defined substrates with multiple arabinose
5
6
7 substituents.
8
9

10 Arabinoxylan-degrading enzymes are usually investigated with rather heterogeneous
11
12 polysaccharides isolated from natural sources by suitable pretreatment procedures. A
13
14 few saccharides containing 2 – 4 xylose units and 1 – 2 arabinose residues can be
15
16
17 isolated from enzymatic degradation of arabinoxylans,⁷ but these simple substrates do
18
19
20
21 not resemble the more densely substituted regions. Thus, well-defined xylans with
22
23
24 several arabinose substituents would be valuable tools for mapping the active site of
25
26
27
28 enzymes implicated in arabinoxylan degradation and to understand the interplay
29
30
31 between different enzymes in order to achieve the most efficient deconstruction.
32
33
34
35
36
37

38 Chemical synthesis offers the possibility to prepare pure and well-defined
39
40
41 oligosaccharides, but the arabinoxylans have so far received relatively little attention.
42
43
44 Xylobiose has been glycosylated with one or two equiv. of an arabinofuranosyl donor to
45
46
47
48 afford tri- and tetrasaccharides.⁸ Recently, arabinoxylans with up to 6 xylose residues
49
50
51 and 1 – 2 arabinose units were prepared by solid-phase synthesis and used for
52
53
54
55
56
57
58
59
60

1
2
3
4 determining the epitopes for monoclonal antibodies and the substrate specificity for
5
6
7 xylanases and arabinofuranosidases.⁹ In one case, two arabinoses were glycosylated
8
9
10 onto the 2 and the 3 position of the same xylose residue although a very low yield was
11
12
13 obtained illustrating the challenge of producing more densely substituted
14
15
16
17 arabinoxylans. Linear xylans, on the other hand, where no additional substituents are
18
19
20
21 present, are more readily assembled by either stepwise or blockwise approaches¹⁰ and
22
23
24 so far linear xylans up to decaxylans have been prepared,¹¹ which includes an
25
26
27
28 approach by cleaving the hydroxymethyl group of glucans¹² and the assembly of *S*-
29
30
31 linked oligoxylans.¹³
32
33
34


35 Herein, we report a novel pre-activation-based synthetic strategy for assembling
36
37
38 arabinoxylans, which has led to the preparation of heptasaccharides **1** – **3** with three
39
40
41 arabinose substituents (Scheme 1). A tetraxylan is chosen as the backbone since this
42
43
44 constitutes a sufficient length to give the necessary enzyme activities.^{5,9a} The three
45
46
47 arabinose substituents are connected to either a xylotriose unit (as in **1**) or a xylobiose
48
49
50 moiety (as in **2** and **3**) in order to resemble the densely substituted region. The
51
52
53
54
55
56
57
58
59
60

preparation of **1** – **3** extends our work on the synthesis of plant cell wall components

where we have previously assembled oligomers of rhamnogalacturonan I and

homogalacturonan by using *n*-pentenyl glycosides as glycosyl donors.¹⁴


Scheme 1. Structures of Arabinoxylans **1** – **3** and Retrosynthetic Strategy



RESULTS AND DISCUSSION

1
2
3 The recent solid-phase synthesis of arabinoxylans was achieved by a stepwise approach
4 where one xylose or arabinose unit was glycosylated at a time onto the growing oligosaccharide
5 chain followed by cleavage from the resin and global deprotection.⁹ Up to ten equiv. of a
6 xylopyranosyl phosphate (prepared from the corresponding tolyl thioglycoside) or an ethyl 1-
7 thioarabinofuranoside was used as the donor for each glycosylation reaction. We opted for a
8 solution-phase strategy involving fewer steps where all the arabinose units are introduced in one
9 coupling reaction (Scheme 1). The linear xylan backbone will be prepared by a pre-activation-
10 based synthesis¹⁵ from the appropriate phenyl 1-thioxylopyranosides where deprotection steps
11 will not be necessary after each glycosylation reaction.¹⁶ The pre-activation-based thioglycoside
12 protocol has previously been used for coupling of a variety of monosaccharides^{15,17} including
13 arabinofuranosides,¹⁸ but it has so far not been applied to the coupling of xylose units. The
14 benzoyl group will serve as a permanent protecting group for the xylose residues since it allows
15 for neighboring group participation from the 2 position to afford the β -linkage in the
16 glycosylations. The arabinose units will also be protected with benzoyl groups and the
17 corresponding *N*-phenyl trifluoroacetimidate **4** will serve as the donor since it has previously
18 given high yields in the coupling with different secondary alcohols.¹⁹


19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 **Scheme 2. Synthesis of Building Blocks 5 and 6**



Several phenyl 1-thioxylopyranoside building blocks were required for the xylan assembly by the pre-activation-based approach. This involves fully benzoylated glycoside **5**, 2,3-di-*O*-benzoylated glycoside **6** and two glycosides with different protecting groups at position 2 and 3. The building blocks were all prepared from phenyl 1-thio-β-D-xylopyranoside (**7**), which is available from xylose in three simple steps involving peracetylation in the presence of 4-(*N,N*-dimethylamino)pyridine (DMAP), thiolation and deacetylation²⁰ with no purification of the intermediates (Scheme 2). Perbenzoylation of **7** then afforded tribenzoate **5** while the preparation of dibenzoate **6** required selective transformations of the hydroxy groups. In xylopyranosides the three hydroxy groups are all equatorial and similar in reactivity making regioselective reactions a particular challenge.²¹ One of the most effective procedures is to block the 2 and the 3 position with an isopropylidene group, which affords only minor amounts of the corresponding 3,4-acetal.²² Thus, treatment of triol **7** with 2-methoxyprop-1-ene and camphorsulfonic acid (CSA) led to the acetonide at position 2 and 3 as the major product.

Subsequently, a *para*-methoxybenzyl group (PMB) was installed at position 4 and the acetonide was removed under acidic conditions to provide diol **8**.²³ The hydroxy groups were benzoylated and excess benzoyl chloride removed with 3-(dimethylamino)-1-propylamine (DMAPA).²⁴ Without further purification the crude product was subjected to PMB deprotection with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) to afford dibenzoate **6**.

Scheme 3. Synthesis of Building Blocks **8** and **10**




The two remaining building blocks required a different protecting group to be installed at position 3 as well as at position 2 and 3 since these sites will be linked to the arabinose units. The temporary protecting group should be an ester to ensure neighboring group participation in the glycosylation, but should at the same time be removable in the presence of the benzoates. A chloroacetate was first considered and installed selectively at position 3, but the following glycosylation with **5** under the pre-activation protocol only furnished a moderate yield of the disaccharide. This was attributed to the use of a silver salt in the glycosylation and a levulinyl (Lev) group was therefore chosen instead. *N,N'*-Dicyclohexylcarbodiimide (DCC) coupling of diol **8** with levulinic acid followed by direct treatment with DDQ furnished dilevulinate **9**

1
2
3 (Scheme 3). For the selective introduction of the temporary protecting group, diol **8** was first
4 subjected to regioselective benzylation at position 2. With benzoyl chloride in pyridine the
5 esterification was slow and actually produced the dibenzoate as the main product. However,
6 when diol **8** was submitted to 1.3 equiv. of benzoyl chloride in a mixture of CH₂Cl₂ and 1 M
7 NaOH under phase transfer catalysis,²⁵ 2-benzoate **10** was obtained in 68% yield with only
8 minor amounts of the corresponding 3-benzoate as a byproduct. Subsequent esterification with
9 levulinic acid and deprotection of the PMB group gave the desired 3-levulinate **11**.

19 The stage was now set for assembling the xylan backbone by the pre-activation-based
20 protocol. The original procedure with tolylthio glycosides called for activation of the donor with
21 *p*-toluenesulfonyl chloride and silver triflate at a temperature of -60 °C to generate the reactive
22 electrophile.¹⁶ The acceptor was then added followed by stirring for 15 min and warming to rt.
23 The procedure was later on modified with the formation of the reactive electrophile at -78 °C
24 followed by adding the acceptor and raising the temperature to rt.¹⁷ Using this latest protocol,
25 donor **5** was coupled to acceptor **6** with phenylsulfonyl chloride and silver triflate²⁶ to give
26 disaccharide **12** in 64% yield (Scheme 4, Method A). The coupling could also be carried out in a
27 one-pot iterative manner¹⁶ by adding more promoter and acceptor to furnish trisaccharide **13**
28 although the yield in this case was only 34% since disaccharide **12** was also formed in 59%
29 yield. Attempts to improve the glycosylations by using Me₂S₂/Tf₂O²⁷ or 1-benzenesulfonyl
30 piperidine/Tf₂O²⁸ as the promoter were unsuccessful and only led to a mixture of products.
31 However, when *p*-nitrobenzenesulfonyl chloride and silver triflate²⁹ was employed as the
32 promoter, the yield of **12** and **13** in the two reactions increased to 79 and 69%, respectively. *p*-
33 Nitrobenzenesulfonyl chloride is commercially available as opposed to phenylsulfonyl chloride
34 and *p*-toluenesulfonyl chloride which are prepared from the corresponding thiols with sulfuryl
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

chloride. In addition, *p*-nitrobenzenesulfonyl chloride is a solid making this reagent procedurally more convenient to deliver while the other two sulfonyl chlorides are liquids. The quality of commercial *p*-nitrobenzenesulfonyl chloride, however, varied and we obtained lower glycosylation yields when using light brown samples with a mp of ~43 °C. Attempts to recrystallize the compound failed, but sublimation yielded material with a light yellow color and a mp of 47 – 51 °C which could be stored at 5 °C for months. When *p*-nitrobenzenesulfonyl chloride purified by sublimation was used, the yields in the glycosylations were consistent. During the couplings, the disulfide byproduct from the activation (i.e. *p*NO₂PhSSPh) precipitated and did therefore not take part in any additional reactions.

Scheme 4. Synthesis of Saccharides 12 and 13



To further improve the glycosylation outcome the influence of the temperature was also investigated. These experiments were performed in a two-neck Schlenk flask that allowed for

1
2
3 measuring the internal temperature in the reaction mixture at any time during the glycosylation.
4
5 The activation of donor **5** was observed to be quick and goes to completion in less than 10 min
6
7 at $-78\text{ }^{\circ}\text{C}$ according to TLC analysis. This is in accordance with investigations on the activation
8
9 of similar thioglycosides under these conditions.³⁰ However, in the ensuing glycosylation
10
11 reaction after adding acceptor **6** the reaction temperature becomes important. If the temperature
12
13 is kept at $-78\text{ }^{\circ}\text{C}$ the glycosylation is sluggish and an intermolecular aglycon transfer³¹ becomes
14
15 a competing reaction leading to the regeneration of donor **5**.³² Under these conditions
16
17 disaccharide **12** was only obtained in 54% yield together with 10% of donor **5** although full
18
19 activation of the donor had been achieved. Thus, at very low temperature the hydroxy group and
20
21 the sulfur in acceptor **6** are both able to serve as a nucleophile in the reaction with the donor.
22
23 Further experiments revealed that the aglycon transfer was mainly observed when the
24
25 temperature was kept below $-60\text{ }^{\circ}\text{C}$. Therefore, three separate experiments were set up where
26
27 acceptor **6** was added at $-55\text{ }^{\circ}\text{C}$ (Figure 1). In the first experiment, the temperature was
28
29 immediately raised to rt over 1 h while in the two other experiments the temperature was first
30
31 maintained at $-55\text{ }^{\circ}\text{C}$ for 10 – 15 min before being raised. The first reaction gave 73% yield of
32
33 disaccharide **12** whereas 85 and 90% yield were obtained in the two other experiments (Figure 1
34
35 and Scheme 4, Method B). This is a notable difference and illustrates the significance of the
36
37 temperature in securing an optimal coupling reaction. If the temperature is too low the aglycon
38
39 transfer becomes a side reaction while if the temperature is too high the activated intermediate
40
41 of donor **5** (which is mostly the corresponding 1,2-benzoxonium ion)³³ presumably decomposes.
42
43 Thus, a coupling temperature of $-55\text{ }^{\circ}\text{C}$ was included in the protocol in the following
44
45 experiments. The reactions were monitored by TLC at $-55\text{ }^{\circ}\text{C}$ and the temperature was only
46
47 raised when the coupling had gone to completion.
48
49
50
51
52
53
54
55
56
57
58
59
60





Figure 1. Reaction Temperatures and Yields for Coupling of 5 and 6 to afford 12


Using this optimized procedure donor **5** was then coupled to Lev-protected acceptor **11** to afford disaccharide **14** in 86% yield (Scheme 5). Unfortunately, when the same coupling was performed in a one-pot fashion with the further addition of acceptor **6**, trisaccharide **15** was only obtained in 28% yield. The addition of 2,4,6-tri-*tert*-butylpyrimidine^{15,34} to mitigate the possible influence of the increasing amount of acid lowered the yield even further to 17%. Instead, large amounts of unreacted acceptor **6** were recovered in both cases. The low yield was not caused by the inability of Lev-protected thioglycosides to serve as donors since the coupling between disaccharide donor **14** and acceptor **6** gave rise to trisaccharide **15** in 69% yield. Possibly, the mediocre result is caused by the Lev group undergoing further transformations due to the longer reaction time in the one-pot protocol. Therefore, the xylan backbones in targets **1 – 3** will be assembled by an iterative glycosylation approach with pre-activation where the glycosylation products once isolated will be immediately used as donors for the next glycosylation step without any further modifications.

Scheme 5. Synthesis of Saccharides 14 and 15



Accordingly, to prepare arabinoxylan **1** disaccharide **14** was then coupled twice with Lev-protected acceptor **11** to afford trisaccharide **16** and tetrasaccharide **17** in 88 and 73% yields, respectively (Scheme 6). The three Lev groups in the 3 positions were removed with hydrazine to give triol **18** in 96% yield. Subsequent glycosylation with four equiv. of imidate donor **4** afforded protected arabinoxylan **19** in 90% yield. The arabinofuranose donor **4** was prepared in four straightforward steps from the parent pentose.^{35,19b} Thus, heptasaccharide **19** has been assembled from monosaccharides **4**, **5** and **11** in a total of five steps and 48% overall yield, which illustrates the advantage of the iterative pre-activation-based strategy.


Scheme 6. Synthesis of Heptasaccharide 19



The same sequence of reactions were then employed to prepare the protected forms of arabinoxylans **2** and **3**. To assemble the former, disaccharide **14** was used again and now coupled with 2,3-Lev-protected acceptor **9** to give trisaccharide **20** in 71% yield (Scheme 7). Further glycosylation with acceptor **6** produced 61% yield of tetrasaccharide **21**, which was subjected to Lev deprotection to afford triol **22**. The arabinoses were then installed with donor **4** to afford heptasaccharide **23** in 95% yield. Likewise, the route to arabinoxylan **3** began with the coupling between donor **5** and 2,3-Lev-protected acceptor **9** to furnish disaccharide **24** in 95% yield (Scheme 8). The following glycosylation with acceptor **11** produced trisaccharide **25** in 52% yield. The moderate yield in this coupling as well as between **20** and **6** in Scheme 7 may

1
2
3 indicate a lower stability of the activated species with a Lev group at the 2 position. Next,
4
5 trisaccharide **25** was reacted with acceptor **6** to give tetrasaccharide **26** in 75% yield, which was
6
7 followed by Lev deprotection to provide triol **27**. The final triple glycosylation with arabinose
8
9 donor **4** gave heptasaccharide **28** in 91% yield. Thus, protected heptasaccharides **23** and **28** have
10
11 each been prepared in five steps from monosaccharides **4**, **5**, **9** and **11** and with overall yields of
12
13 34 and 33%, respectively. In all cases, the three arabinoses were installed in very high yields
14
15 with imidate donor **4** proving the strength of this method even for forming densely substituted
16
17 substrates.
18
19
20
21
22
23


24 Scheme 7. Synthesis of Heptasaccharide **23**



53 The β -linkages of the xylose backbones were confirmed by measuring the J_{CH} coupling
54 constants for the anomeric carbon atoms, which were found to be between 155 and 165 Hz
55
56
57
58
59
60

where a coupling around 160 Hz is characteristic of a β -glycosidic bond.³⁶ The anomeric carbon atoms for the β -glycosidic linkages were located in the $\delta = 99.0 - 102.5$ ppm range, which is in accordance with earlier observations.¹¹ Xylopyranosides are more conformationally flexible than e.g. glucopyranosides²¹ and the $J_{H-1,H-2}$ coupling constants can therefore not always be used to determine their anomeric configuration. On the contrary, for the arabinofuranose substituents the $J_{H-1,H-2}$ coupling constants could be used to verify the α -linkages since small values of 0 – 2 Hz are characteristic for the α anomers.³⁷

Scheme 8. Synthesis of Heptasaccharide 28




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Deprotection of the three heptasaccharides is a two-step process since the thiophenyl moiety and the benzoates are removed under different conditions. In the first approach, the thiophenyl group was cleaved in the first step with NBS in a mixture of acetone and water.³⁸ This transformation proceeded uneventfully for all three heptasaccharides and gave rise to reducing sugars **29** – **31** in 92 – 93% yield (Scheme 9). The products were isolated as α/β mixtures with the α anomers as the major components according to NMR. The following Zemplén reaction to remove the benzoates³⁹ was carried out with approx. 20 equiv. of NaOMe in a 1:1 methanol – dichloromethane solution for up to 7 days to achieve complete deprotection of all benzoyl groups. Unfortunately, a mixture of saccharides of different length were obtained and it became apparent that the alkaline deprotection procedure had been accompanied by a so-called peeling reaction (endwise degradation).⁴⁰ In this process reducing saccharides are degraded with one monosaccharide at the time from the reducing end through aldose – ketose isomerizations and β -alkoxy eliminations.⁴⁰ To circumvent this side reaction, the order of the deprotections were reversed. First, the protected heptasaccharides were subjected to cleavage of the benzoyl groups, which proceeded cleanly to give compounds **32** – **34** in 77 – 91% yield. Then, attempts were made to remove the thiophenyl moiety under the same conditions as applied to the perbenzoylated substrates. However, these conditions now resulted in partial hydrolysis of the arabinofuranose residues, which is presumably due to the higher acid lability of these substituents in the absence of the benzoyl groups. Hence, 2,6-lutidine was added during the deprotections⁴¹ to ensure a pH of the reaction media around 6, which led to clean removal of the thiophenyl groups to afford the completely deprotected target structures **1** – **3** in 84 – 97% yield. Heptasaccharides **1** – **3** were fully characterized by 800 MHz NMR spectroscopy and although they were obtained as α/β mixtures, the differences in the ¹H and the ¹³C chemical shifts between

the α and the β anomers were only observed for the reducing end xylose unit and when an arabinose residue was attached to this moiety (as in **1**).

Scheme 9. Deprotection of Heptasaccharides **19**, **23** and **28**



In conclusion, three arabinoxylan heptasaccharides have been assembled in seven steps from monosaccharide building blocks in overall yields of 36 (**1**), 24 (**2**) and 27% (**3**). The key transformations are the pre-activation-based thioglycoside glycosylation to prepare the tetra- and pentasaccharide backbones and the triple glycosylation with imidate donor **4** to install the arabinose substituents. *p*-Nitrobenzenesulfenyl chloride was employed as a potent and convenient source of the sulfenium electrophile for the pre-activation of the phenyl thioglycoside donors. In the ensuing glycosylation reaction, temperature monitoring revealed an optimum temperature for the coupling where side reactions are diminished. The strategy allows for the expeditious

1
2
3 preparation of a diverse set of densely substituted arabinoxylans in good overall yields as
4
5 valuable substrates for characterization of hemicellulose-degrading enzymes.
6
7

10 EXPERIMENTAL SECTION

11
12 **General Information.** Starting materials, reagents and solvents were purchased from
13
14 commercial suppliers and used without further purification unless otherwise noted. All solvents
15
16 were of analytical HPLC grade. Anhyd solvents were obtained from an Innovative Technology
17
18 PS-MD-7 PureSolv solvent purification system except for CH_2Cl_2 and toluene for
19
20 glycosylations, which were dried over 3\AA mol sieves prior to use. All reactions were performed
21
22 under inert atm (N_2) in oven-dried glassware. TLC was carried out using Merck aluminium
23
24 sheets pre-coated with 0.25 mm silica gel, C-60 F₂₅₄ plates. TLC plates were inspected under
25
26 UV light or visualized by charring after dipping in a cerium ammonium sulfate solution (1%
27
28 cerium(IV)sulfate and 2.5% ammonium heptamolybdate in a 10% sulfuric acid solution). Flash
29
30 column chromatography was performed using Merck Geduran silica gel 60 \AA (40–63 μm) while
31
32 dry column chromatography⁴² was accomplished using Merck silica gel 60 \AA (15–40 μm).
33
34
35
36
37 NMR spectra were recorded with a Bruker Ascend™ 400 or a Bruker Ascend™ 800
38
39 spectrometer. The chemical shift are reported in ppm relative to the residual solvent peak from
40
41 CDCl_3 ($\delta_{\text{H}} = 7.26$ ppm, $\delta_{\text{C}} = 77.16$ ppm), CD_3OD ($\delta_{\text{H}} = 4.87$ ppm, $\delta_{\text{C}} = 49.00$ ppm) or D_2O (δ_{H}
42
43 = 4.79 ppm). Assignment of ^1H and ^{13}C resonances were based on APT, DQF-COSY, HSQC,
44
45 H2BC, HMBC, TOCSY and HSQC-TOSCY experiments. Xylose residues are numbered A – D
46
47 from the reducing end while arabinose substituents are labelled E – G as shown in Scheme 6 –
48
49
50
51
52 8. Optical rotation was measured on a Perkin Elmer Model 341 polarimeter. HRMS analysis
53
54
55
56
57
58
59
60

1
2
3 was performed on either a UHPLC-QTOF system (Dionex UltiMate 3000 and Bruker MaXis)
4
5 with an electrospray ionization (ESI) source or a MALDI-TOF system (Bruker Solarix XR 7T).
6
7
8
9

10 *Phenyl 1-thio-β-D-xylopyranoside (7)*. D-Xylose (29.9 g, 199 mmol), Et₃N (223 mL, 160 mmol)
11
12 and DMAP (4.90 g, 40.1 mmol) were suspended in CH₂Cl₂ (300 mL) at 0 °C. Upon dropwise
13
14 addition of Ac₂O (94.0 mL, 996 mmol) the solution turned yellow. After stirring for 2 h at rt
15
16 TLC analysis (hexane/EtOAc 1:1) showed the consumption of the starting material and
17
18 formation of the product. The mixture was washed with 1 M HCl and brine followed by
19
20 concentration in vacuo. The crude xylose tetraacetate was dissolved in dry CH₂Cl₂ (160 mL)
21
22 followed by addition of PhSH (25.0 mL, 243 mmol) and BF₃·OEt₂ (76.0 mL, 600 mmol) at 0 °C
23
24 under N₂ atm. The reaction was stirred overnight at rt where it turned purple. TLC analysis
25
26 (hexane/EtOAc 2:1) showed full consumption of the starting material and formation of the
27
28 product. The solution was diluted with CH₂Cl₂ and washed with saturated aq NaHCO₃ and
29
30 brine, dried over Na₂SO₄, filtered and concd in vacuo. The residue was dissolved in methanol
31
32 (160 mL) and solid sodium was added until a basic medium was obtained. After 1 h TLC
33
34 analysis (hexane/EtOAc 2:1) showed the consumption of the starting material and formation of
35
36 the product. The solution was neutralized with Amberlite IR-120 (H⁺), filtered and concd in
37
38 vacuo. The residue was purified by dry column chromatography (10% acetone in toluene, 5%
39
40 gradient) followed by crystallization from hexane and acetone to afford a white crystalline
41
42 product (19.2 g, 40%). *R*_f 0.27 (toluene/acetone 1:1). Mp 142.3–143.8 °C (lit.²⁰ 143–145 °C). ¹H
43
44 NMR (400 MHz, CD₃OD) δ 7.54–7.51 (m, 2H), 7.34–7.26 (m, 3H), 4.56 (d, *J*_{1,2} = 9.3 Hz, 1H,
45
46 H1), 3.95 (dd, *J*_{5,5'} = 11.3 Hz, *J*_{4,5} = 5.2 Hz, 1H, H5), 3.51–3.45 (m, 1H, H4), 3.38–3.31 (m, 3H,
47
48 H3, OH), 3.26–3.19 (m, 2H, H2, H5'). ¹³C{¹H} NMR (101 MHz, CD₃OD) δ 134.9, 133.1 (×2),
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 129.9 (×2), 128.5, 90.1 (C1), 79.2 (C3), 73.7 (C2), 70.9 (C4), 70.4 (C5). NMR data are in
4
5 accordance with literature values.⁴³
6
7

8
9
10 *Phenyl 2,3,4-tri-O-benzoyl-1-thio-β-D-xylopyranoside (5)*. Triol **7** (5.12 g, 21.1 mmol) was
11
12 dissolved in pyridine (45 mL) and BzCl (7.4 mL, 63.4 mmol) was added. The reaction was
13
14 stirred at rt for 1 h and excess of BzCl was quenched by the addition of methanol (10 mL)
15
16 followed by stirring for an additional 10 min. The mixture was diluted with CH₂Cl₂ and washed
17
18 with 1 M HCl and water, dried over Na₂SO₄, filtered, and concd in vacuo. The residue was
19
20 purified by flash column chromatography to afford **5** as white crystals (5.1 g, 84%). *R*_f 0.28
21
22 (heptane/EtOAc 7:3). Mp 102.9–104.3 °C (EtOH). ¹H NMR (400 MHz, CDCl₃) δ 8.05 (dd, *J*=
23
24 8.2, 1.1 Hz, 2H), 8.01–7.98 (m, 4H), 7.56–7.51 (m, 5H), 7.42–7.32 (m, 9H), 5.78 (t, *J* = 6.6 Hz,
25
26 1H, H3), 5.46 (t, *J* = 6.3 Hz, 1H, H2), 5.33–5.26 (m, 2H, H1, H4), 4.71 (dd, *J*_{5eq,5ax} = 12.3 Hz,
27
28 *J*_{5eq,4} = 4.0 Hz, 1H, H5_{eq}), 3.83 (dd, *J*_{5ax,5eq} = 12.3 Hz, *J*_{5ax,4} = 6.5 Hz, 1H, H5_{ax}). ¹³C {¹H} NMR
29
30 (101 MHz, CDCl₃) δ 165.6, 165.3, 165.3, 133.6, 133.5, 133.5, 133.2, 132.8 (×2), 130.2 (×2),
31
32 130.1 (×2), 130.1 (×2), 129.3 (×2), 129.2 (×2), 129.1, 128.6 (×2), 128.6 (×2), 128.5 (×2), 128.3,
33
34 86.5 (C1), 70.6 (C3), 70.1 (C2), 68.8 (C4), 63.7 (C5). Anal. calcd for C₃₂H₂₆O₇S: C, 69.30; H,
35
36 4.73; S, 5.78. Found: C, 69.26; H, 4.70; S, 5.66. NMR data are in accordance with literature
37
38 values.⁴⁴
39
40
41
42
43
44
45
46

47 *Phenyl 4-O-(p-methoxy)benzyl-1-thio-β-D-xylopyranoside (8)*. To a solution of triol **7** (29.5 g,
48
49 122 mmol) in dry DMF (200 mL) was added CSA (2.83 g, 12.0 mmol) and the mixture was
50
51 heated to 60 °C. 2-Methoxy prop-1-ene (37.5 mL, 366 mmol) was added slowly to the solution.
52
53 The reaction was stirred for 1 h and then cooled to rt and quenched with Et₃N (30 mL).
54
55
56
57
58
59
60

Concentration in vacuo followed by purification of the residue by flash column chromatography (heptane/EtOAc/CH₂Cl₂ 4:1:1) afforded phenyl 2,3-*O*-isopropylidene-1-thio-β-D-xylopyranoside as a colorless oil (24.1 g, 70%). *R*_f 0.62 (toluene/acetone 3:1). ¹H NMR (400 MHz, CDCl₃) δ 7.58–7.55 (m, 2H), 7.34–7.30 (m, 3H), 4.80 (d, *J*_{1,2} = 9.6 Hz, 1H, H1), 4.13 (dd, *J*_{5,5'} = 11.6 Hz, *J*_{4,5} = 5.3 Hz, 1H, H5_{eq}), 4.03–3.96 (m, 1H, H4), 3.54 (t, *J* = 9.1 Hz, 1H, H3), 3.27–3.22 (m, 2H, H2, H5_{ax}), 2.17 (d, *J*_{OH,4} = 4.0 Hz, 1H, OH), 1.49 (s, 3H), 1.45 (s, 3H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 133.0 (×2), 132.0, 129.0 (×2), 128.3, 111.5, 85.6 (C1), 83.0 (C3), 75.3 (C2), 70.0 (C5), 69.2 (C4), 26.8, 26.7. NMR data are in accordance with literature values.⁴⁵ The intermediate (17.5 g, 62.0 mmol) was dissolved in dry DMF (120 mL) and NaH (3.0 g, 74.4 mmol, 60% in mineral oil) was added at 0 °C. After 10 min PMBCl (10.9 mL, 80.6 mmol, 1.3 equiv.) was added. The mixture was stirred at rt for 16 h and then quenched with 10% HCl (28 mL) and diluted with CH₂Cl₂. The solution was washed with saturated aq NaHCO₃, dried over Na₂SO₄, filtered, and concd in vacuo. The crude residue was dissolved in CH₂Cl₂/MeOH 1:1 (200 mL), CSA (14.4 g, 62.0 mmol) was added and the mixture was allowed to stir at rt overnight. The reaction was quenched by addition of Et₃N and concd in vacuo. The residue was purified by flash column chromatography (heptane/EtOAc 3:2) to furnish compound **8** as a white crystalline solid (20.5 g, 91% over 2 steps). *R*_f 0.47 (hexane/EtOAc 1:1). Mp 72.7–75.6 °C (EtOH). [*α*]_D²⁵ –55.9 (*c* 1, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 7.53–7.50 (m, 2H), 7.33–7.29 (m, 3H), 7.27–7.23 (m, 2H), 6.90–6.86 (m, 2H), 4.61 (d, *J*_{gem} = 11.4 Hz, 1H, PMB), 4.58–4.54 (m, 2H, H1 (*J*_{1,2} = 8.9 Hz) PMB), 4.07 (dd, *J*_{5eq,5ax} = 11.5 Hz, *J*_{5eq,4} = 4.8 Hz, 1H, H5_{eq}), 3.80 (s, 3H, OMe), 3.66 (t, *J* = 8.4 Hz, 1H, H3), 3.49–3.43 (m, 1H, H4), 3.40 (t, *J* = 8.6 Hz, 1H, H2), 3.27 (dd, *J*_{5ax,5eq} = 11.5 Hz, *J*_{5ax,4} = 9.7 Hz, 1H, H5_{ax}), 2.52 (bs, 2H, OH). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 159.7, 132.8 (×2), 132.3, 130.1, 129.7 (×2), 129.2 (×2),

1
2
3 128.3, 114.2 ($\times 2$), 88.8 (C1), 76.6 (C4), 76.5 (C3), 72.8 (PMB), 72.1 (C2), 67.1 (C5), 55.4
4
5 (OMe). HRMS: m/z calcd for $C_{19}H_{22}O_5SNa$ $[M + Na]^+$ 385.1080, found 385.1090.
6
7
8
9

10 *Phenyl 2,3-di-O-benzoyl-1-thio- β -D-xylopyranoside (6)*. Compound **8** (1.97 g, 5.43 mmol) was
11 dissolved in CH_2Cl_2 (32 mL) and Et_3N (2.4 mL, 17.2 mmol), DMAP (0.069 g, 0.56 mmol) and
12 BzCl (1.9 mL, 16.4 mmol) were added at $0^\circ C$. After 5 min the reaction was left to stir at rt
13 overnight. 3-(Dimethylamino)-1-propylamine (2.2 mL, 17.5 mmol) was added and the mixture
14 was stirred for 1.5 h followed by washing with 1 M HCl and brine, drying over $MgSO_4$,
15 filtration, and concentration in vacuo. The product phenyl 2,3-di-O-benzoyl-4-O-(*p*-
16 methoxy)benzyl-1-thio- β -D-xylopyranoside (3.12 g, 100%) was obtained as a light yellow syrup
17 and used without further purification. R_f 0.28 (hexane/EtOAc 4:1). $[\alpha]^{25}_D +62.0$ (c 1.0, $CHCl_3$).
18
19 1H NMR (400 MHz, $CDCl_3$) β 7.99–7.95 (m, 3H), 7.56–7.27 (m, 12H), 7.15–7.12 (m, 2H),
20 6.74–6.71 (m, 2H), 5.60 (t, $J = 8.0$ Hz, 1H, H3), 5.34 (t, $J = 8.0$ Hz, 1H, H2), 5.04 (d, $J_{1,2} = 8.1$
21 Hz, 1H, H1), 4.56 (d, $J_{gem} = 11.9$ Hz, 1H, PMB), 4.52 (d, $J_{gem} = 11.8$ Hz, 1H, PMB), 4.28 (dd,
22 $J_{5eq,5ax} = 11.9$ Hz, $J_{5eq,4} = 4.6$ Hz, 1H, H5_{eq}), 3.80–3.76 (m, 1H, H4), 3.75 (s, 3H, OMe), 3.57
23 (dd, $J_{5ax,5eq} = 11.9$ Hz, $J_{5ax,4} = 8.5$ Hz, 1H, H5_{ax}). ^{13}C $\{^1H\}$ NMR (101 MHz, $CDCl_3$) δ 165.7,
24 165.4, 159.5, 133.4 ($\times 2$), 133.2, 132.5 ($\times 2$), 130.1 ($\times 2$), 130.0 ($\times 2$), 129.7, 129.7 ($\times 2$), 129.5,
25 129.4, 129.1 ($\times 2$), 128.5 ($\times 2$), 128.5 ($\times 2$), 128.1, 113.9 ($\times 2$), 87.0 (C1), 73.9 (C4), 73.6 (C3),
26 72.5 (PMB), 70.6 (C2), 66.4 (C5), 55.4 (OMe). HRMS: m/z calcd for $C_{33}H_{30}O_7SNa$ $[M + Na]^+$
27 593.1604, found 593.1617. NMR data are in accordance with literature values.¹³ To a well
28 stirred emulsion of the above PMB ether (0.696 g, 1.22 mmol) in CH_2Cl_2 (10 mL) and H_2O (1.0
29 mL), DDQ (0.415 g, 1.83 mmol) was added. The reaction was stirred in the dark for 7 h and
30 then filtered through a large pad of Celite. The filtrate was washed with saturated aq $NaHCO_3$
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 and brine, filtered and concd in vacuo. Purification by flash column chromatography
4
5 (pentane/EtOAc 4:1) yielded **6** as a white crystalline compound (0.514 g, 94%). R_f 0.53
6
7 (hexane/EtOAc 2:1). Mp 133.1–134.0 °C (EtOH). $[\alpha]^{25}_D$ +66.1 (c 1.0, CHCl_3). ^1H NMR (400
8
9 MHz, CDCl_3) δ 8.05–7.99 (m, 4H), 7.57–7.49 (m, 4H), 7.43–7.39 (m, 4H), 7.35–7.29 (m, 3H),
10
11 5.43 (t, $J = 7.4$ Hz, 1H, H2), 5.33 (t, $J = 7.4$ Hz, 1H, H3), 5.09 (d, $J_{1,2} = 7.3$ Hz, 1H, H1), 4.45
12
13 (dd, $J_{5\text{eq},5\text{ax}} = 12.0$ Hz, $J_{5\text{eq},4} = 4.4$ Hz, 1H, H5_{eq}), 4.00 (q, $J = 6.9$ Hz, 1H, H4), 3.61 (dd, $J_{5\text{ax},5\text{eq}} =$
14
15 12.0 Hz, $J_{5\text{ax},4} = 7.9$ Hz, 1H, H5_{ax}), 3.03 (s, 1H, OH). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 167.1,
16
17 165.2, 133.8, 133.6, 133.0, 132.8 ($\times 2$), 130.2 ($\times 2$), 130.0 ($\times 2$), 129.3, 129.2 ($\times 2$), 128.9, 128.7
18
19 ($\times 2$), 128.7 ($\times 2$), 128.3, 86.8 (C1), 76.0 (C3), 70.2 (C2), 68.4 (C4), 67.6 (C5). HRMS: m/z calcd
20
21 for $\text{C}_{25}\text{H}_{22}\text{O}_6\text{SNa}$ $[\text{M} + \text{Na}]^+$ 473.1029, found 473.1048.
22
23
24
25
26
27
28

29 *Phenyl 2,3-di-O-levulinoyl-1-thio-β-D-xylopyranoside (9)*. A solution of diol **8** (1.98 g, 5.47
30
31 mmol) and 4-oxopentanoic acid (1.61 g, 13.8 mmol) in CH_2Cl_2 (8.5 mL) was cooled to 0 °C.
32
33 Then a solution of DMAP (0.070 g, 0.58 mmol) and DDC (3.42 g, 16.8 mmol) in CH_2Cl_2 (1.0
34
35 mL) was added. The reaction stirred at rt overnight and then filtered and concd. The residue was
36
37 dissolved in CH_2Cl_2 (20 mL) and H_2O (2.0 mL) and stirred vigorously. DDQ (1.86 g, 8.21
38
39 mmol) was added and the mixture was stirred at rt for 4.5 h. Then the solution was filtered
40
41 through a large pad of Celite and washed with saturated aq NaHCO_3 and brine, dried over
42
43 Na_2SO_4 , filtered, and concd in vacuo. The residue was purified by flash column chromatography
44
45 (hexane/EtOAc 1:1) to provide **9** as a light yellow wax (1.78 g, 74% over 2 steps). R_f 0.58
46
47 (EtOAc). $[\alpha]^{25}_D$ –23.4 (c 1.0, CHCl_3). ^1H NMR (400 MHz, CDCl_3) δ 7.48–7.46 (m, 2H),
48
49 7.33–7.29 (m, 3H), 5.02 (t, $J = 8.4$ Hz, 1H, H3), 4.94 (t, $J = 8.7$ Hz, 1H, H2), 4.73 (d, $J_{1,2} = 8.8$
50
51 Hz, 1H, H1), 4.19 (dd, $J_{5\text{eq},5\text{ax}} = 11.7$ Hz, $J_{5\text{eq},4} = 5.1$ Hz, 1H, H5_{eq}), 3.84–3.79 (m, 1H, H4), 3.39
52
53
54
55
56
57
58
59
60

(dd, $J_{5_{ax},5_{eq}} = 11.7$ Hz, $J_{5_{ax},4} = 9.5$ Hz, 1H, H5_{ax}), 3.16 (bs, 1H, OH), 2.89–2.51 (m, 8H), 2.17 (2 × s, 6H). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 208.0, 206.4, 173.0, 171.4, 132.7, 132.6 (×2), 129.1 (×2), 128.2, 86.8 (C1), 76.5 (C3), 69.7 (C2), 68.6 (C5), 68.5 (C4), 38.4, 37.9, 29.9, 29.9, 28.3, 28.1. HRMS: m/z calcd for $\text{C}_{21}\text{H}_{26}\text{O}_8\text{SNa}$ $[\text{M} + \text{Na}]^+$ 461.1241, found 461.1242.

Phenyl 2-O-benzoyl-4-O-(p-methoxy)benzyl-1-thio- β -D-xylopyranoside (10). To a vigorously stirred solution of diol **8** (1.0 g, 2.76 mmol), tetrabutylammonium hydrogen sulfate (0.187 g, 0.55 mmol) and BzCl (0.43 mL, 3.72 mmol) in CH_2Cl_2 (50 mL) at -5 °C was added a 1 M aq NaOH solution (7.0 mL, 7.0 mmol). The mixture was stirred for 30 min (TLC indicated traces of **8** and the dibenzoylated product). The organic layer was separated, washed with H_2O , dried over Na_2SO_4 , filtered, and concd in vacuo. The residue was purified by flash column chromatography (hexane/EtOAc 8.5:1.5) affording compound **10** as white crystals (0.877 g, 68%). R_f 0.35 (hexane/EtOAc 3:1). Mp 101.7–102.4 °C (EtOH). $[\alpha]^{25}_{\text{D}} -13.2$ (c 1.0, CHCl_3). ^1H NMR (400 MHz, CDCl_3) δ 8.09–8.07 (m, 2H), 7.61–7.57 (m, 1H), 7.48–7.41 (m, 4H), 7.29–7.24 (m, 5H), 6.89–6.86 (m, 2H), 5.04 (dd, $J_{1,2} = 9.5$ Hz, $J_{2,3} = 9.0$ Hz, 1H, H2), 4.78 (d, $J_{1,2} = 9.6$ Hz, 1H, H1), 4.65 (d, $J_{\text{gem}} = 11.5$ Hz, 1H, PMB), 4.60 (d, $J_{\text{gem}} = 11.5$ Hz, 1H, PMB), 4.10 (dd, $J_{5_{eq},5_{ax}} = 11.5$ Hz, $J_{5_{eq},4} = 5.1$ Hz, 1H, H5_{eq}), 3.85 (t, $J = 8.8$ Hz, 1H, H3), 3.81 (s, 3H, OMe), 3.61–3.55 (m, 1H, H4), 3.30 (dd, $J_{5_{ax},5_{eq}} = 11.5$ Hz, $J_{5_{ax},4} = 10.1$ Hz, 1H, H5_{ax}), 2.62 (bs, 1H, OH). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 166.1, 159.7, 133.5, 132.8 (×2), 132.7, 130.1 (×2), 130.1, 129.8, 129.7 (×2), 129.1 (×2), 128.6 (×2), 128.2, 114.2 (×2), 86.8 (C1), 77.4 (C4), 76.1 (C3), 73.1 (PMB), 73.0 (C2), 67.6 (C5), 55.4 (OMe). HRMS: m/z calcd for $\text{C}_{26}\text{H}_{26}\text{O}_6\text{SNa}$ $[\text{M} + \text{Na}]^+$ 489.1342, found 489.1343.

1
2
3
4 *Phenyl 2-O-benzoyl-3-O-levulinoyl-1-thio-β-D-xylopyranoside (11)*. To a solution of **10** (3.94 g,
5
6 8.45 mmol) and 4-oxopentanoic acid (1.22 g, 10.5 mmol) in CH₂Cl₂ (14 mL) cooled to 0 °C was
7
8 added DMAP (0.104 g, 0.85 mmol) and DCC (2.61 g, 12.6 mmol). The solution was left to stir
9
10 at rt for 3.5 h, then filtered and concd in vacuo. The residue was dissolved in CH₂Cl₂ (60 mL)
11
12 and H₂O (6.0 mL) and stirred vigorously. DDQ (2.88 g, 12.7 mmol) was added and the mixture
13
14 was left to stir at rt overnight in the dark. Then the solution was filtered through a large pad of
15
16 Celite and washed with saturated aq NaHCO₃ and brine, dried over Na₂SO₄, filtered, and concd
17
18 in vacuo. The residue was purified by flash column chromatography (hexane/EtOAc 2:1)
19
20 affording **11** as a white crystalline solid (2.935 g, 78% over 2 steps). *R*_f 0.28 (hexane/EtOAc
21
22 1:1). Mp 88.7–90.3 °C (EtOH). [α]_D²⁵ –1.8 (*c* 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ
23
24 8.03–8.01 (m, 2H), 7.59 (t, *J* = 7.4 Hz, 1H), 7.47–7.44 (m, 4H), 7.29–7.28 (m, 3H), 5.23–5.16
25
26 (m, 2H, H₂, H₃), 4.94–4.91 (m, 1H, H₁), 4.29 (dd, *J*_{5eq,5ax} = 11.8 Hz, *J*_{5eq,4} = 4.9 Hz, 1H, H_{5eq}),
27
28 3.94–3.89 (m, 1H, H₄), 3.49 (dd, *J*_{5ax,5eq} = 11.7 Hz, *J*_{5ax,4} = 9.1 Hz, 1H, H_{5ax}), 3.14 (bs, 1H,
29
30 OH), 2.81–2.65 (m, 2H), 2.58–2.51 (m, 1H), 2.44–2.37 (m, 1H), 2.11 (s, 3H). ¹³C {¹H} NMR
31
32 (101 MHz, CDCl₃) δ 207.7, 172.8, 165.3, 133.6, 132.9, 132.7 (×2), 130.0 (×2), 129.4, 129.1
33
34 (×2), 128.7 (×2), 128.2, 87.0 (C₁), 76.4 (C₃), 70.3 (C₂), 68.5 (C₄), 68.3 (C₅), 38.5, 29.8, 28.3.
35
36 HRMS: *m/z* calcd for C₂₃H₂₄O₇SNa [M + Na]⁺ 467.1135, found 467.1136.
37
38
39
40
41
42
43
44

45 **Temperature Screening (General Procedure A)**. Crushed mol sieves (3Å, 1.0 g) were added a
46
47 2-neck Schlenk flask, where the middle neck was fitted with a glass stopper and the other with a
48
49 septum. The flask was placed under vacuum, heated with a heatgun and then subjected to an atm
50
51 of N₂ and cooled to rt. Donor **5** (0.200 g, 0.36 mmol) dissolved in dry CH₂Cl₂ (3.0 mL) was
52
53 added to the flask together with AgOTf (0.185 g, 0.72 mmol) dissolved in dry toluene (2.0 mL).
54
55
56
57
58
59
60

1
2
3 Stirring of the reaction was initiated and the glass stopper was exchanged for a thermometer,
4
5 and the solution was cooled to $-65\text{ }^{\circ}\text{C}$. $p\text{NO}_2\text{PhSCl}$ (0.068 g, 0.36 mmol) was dissolved in dry
6
7 CH_2Cl_2 (0.5 mL) and slowly added followed by stirring for an additional 10 min. Acceptor **6**
8
9 (0.146 g, 0.33 mmol) was dissolved in dry CH_2Cl_2 (0.5 mL) and added quickly. The temperature
10
11 was raised as indicated in Figure 1, where the reaction was allowed to warm to $-15\text{ }^{\circ}\text{C}$ at which
12
13 point Et_3N (0.15 mL, 1.08 mmol) was added. The solution was filtered through a pad of Celite
14
15 and concd in vacuo. The residue was purified by flash column chromatography to afford the
16
17 product **12**.
18
19
20
21
22
23

24 *Phenyl 2,3,4-tri-O-benzoyl- β -D-xylopyranosyl-(1 \rightarrow 4)-2,3-di-O-benzoyl-1-thio- β -D-*
25
26 *xylopyranoside (12)*. To a Schlenk flask containing 3Å mol sieves (1.20 g) was added donor **5**
27
28 (0.200 g, 0.36 mmol) dissolved in dry CH_2Cl_2 (6.0 mL) and AgOTf (0.185 g, 0.72 mmol)
29
30 dissolved in dry Et_2O (5.0 mL). The mixture was stirred for 1 h at rt and then cooled to $-78\text{ }^{\circ}\text{C}$
31
32 at which point PhSCl (0.062 g, 0.43 mmol) or $p\text{NO}_2\text{PhSCl}$ (0.082 g, 0.43 mmol) dissolved in
33
34 dry CH_2Cl_2 (1.0 mL) was added dropwise followed by stirring for 10 min. Acceptor **6** (0.146 g,
35
36 0.33 mmol) dissolved in dry CH_2Cl_2 (0.5 mL) was then added slowly. The reaction was stirred
37
38 at $-78\text{ }^{\circ}\text{C}$ for 15 min, then for 15 min at rt after which Et_3N (0.21 mL, 2.88 mmol) was added.
39
40
41 The mixture was filtered through a pad of Celite, concd in vacuo, and purified by flash column
42
43 chromatography (pentane/ EtOAc 3:2) to give **12** as a white amorphous solid (0.186 g, 64% with
44
45 PhSCl and 0.229 g, 79% with $p\text{NO}_2\text{PhSCl}$). Alternatively, general procedure A was employed
46
47 to afford 0.260 g (90%) of **12**. R_f 0.18 (heptane/ EtOAc 7:3). $[\alpha]_D^{25} -22$ (c 1.0, CHCl_3). $^1\text{H NMR}$
48
49 (400 MHz, CDCl_3) δ 8.02–7.92 (m, 10H), 7.56–7.50 (m, 5H), 7.45–7.27 (m, 15H), 5.69–5.62
50
51 (m, 2H, $\text{H}3^{\text{A}}$, $\text{H}3^{\text{B}}$), 5.34 (t, $J = 8.0$ Hz, 1H, $\text{H}2^{\text{A}}$), 5.24 (dd, $J_{2\text{B},3\text{B}} = 6.5$ Hz, $J_{1\text{B},2\text{B}} = 5.0$ Hz, 1H,
52
53
54
55
56
57
58
59
60

1
2
3 H2^B), 5.06–5.01 (m, 2H, H1^A, H4^B), 4.95 (d, $J_{1B,2B} = 4.8$ Hz, 1H, H1^B), 4.27 (dd, $J_{5Aeq,5Aax} =$
4
5 12.2 Hz, $J_{5Aeq,4A} = 4.7$ Hz, 1H, H5^A_{eq}), 4.09 (td, $J = 8.3$ Hz, $J_{5Aeq,4A} = 4.9$ Hz, 2H, H4^A), 4.03
6
7 (dd, $J_{5Beq,5Bax} = 12.4$ Hz, $J_{5Beq,4B} = 3.9$ Hz, 1H, H5^B_{eq}), 3.55 (dd, $J_{5Aax,5Aeq} = 12.1$ Hz, $J_{5Aax,4A} =$
8
9 8.6 Hz, 1H, H5^A_{ax}), 3.43 (dd, $J_{5Bax,5Beq} = 12.4$ Hz, $J_{5Bax,4B} = 6.3$ Hz, 1H, H5^B_{ax}). ¹³C{¹H} NMR
10
11 (101 MHz, CDCl₃) δ 165.6, 165.5, 165.4, 165.4, 165.2, 133.5 ($\times 2$), 133.5, 133.4 ($\times 2$), 132.8,
12
13 132.7 ($\times 2$), 130.1 ($\times 2$), 130.0 ($\times 2$), 130.0 ($\times 2$), 129.9 ($\times 2$), 129.9 ($\times 2$), 129.5, 129.4, 129.3,
14
15 129.2, 129.1 ($\times 2$), 129.0, 128.6 ($\times 6$), 128.5 ($\times 4$), 128.2, 99.8 (C1^B), 86.8 (C1^A), 75.2 (C4^A), 73.1
16
17 (C3^A), 70.6 (C2^A), 70.2 (C2^B), 69.7 (C3^B), 68.6 (C4^B), 65.6 (C5^A), 60.9 (C5^B). HRMS: m/z
18
19 calcd for C₅₁H₄₂O₁₃SNa [M + Na]⁺ 917.2238, found 917.2258.
20
21
22
23
24
25

26
27 *Phenyl 2,3,4-tri-O-benzoyl- β -D-xylopyranosyl-(1 \rightarrow 4)-2,3-di-O-benzoyl- β -D-xylopyranosyl-*
28
29 *(1 \rightarrow 4)-2,3-di-O-benzoyl-1-thio- β -D-xylopyranoside (13).* To a Schlenk flask containing 3Å mol
30
31 sieves (1.20 g) was added donor **5** (0.200 g, 0.36 mmol) dissolved in dry CH₂Cl₂ (6.0 mL) and
32
33 AgOTf (0.185 g, 0.72 mmol) dissolved in dry Et₂O (5.0 mL). The mixture was stirred for 1 h at
34
35 rt and then cooled to -78 °C at which point PhSCl (0.062 g, 0.43 mmol) or *p*NO₂PhSCl (0.082
36
37 g, 0.43 mmol) dissolved in dry CH₂Cl₂ (1.0 mL) was added dropwise followed by stirring for 10
38
39 min. Acceptor **6** (0.146 g, 0.33 mmol) dissolved in dry CH₂Cl₂ (0.5 mL) was then added slowly.
40
41 The reaction was stirred at -78 °C for 15 min and then for 15 min at rt. AgOTf (0.093 g, 0.36
42
43 mmol) dissolved in dry Et₂O (2.0 mL) was added and the mixture was cooled to -78 °C over 20
44
45 min. PhSCl (0.058 g, 0.40 mmol) or *p*NO₂PhSCl (0.075 g, 0.40 mmol) dissolved in dry CH₂Cl₂
46
47 (1.0 mL) was added dropwise followed by stirring for 10 min. Acceptor **6** (0.132 g, 0.29 mmol)
48
49 dissolved in dry CH₂Cl₂ (0.4 mL) was added slowly. The reaction was stirred at -78 °C for 15
50
51 min, then for 15 min at rt after which Et₃N (0.4 mL, 5.44 mmol) was added. The mixture was
52
53
54
55
56
57
58
59
60

1
2
3 filtered through a pad of Celite, concd in vacuo, and purified by flash column chromatography
4 (pentane/acetone 7.5:2.5) to give a mixture of **12** (0.169 g, 59%) and **13** (0.134 g, 34%) (with
5 PhSCl) or only **13** as a white amorphous solid (0.249 g, 69%, with *p*NO₂PhSCl). ¹H NMR (400
6 MHz, CDCl₃) δ 7.99–7.89 (m, 14H), 7.57–7.47 (m, 7H), 7.42–7.25 (m, 19H), 5.62–5.57 (m, 2H,
7 H^{3A}, H^{3C}), 5.52 (t, *J* = 8.1 Hz, 1H, H^{3B}), 5.31–5.27 (m, 1H, H^{2A}), 5.19–5.14 (m, 2H, H^{2B},
8 H^{2C}), 5.01 (td, *J* = 6.4 Hz, *J* = 4.1 Hz, 1H, H^{4C}), 4.97 (d, *J* = 8.1 Hz, 1H, H^{1A}), 4.74–4.71 (m,
9 2H, H^{1B}, H^{1C}), 4.14 (dd, *J*_{5Aeq,5Aax} = 12.0 Hz, *J*_{5Aeq,4A} = 4.7 Hz, 1H, H^{5A}_{eq}), 4.00–3.94 (m, 2H,
10 H^{4A}, H^{5C}_{eq}), 3.82 (td, *J* = 8.4, 5.0 Hz, 1H, H^{4B}), 3.55 (dd, *J*_{5Beq,5Bax} = 12.3 Hz, *J*_{5Beq,4B} = 4.8 Hz,
11 1H, H^{5B}_{eq}), 3.46 (dd, *J*_{5Aax,5Aeq} = 12.1 Hz, *J*_{5Aax,4A} = 8.7 Hz, 1H, H^{5A}_{ax}), 3.35 (dd, *J*_{5Cax,5Ceq} =
12 12.3 Hz, *J*_{5Cax,4C} = 6.3 Hz, 1H, H^{5C}_{ax}), 3.16 (dd, *J*_{5Bax,5Beq} = 12.3 Hz, *J*_{5Bax,4B} = 8.6 Hz, 1H,
13 H^{5B}_{ax}). ¹³C {¹H} NMR (101 MHz, CDCl₃) δ 165.5, 165.5, 165.5, 165.4, 165.3, 165.2, 165.0,
14 133.5 (×2), 133.5, 133.5, 133.4 (×2), 133.3, 132.8, 132.6 (×2), 130.1 (×2), 130.0 (×2), 130.0
15 (×2), 129.9 (×2), 129.9 (×2), 129.8 (×2), 129.8 (×2), 129.6, 129.5, 129.5, 129.3 (×2), 129.2,
16 129.1 (×2), 129.0, 128.6 (×2), 128.6 (×2), 128.6 (×4), 128.5 (×2), 128.5 (×2), 128.4 (×2), 128.2,
17 101.1 (C^{1B}), 99.6 (C^{1C}), 86.8 (C^{1A}), 75.8 (C^{4A}), 74.9 (C^{4B}), 73.1 (C^{3A}), 72.1 (C^{3B}), 71.6 (C^{2B}),
18 70.5 (C^{2A}), 70.2 (C^{2C}), 69.7 (C^{3C}), 68.6 (C^{4C}), 65.9 (C^{5A}), 62.3 (C^{5B}), 60.9 (C^{5C}). HRMS: *m/z*
19 calcd for C₇₀H₅₈O₁₉SNa [M + Na]⁺ 1257.3185, found 1257.3192.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 *Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-2-O-benzoyl-3-O-levulinoyl-1-thio-β-D-*
46 *xylopyranoside (14)*. Disaccharide **14** was obtained in 86% yield as a light yellow amorphous
47 solid following general procedure A. *R*_f 0.42 (hexane/EtOAc 1:1). [α]_D²⁵ –60.4 (*c* 1.0, CHCl₃).
48 ¹H NMR (400 MHz, CDCl₃) δ 8.05–8.00 (m, 4H), 7.97–7.92 (m, 4H), 7.62–7.45 (m, 4H),
49 7.42–7.32 (m, 10H), 7.28–7.25 (m, 3H), 5.70 (t, *J* = 6.5 Hz, 1H, H^{3B}), 5.40 (t, *J* = 8.3 Hz, 1H,
50
51
52
53
54
55
56
57
58
59
60

1
2
3 H3^A), 5.28–5.23 (m, 2H, H2^B, H4^B), 5.16 (t, $J = 8.4$ Hz, 1H, H2^A), 4.94 (d, $J_{1B,2B} = 4.7$ Hz, 1H,
4 H1^B), 4.89 (d, $J_{1A,2A} = 8.6$ Hz, 1H, H1^A), 4.48 (dd, $J_{5Beq,5Bax} = 12.4$, Hz, $J_{5Beq,4B} = 3.8$ Hz, 1H,
5 H5^B_{eq}), 4.19 (dd, $J_{5Aeq,5Aax} = 12.0$ Hz, $J_{5Aeq,4A} = 4.9$ Hz, 1H, H5^A_{eq}), 3.99 (td, $J = 8.7$, Hz, $J_{5Aeq,4A}$
6 = 5.0 Hz, 1H, H4^A), 3.75 (dd, $J_{5Bax,5Beq} = 12.4$ Hz, $J_{5Bax,4B} = 6.1$ Hz, 1H, H5^B_{ax}), 3.44 (dd,
7 $J_{5Aax,5Aeq} = 12.0$ Hz, $J_{5Aax,4A} = 9.1$ Hz, 1H, H5^A_{ax}), 2.65–2.41 (m, 4H), 1.97 (s, 3H). ¹³C {¹H}
8 NMR (101 MHz, CDCl₃) δ 205.9, 171.8, 165.6, 165.4, 165.4, 165.2, 133.6, 133.5 ($\times 2$), 133.5,
9 132.8 ($\times 2$), 132.5, 130.1 ($\times 2$), 130.1 ($\times 2$), 130.1 ($\times 2$), 130.0 ($\times 2$), 129.5, 129.4, 129.2, 129.1,
10 129.1 ($\times 2$), 128.6 ($\times 2$), 128.6 ($\times 2$), 128.6 ($\times 3$), 128.2, 99.4 (C1^B), 86.7 (C1^A), 74.6 (C4^A), 73.2
11 (C3^A), 70.7 (C2^A), 70.3 (C2^B), 69.6 (C3^B), 68.7 (C4^B), 66.1 (C5^A), 61.1 (C5^B), 37.9, 29.6, 28.1.
12
13
14
15
16
17
18
19
20
21
22
23
24 HRMS: m/z calcd for C₄₉H₄₄O₁₄SNa [M + Na]⁺ 911.2344, found 911.2340.
25
26
27
28

29 **Glycosylation with a Thiophenyl Donor (General Procedure B).** Crushed mol sieves (3Å)
30 were added to a 2-neck Schlenk flask, where the middle neck was fitted with a glass stopper and
31 the other with a septum. The flask was placed under vacuum, heated with a heatgun and then
32 subjected to an atm of N₂ and cooled to rt. The donor (1.0 equiv) dissolved in dry CH₂Cl₂ ($c =$
33 0.12 M) was added to the flask together with AgOTf (2.0 equiv.) dissolved in dry toluene ($c =$
34 0.36 M). Stirring of the reaction was initiated and the glass stopper was exchanged for a
35 thermometer, and the solution was cooled to –65 °C. *p*NO₂PhSCl (1.0 equiv) was dissolved in
36 dry CH₂Cl₂ ($c = 0.72$ M) and slowly added to avoid raising the temperature above –60 °C. The
37 mixture was left to stir for approximately 10 min or until complete activation. The acceptor (0.9
38 equiv.) was dissolved in dry CH₂Cl₂ ($c = 0.65$ M) and added quickly. The temperature was kept
39 between –55 °C and –50 °C until TLC analysis showed completion of the reaction. Afterwards
40 the mixture was allowed to warm to –15 °C over 10 min at which point Et₃N (3.0 equiv) was
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 added. The solution was filtered through a pad of Celite and concd in vacuo. The residue was
4
5 then purified by flash column chromatography to afford the product.
6
7

8
9
10 *Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-2-O-benzoyl-3-O-levulinoyl-β-D-*
11
12 *xylopyranosyl-(1→4)-2,3-di-O-benzoyl-1-thio-β-D-xylopyranoside (15)*. Crushed mol sieves
13
14 (3Å, 1.0 g) were added a 2-neck Schlenk flask, where the middle neck was fitted with a glass
15
16 stopper and the other with a septum. The flask was placed under vacuum, heated with a heatgun
17
18 and then subjected to an atm of N₂ and cooled to rt. Donor **5** (0.200 g, 0.36 mmol) dissolved in
19
20 dry CH₂Cl₂ (3.0 mL) and AgOTf (0.185 g, 0.72 mmol) dissolved in dry toluene (2.0 mL) were
21
22 added. Stirring of the reaction was initiated and the glass stopper was exchanged for a
23
24 thermometer, and the solution was cooled to –65 °C. *p*NO₂PhSCl (0.068 g, 0.36 mmol)
25
26 dissolved in dry CH₂Cl₂ (0.5 mL) was added dropwise followed by stirring for an additional 10
27
28 min. Acceptor **11** (0.144 g, 0.33 mmol) dissolved in dry CH₂Cl₂ (0.5 mL) was quickly added.
29
30 The reaction was stirred at a temperature between –55 °C and –50 °C for 15 min and then for 15
31
32 min at rt. The mixture was cooled to –65 °C. AgOTf (0.093 g, 0.36 mmol) dissolved in dry
33
34 toluene (1.0 mL) was added followed by stirring for an additional 20 min. *p*NO₂PhSCl (0.062 g,
35
36 0.33 mmol) dissolved in dry CH₂Cl₂ (0.4 mL) was added dropwise followed by stirring for 10
37
38 min. Acceptor **6** (0.132 g, 0.29 mmol) dissolved in dry CH₂Cl₂ (0.4 mL) was added quickly. The
39
40 reaction was stirred at a temperature between –55 °C and –50 °C for 15 min and then allowed to
41
42 warm to –15 °C over 15 min, at which point Et₃N (0.8 mL, 0.54 mmol) was added. The mixture
43
44 was filtered through a pad of Celite, concd in vacuo, and purified by flash column
45
46 chromatography (heptane/acetone 3:2). The product **15** was obtained as a white amorphous solid
47
48 (0.124 g, 28%). Alternatively, general procedure B was employed with crushed mol sieves (1.0
49
50
51
52
53
54
55
56
57
58
59
60

g), donor **14** (0.490 g, 0.551 mmol), AgOTf (0.283 g, 1.10 mmol), *p*NO₂PhSCl (0.105 g, 0.551 mmol) and acceptor **6** (0.223 g, 0.496 mmol). Reaction time 15 min then Et₃N (0.232 mL, 1.65 mmol). Purification by flash column chromatography gave product **15** (0.420 g, 69%). *R*_f 0.22 (heptane/acetone 3:2). [α]_D²⁵ -24.0 (*c* 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 8.01–7.90 (m, 12H), 7.60–7.25 (m, 23H), 5.65 (t, *J* = 6.6 Hz, 1H, H^{3C}), 5.58 (t, *J* = 7.9 Hz, 1H, H^{3A}), 5.31–5.20 (m, 3H, H^{2A}, H^{3B}, H^{4C}), 5.16 (dd, *J*_{2C,3C} = 6.4 Hz, *J*_{1C,2C} = 4.9 Hz, 1H, H^{2C}), 5.02 (dd, *J*_{2B,3B} = 8.2 Hz, *J*_{1B,2B} = 6.6 Hz, 1H, H^{2B}), 4.96 (d, *J*_{1A,2A} = 8.1 Hz, 1H, H^{1A}), 4.71 (d, *J*_{1C,2C} = 4.7 Hz, 1H, H^{1C}), 4.66 (d, *J*_{1B,2B} = 6.4 Hz, 1H, H^{1B}), 4.40 (dd, *J*_{5Ceq,5Cax} = 12.4 Hz, *J*_{5Ceq,4C} = 3.8 Hz, 1H, H^{5C}_{eq}), 4.13 (dd, *J*_{5Aeq,5Aax} = 12.0 Hz, *J*_{5Aeq,4A} = 4.7 Hz, 1H, H^{5A}_{eq}), 3.95 (td, *J* = 8.2 Hz, *J*_{5Aeq,4A} = 4.9 Hz, 1H, H^{4A}), 3.74–3.65 (m, 2H, H^{4B}, H^{5C}_{ax}), 3.51 (dd, *J*_{5Beq,5Bax} = 12.2 Hz, *J*_{5Beq,4B} = 4.8 Hz, 1H, H^{5B}_{eq}), 3.45 (dd, *J*_{5Aax,5Aeq} = 12.0 Hz, *J*_{5Aax,4A} = 8.6 Hz, 1H, H^{5A}_{ax}), 3.09 (dd, *J*_{5Bax,5Beq} = 12.2 Hz, *J*_{5Bax,4B} = 8.7 Hz, 1H, H^{5B}_{ax}), 2.64–2.37 (m, 4H, 1.97 (s, 3H). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 205.9, 171.8, 165.6, 165.4, 165.3, 165.3, 165.2, 165.0, 133.6, 133.5 (×2), 133.5, 133.4, 133.3, 132.8, 132.6 (×2), 130.1 (×3), 130.0 (×2), 129.9 (×2), 129.8 (×2), 129.6, 129.4, 129.4, 129.4, 129.3, 129.2, 129.1 (×2), 128.6 (×2), 128.6 (×3), 128.5 (×3), 128.5 (×2), 128.4 (×2), 128.2, 101.0 (C^{1B}), 99.1 (C^{1C}), 86.7 (C^{1A}), 75.6 (C^{4A}), 74.2 (C^{4B}), 73.1 (C^{3A}), 71.9 (C^{3B}), 71.6 (C^{2B}), 70.4 (C^{2A}), 70.1 (C^{2C}), 69.6 (C^{3C}), 68.7 (C^{4C}), 65.9 (C^{5A}), 62.2 (C^{5B}), 61.0 (C^{5C}), 37.8, 29.6, 28.0. HRMS: *m/z* calcd. for C₆₈H₆₀O₂₀SNa [M + Na]⁺ 1251.3291, found 1251.3308.

2,3,5-Tri-O-benzoyl-L-arabinofuranosyl N-phenyl-2,2,2-trifluoroacetimidate (4). AcCl (5.0 mL, 70.3 mmol) was slowly added to methanol (60.0 mL) cooled to 0 °C. The methanolic HCl solution was then added to a vigorously stirred suspension of L-arabinose (10.0 g, 66.8 mmol) in

1
2
3 methanol (200 mL). The reaction was left to stir at rt for 4.5 h, then the mixture was concd
4
5 under co-evaporation with CH₂Cl₂. The resulting residue was cooled to 0 °C and dissolved in
6
7 pyridine (60.0 mL). BzCl (62.0 mL, 534 mmol) was slowly added over 25 min and the reaction
8
9 was left to stir at rt overnight. H₂O (5.0 mL) was added to quench the reaction which was stirred
10
11 for an additional 5 min. The mixture was then diluted with CH₂Cl₂ (500 mL) and washed with
12
13 H₂O, 1 M HCl, and NaHCO₃, dried over MgSO₄, filtered, and concd in vacuo under co-
14
15 evaporation with toluene. The residue was crystallized from absolute EtOH with the addition of
16
17 pentane to promote the crystallization. The intermediate methyl 2,3,5-tri-*O*-benzoyl- α -L-
18
19 arabinofuranoside was obtained as white crystals (14.13 g, 44% over 2 steps). *R*_f 0.44
20
21 (hexane/EtOAc 2:1). Mp 95.3–97.7 °C (EtOH) (lit.³⁵ 106 °C). ¹H NMR (400 MHz, CDCl₃) δ
22
23 8.09–8.00 (m, 6H), 7.62–7.56 (m, 2H), 7.53–7.49 (m, 1H), 7.46 (t, *J* = 7.7 Hz, 2H), 7.40 (t, *J* =
24
25 7.8 Hz, 2H), 7.30 (t, *J* = 7.8 Hz, 2H), 5.59 (d, *J* = 5.2 Hz, 1H, H3), 5.52 (d, *J* = 1.3 Hz, 1H, H2),
26
27 5.18 (s, 1H, H1), 4.85 (dd, *J*_{5,5'} = 11.9 Hz, *J*_{4,5} = 3.4 Hz, 1H, H5), 4.70 (dd, *J*_{5,5'} = 11.9 Hz, *J*_{4,5'} =
28
29 4.8 Hz, 1H, H5'), 4.59–4.56 (m, 1H, H4), 3.50 (s, 3H, OMe). ¹³C{¹H} NMR (101 MHz, CDCl₃)
30
31 δ 166.4, 166.0, 165.6, 133.7, 133.6, 133.2, 130.1 (×2), 130.0 (×2), 129.9 (×2), 129.2, 129.2,
32
33 128.7 (×2), 128.6 (×2), 128.5 (×2), 107.0 (C1), 82.4 (C2), 81.0 (C4), 78.1 (C3), 63.9 (C5), 55.2
34
35 (OMe). NMR data are in accordance with literature values.³⁵ The above glycoside (1.92 g, 4.03
36
37 mmol) was dissolved in 80% AcOH (13.0 mL) and 33% HBr·AcOH (11.0 mL, 60.7 mmol) was
38
39 added under an atm of N₂. The solution was let stir at rt for 3.5 h. EtOAc was added and the
40
41 mixture washed with brine and H₂O, dried over MgSO₄, filtered, and concd in vacuo. The
42
43 residue was purified by flash column chromatography (hexane/EtOAc 7:1) to afford 2,3,5-tri-*O*-
44
45 benzoyl-L-arabinofuranose (α : β ratio 1:0.35) as a white foam (1.445 g, 77%). *R*_f 0.41
46
47 (hexane/EtOAc 2:1). ¹H NMR (400 MHz, CDCl₃) δ 8.13–8.03 (m, 8H), 7.66–7.32 (m, 12H),
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 5.91 (t, $J = 5.4$ Hz, 0.35H, H3 $_{\beta}$), 5.86 (d, $J = 4.4$ Hz, 0.34H, H1 $_{\beta}$), 5.71 (s, 1H, H1 $_{\alpha}$), 5.61–5.57
4
5 (m, 2.35H, H2 $_{\alpha}$, H3 $_{\alpha}$, H2 $_{\beta}$), 4.89–4.75 (m, 2.70H, H4 $_{\alpha}$, H5 $_{\alpha}$, H5 $_{\beta}$, H5' $_{\beta}$), 4.69 (dd, $J = 11.5, 5.2$
6
7 Hz, 1H, H5' $_{\alpha}$), 4.51–4.47 (m, 0.34H, H4 $_{\beta}$). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl $_3$) δ 166.6, 166.4,
8
9 166.1, 166.0, 165.9, 165.7, 134.0, 133.8, 133.7, 133.7, 133.3, 133.3, 132.2, 130.3, 130.1, 130.1,
10
11 130.0, 129.9, 129.9, 129.8, 129.1, 129.1, 128.7, 128.7, 128.6, 128.5, 128.5, 128.4, 101.2 (C1 $_{\alpha}$),
12
13 95.7 (C1 $_{\beta}$), 82.6 (C2 $_{\alpha}$), 81.6 (C4 $_{\alpha}$), 79.3 (C4 $_{\beta}$), 78.1 (C3 $_{\alpha}$), 77.8 (C2 $_{\beta}$), 76.7 (C3 $_{\beta}$), 65.9 (C5 $_{\beta}$),
14
15 64.1 (C5 $_{\alpha}$). NMR data are in accordance with literature values.^{19a} The above hemiacetal (0.500
16
17 g, 1.08 mmol) was dissolved in acetone (4.0 mL) followed by addition of Cs $_2$ CO $_3$ (0.704 g, 2.16
18
19 mmol) and 2,2,2-trifluoro-*N*-phenylacetimidoyl chloride (0.35 mL, 2.21 mmol). The mixture
20
21 was left to stir at rt for 2 h and then filtered through a pad of Celite, concd in vacuo, and purified
22
23 by flash column chromatography (hexane/EtOAc 10:1). The product **4** was obtained a yellow
24
25 foam. It was possible to separate some of the α isomer (0.281 g, 41%) from the mixture (α/β
26
27 ratio 1:2) (0.292 g, 43%) (overall α/β ratio 2:1). $R_f(\alpha)$ 0.57 and $R_f(\beta)$ 0.52 (hexane/EtOAc 3:1).
28
29 ^1H NMR (400 MHz, CDCl $_3$) δ 8.10–8.08 (m, 5H), 8.06–8.04 (m, 3H), 8.01 (d, $J = 7.5$ Hz, 1H),
30
31 7.66–7.38 (m, 11.5H), 7.32–7.26 (m, 3H), 7.17 (t, $J = 7.8$ Hz, 2H), 7.09 (t, $J = 7.5$ Hz, 0.5H),
32
33 7.03 (t, $J = 7.5$ Hz, 1H), 6.88 (d, $J = 7.8$ Hz, 1H), 6.83 (bs, 1H, H1 $_{\beta}$), 6.58 (d, $J = 7.7$ Hz, 2H),
34
35 6.02–5.99 (m, 1H, H3 $_{\beta}$), 5.93–5.91 (m, 1H, H2 $_{\beta}$), 5.80 (s, 0.5H, H2 $_{\alpha}$), 5.67–5.66 (m, 0.5H,
36
37 H3 $_{\alpha}$), 4.84–4.80 (m, 2H, H4 $_{\alpha}$, H5 $_{\alpha}$, H5 $_{\beta}$), 4.76–4.71 (m, 1.5H, H5' $_{\alpha}$, H5' $_{\beta}$), 4.67–4.63 (m, 1H,
38
39 H4 $_{\beta}$). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl $_3$) δ 166.3, 166.2, 165.9, 165.7, 165.6, 165.2, 143.5,
40
41 143.4, 134.0, 134.0 ($\times 2$), 133.9 ($\times 2$), 133.3 ($\times 2$), 130.1 ($\times 4$), 130.1 ($\times 4$), 130.0 ($\times 4$), 129.9 ($\times 4$),
42
43 129.7, 129.7, 129.0, 128.9 ($\times 2$), 128.8 ($\times 4$), 128.7, 128.5 ($\times 4$), 124.6, 124.3, 119.7, 119.3, 102.4
44
45 (C1 $_{\alpha}$), 97.0 (C1 $_{\beta}$), 84.4 (C4 $_{\alpha}$), 80.9 (C2 $_{\alpha}$), 80.7 (C4 $_{\beta}$), 77.4 (C3 $_{\alpha}$), 76.3 (C2 $_{\beta}$), 75.7 (C3 $_{\beta}$), 65.0
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(C5_β), 63.7 (C5_α). HRMS: *m/z* calcd for C₃₄H₂₆F₃NO₈Na [M + Na]⁺ 656.1503, found 656.1499.

NMR data are in accordance with literature values.¹⁹

Deprotection of Levulinoyl Groups (General Procedure C). The linear xylan saccharide (1.0 equiv) was dissolved in AcOH (*c* = 0.40 M) and pyridine (*c* = 0.20 M). A mixture of a 50% solution of hydrazine hydrate (10.0 equiv) in AcOH (*c* = 20 M) and pyridine (*c* = 10 M) was added. The reaction was left to stir at rt until TLC showed consumption of the starting material and formation of the product. The reaction was stopped by the addition of acetone (300 equiv) and left to stir at rt for 30 min. EtOAc was added and the mixture was washed with 1 M HCl, NaHCO₃, and H₂O, dried over Na₂SO₄, filtered, and *concd* in *vacuo*. The product was obtained in sufficient purity and did not need further purification.

Glycosylation with *N*-Phenyl-2,2,2-trifluoroacetimidate 4 (General Procedure D). The partially deprotected xylan (1.0 equiv) and donor 4 (4.0 equiv) were co-evaporated twice with toluene and left under high vacuum overnight. The mixture was dissolved in dry CH₂Cl₂ (20 mL/0.1 g deprotected xylan) and cooled to −40 °C. TMSOTf (0.1 equiv) was added from a freshly made stock solution of TMSOTf in dry CH₂Cl₂. The reaction was kept at −40 °C for the time indicated after which Et₃N (0.7 equiv) was added, and the mixture was *concd* in *vacuo*. The residue was purified by flash column chromatography to afford the desired product.

Deprotection of Anomeric Thiophenyl Group to Prepare 29 – 31 (General Procedure E).

The arabinoxylan saccharide (1.0 equiv) was dissolved in acetone/H₂O 9:1 (15 mL/mmol) and NBS (4.0 equiv) was added. The mixture was left to stir at rt until TLC analysis showed

1
2
3 completion of the reaction. EtOAc was added to the mixture, which was then washed with
4
5 NaHCO₃ and H₂O, dried over Na₂SO₄, filtered, and concd in vacuo. The residue was purified by
6
7 flash column chromatography.
8
9

10
11
12 **Deprotection of Benzoyl Groups (General Procedure F).** The arabinoxylan saccharide (1.0
13
14 equiv) was dissolved in CH₂Cl₂/MeOH 1:1 (*c* = 0.01 M). A 1 M NaOMe solution was added
15
16 until the solution had obtained a pH of 11. The reaction was stirred at rt until TLC analysis
17
18 (EtOAc/MeOH/H₂O/AcOH 6:3:0.8:0.2) showed completion of the reaction (about 1 – 3 h). The
19
20 mixture was diluted with water and washed with Et₂O and EtOAc and concd followed by
21
22 purification with Sep-Pak C18 Plus Short Cartridge (Waters) with a gradient of H₂O to
23
24 H₂O/MeOH 1:1.
25
26
27
28
29
30

31 **Deprotection of Anomeric Thiophenyl Group to Prepare 1 – 3 (General Procedure G).** The
32
33 partially deprotected arabinoxylan saccharide (1.0 equiv) was dissolved in MeCN/H₂O 5:1 (*c* =
34
35 0.02 M) followed by addition of 2,6-lutidine (1.5 equiv) and NBS (3.0 equiv). The mixture was
36
37 stirred at rt until TLC analysis (EtOAc/MeOH/H₂O/AcOH 6:3:0.8:0.2) showed completion of
38
39 the reaction. The mixture was concd in vacuo. The crude residue was dissolved in MeOH (0.5
40
41 mL) followed by addition of ice-cold Et₂O, and precipitation for 30 min in the freezer. The
42
43 mixture was then centrifuged and the supernatant removed. The precipitation process was
44
45 repeated three times in total. The resulting residue was purified by gel filtration (PD miniTrap
46
47 G-10 column).
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 *Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-2-O-benzoyl-3-O-levulinoyl-β-D-*
5
6 *xylopyranosyl-(1→4)-2-O-benzoyl-3-O-levulinoyl-1-thio-β-D-xylopyranoside (16)*. General
7
8 procedure B with crushed mol sieves (1.0 g), donor **14** (0.250 g, 0.28 mmol), AgOTf (0.145 g,
9
10 0.56 mmol), *p*NO₂PhSCl (0.053 g, 0.28 mmol) and acceptor **11** (0.113 g, 0.25 mmol). Reaction
11
12 time 15 min then Et₃N (0.12 mL, 0.84 mmol). Purification by flash column chromatography
13
14 (heptane/EtOAc 4:3) gave **16** as a white amorphous solid (0.274 g, 88%). *R*_f 0.37
15
16 (heptane/EtOAc 1:1). [α]²⁵_D -57.7 (*c* 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 8.05–7.95 (m,
17
18 10H), 7.63–7.46 (m, 9H), 7.43–7.35 (m, 8H), 7.29–7.25 (m, 3H), 5.73 (t, *J* = 6.6 Hz, 1H, H3^C),
19
20 5.37–5.26 (m, 4H, H3^A, H3^B, H2^C, H4^C), 5.13 (t, *J*_{1A,2A} = 8.3 Hz, 1H, H2^A), 5.05 (dd, *J*_{2B,3B} =
21
22 8.3 Hz, *J*_{1B,2B} = 6.5 Hz, 1H, H2^B), 4.96 (d, *J*_{1C,2C} = 4.7 Hz, 1H, H1^C), 4.86 (d, *J*_{1A,2A} = 8.4 Hz,
23
24 1H, H1^A), 4.67 (d, *J*_{1B,2B} = 6.4 Hz, 1H, H1^B), 4.48 (dd, *J*_{5Ceq,5Cax} = 12.4 Hz, *J*_{5Ceq,4C} = 3.9 Hz, 1H,
25
26 H5^C_{eq}), 4.11–4.04 (m, 2H, H5^A_{eq}, H5^B_{eq}), 3.98–3.97 (m, 1H, H4^B), 3.88–3.82 (m, 1H, H4^A), 3.77
27
28 (dd, *J*_{5Cax,5Ceq} = 12.4 Hz, *J*_{5Cax,4C} = 6.2 Hz, 1H, H5^C_{ax}), 3.42 (dd, *J*_{5Bax,5Beq} = 12.1 Hz, *J*_{5Bax,4B} = 8.4
29
30 Hz, 1H, H5^B_{ax}), 3.36 (dd, *J*_{5Aax,5Aeq} = 11.9 Hz, *J*_{5Aax,4A} = 9.0 Hz, 1H, H5^A_{ax}), 2.66–2.39 (m, 8H),
31
32 2.06 (s, 3H), 2.01 (s, 3H). ¹³C {¹H} NMR (101 MHz, CDCl₃) δ 206.0, 205.9, 171.8, 171.6,
33
34 165.6, 165.3, 165.3, 165.1 (×2), 133.5, 133.5, 133.4, 133.4, 132.6 (×2), 132.6, 130.1 (×2), 130.0
35
36 (×2), 130.0 (×2), 129.9 (×2), 129.9 (×2), 129.5, 129.4, 129.4, 129.2, 129.1, 129.0 (×2), 128.6
37
38 (×2), 128.5 (×4), 128.5 (×4), 128.1, 100.5 (C1^B), 99.2 (C1^C), 86.5 (C1^A), 74.9 (C4^A), 74.3 (C4^B),
39
40 73.0 (C3^A), 71.9 (C3^B), 71.4 (C2^B), 70.4, 70.2 (C2^A, C2^C), 69.7 (C3^C), 68.7 (C4^C), 66.1 (C5^A),
41
42 62.4 (C5^B), 61.1 (C5^C), 37.8, 37.8, 29.8, 29.6, 28.0, 28.0. HRMS: *m/z* calcd for C₆₆H₆₂O₂₁SNa
43
44 [M + Na]⁺ 1245.3396, found 1245.3379.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 *Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-2-O-benzoyl-3-O-levulinoyl-β-D-*
5
6 *xylopyranosyl-(1→4)-2-O-benzoyl-3-O-levulinoyl-β-D-xylopyranosyl-(1→4)-2-O-benzoyl-3-O-*
7
8 *levulinoyl-1-thio-β-D-xylopyranoside (17)*. General procedure B with crushed mol sieves (1.5 g),
9
10 donor **16** (0.794 g, 0.65 mmol), AgOTf (0.332 g, 1.29 mmol), *p*NO₂PhSCl (0.123 g, 0.65 mmol)
11
12 and acceptor **11** (0.262 g, 0.59 mmol). Reaction time 60 min then Et₃N (0.27 mL, 1.95 mmol).
13
14 Purification by flash column chromatography (heptane/acetone 2:1) gave **17** as a white
15
16 amorphous solid (0.670 g, 73%). *R*_f 0.34 (hexane/acetone 1:1). [α]²⁵_D -46.6 (*c* 1.0, CHCl₃). ¹H
17
18 NMR (400 MHz, CDCl₃) δ 8.01–7.92 (m, 12H), 7.60–7.31 (m, 20H), 7.25–7.21 (m, 3H), 5.70
19
20 (t, *J* = 6.5 Hz, 1H, H3^D), 5.32 (t, *J*_{2B,3B} = 8.2 Hz, 1H, H3^B), 5.28–5.24 (m, 3H, H3^A, H2^D, H4^D),
21
22 (t, *J* = 6.5 Hz, 1H, H3^D), 5.32 (t, *J*_{2B,3B} = 8.2 Hz, 1H, H3^B), 5.28–5.24 (m, 3H, H3^A, H2^D, H4^D),
23
24 5.20 (t, *J*_{2C,3C} = 8.0 Hz, 1H, H3^C), 5.08 (t, *J*_{1A,2A} = 8.3 Hz, 1H, H2^A), 5.04 (dd, *J*_{2B,3B} = 8.4 Hz,
25
26 *J*_{1B,2B} = 6.5 Hz, 1H, H2^B), 4.97 (dd, *J*_{2C,3C} = 8.3 Hz, *J*_{1C,2C} = 6.4 Hz, 1H, H2^C), 4.93 (d, *J*_{1D,2D} =
27
28 4.7 Hz, 1H, H1^D), 4.81 (d, *J*_{1A,2A} = 8.4 Hz, 1H, H1^A), 4.64 (d, *J*_{1B,2B} = 6.5 Hz, 1H, H1^B), 4.57 (d,
29
30 *J*_{1C,2C} = 6.3 Hz, 1H, H1^C), 4.46 (dd, *J*_{5Deq,5Dax} = 12.4 Hz, *J*_{5Deq,4D} = 3.9 Hz, 1H, H5^D_{eq}), 4.05–4.00
31
32 (m, 2H, H5^A, H5^B_{eq}), 3.95–3.89 (m, 2H, H4^B, H5^C), 3.80–3.72 (m, 3H, H4^A, H4^C, H5^D_{ax}), 3.38
33
34 (dd, *J*_{5Bax,5Beq} = 12.1 Hz, *J*_{5Bax,4B} = 8.5 Hz, 1H, H5^B_{ax}), 3.33–3.25 (m, 2H, H5^A, H5^C),
35
36 2.66–2.32 (m, 12H), 2.03 (s, 3H), 2.01 (s, 3H), 1.97 (s, 3H). ¹³C{¹H} NMR (101 MHz, CDCl₃)
37
38 δ 206.1, 206.0, 205.9, 171.8, 171.7, 171.5, 165.6, 165.3, 165.2, 165.2, 165.1, 165.1, 133.5,
39
40 133.5 (×2), 133.4 (×2), 132.6 (×2), 132.5, 130.1 (×2), 130.0 (×2), 130.0 (×2), 129.9 (×3), 129.9
41
42 (×3), 129.5, 129.4 (×2), 129.4, 129.2, 129.1, 129.0 (×2), 128.6 (×3), 128.5 (×4), 128.5 (×4),
43
44 128.1, 100.4 (×2, C1^B, C1^C), 99.2 (C1^D), 86.5 (C1^A), 74.7 (C4^A), 74.5, 74.3 (C4^B, C4^C), 72.9
45
46 (C3^A), 71.9 (C3^B), 71.6 (C3^C), 71.4 (C2^B), 71.1 (C2^C), 70.4 (C2^A), 70.2 (C2^D), 69.6 (C3^D), 68.7
47
48 (C4^D), 66.0 (C5^A), 62.4 (2C, C5^B, C5^C), 61.0 (C5^D), 37.8, 37.8, 37.8, 29.8, 29.8, 29.6, 28.0,
49
50 28.0, 27.9. HRMS: *m/z* calcd for C₈₃H₈₀O₂₈SNa [M + Na]⁺ 1579.4449, found 1579.4427.
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6 *Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-2-O-benzoyl-β-D-xylopyranosyl-(1→4)-2-*
7
8 *O-benzoyl-β-D-xylopyranosyl-(1→4)-2-O-benzoyl-1-thio-β-D-xylopyranoside (18)*. General
9
10 procedure C with tetrasaccharide **17** (0.146 g, 0.094 mmol) and 50% solution of hydrazine
11 hydrate (59 μL, 0.94 mmol). Reaction time 3 h. Eluent for TLC (heptane/EtOAc 1:1). The
12 product obtained was a white powder (0.113 g, 96%). R_f 0.37 (heptane/acetone 1:1). $[\alpha]^{25}_D$
13 -36.6 (c 1.0, CHCl_3). $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.06–8.00 (m, 6H), 7.96–7.90 (m, 6H),
14 7.62–7.32 (m, 22H), 7.24–7.20 (m, 1H), 5.82 (t, $J = 8.9$ Hz, 1H, H3^D), 5.42 (dd, $J_{2D,3D} = 9.0$ Hz,
15 $J_{1D,2D} = 7.0$ Hz, 1H, H1^D), 5.37 (td, $J_{3D,4D} = 8.9$ Hz, $J = 4.5$ Hz, 1H, H4^D), 5.11–5.07 (m, 1H,
16 H2^C), 5.05–5.00 (m, 2H, H2^A, H2^B), 4.83 (d, $J_{1D,2D} = 7.0$ Hz, 1H, H1^D), 4.64 (d, $J_{1A,2A} = 9.9$ Hz,
17 1H, H1^A), 4.50 (d, $J_{1C,2C} = 7.9$ Hz, 1H, H1^C), 4.47–4.43 (m, 2H, H1^B, H5^D), 3.87–3.77 (m, 4H,
18 H3^A, H3^C, H4^C, H5^C), 3.75–3.55 (m, 7H, H3^A, H4^A, H3^B, H4^B, H5^B, H5^D), 3.31–3.15 (m, 3H,
19 H5^A, H5^B, H5^C). $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 165.6 (×2), 165.5, 165.5, 165.4, 165.2,
20 133.8, 133.7, 133.6, 133.5, 133.5, 133.3, 132.8 (×3), 132.3, 130.0 (×2), 129.9 (×2), 129.9 (×2),
21 129.9 (×2), 129.8 (×2), 129.7 (×4), 129.5, 129.4, 129.1, 129.0, 129.0 (×3), 128.9, 128.9, 128.9,
22 128.8 (×2), 128.7 (×2), 128.6 (×2), 128.6 (×2), 128.5 (×2), 128.5 (×2), 128.3, 128.1, 102.2,
23 102.1 (C1^B, C1^C, $J_{\text{C-H}} = 157$ Hz, $J_{\text{C-H}} = 163$ Hz), 101.8 (C1^D, $J_{\text{C-H}} = 162$ Hz), 86.6 (C1^A, $J_{\text{C-H}} =$
24 155 Hz), 80.6 (C4^A), 80.3 (C4^B), 80.1 (C4^C), 74.8 (C3^A), 73.2, 73.1, 73.0, 73.0 (C2^B, C3^B, C2^C,
25 C3^C), 72.0 (C2^A), 71.3 (C3^D), 71.1 (C2^D), 69.3 (C4^D), 67.1 (C5^A), 63.5 (2C, C5^B, C5^C), 62.8
26 (C5^D). HRMS: m/z calcd for $\text{C}_{68}\text{H}_{62}\text{O}_{22}\text{SNa}$ $[\text{M} + \text{Na}]^+$ 1285.3346, found 1285.3346.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 *Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-[2,3,5-tri-O-benzoyl-α-L-*
53 *arabinofuranosyl-(1→3)]-2-O-benzoyl-β-D-xylopyranosyl-(1→4)-[2,3,5-tri-O-benzoyl-α-L-*
54
55
56
57
58
59
60

1
2
3
4 *arabinofuranosyl-(1→3)]-2-O-benzoyl-β-D-xylopyranosyl-(1→4)-[2,3,5-tri-O-benzoyl-α-L-*
5
6 *arabinofuranosyl-(1→3)]-2-O-benzoyl-1-thio-β-D-xylopyranoside (19)*. General procedure D
7
8 with acceptor **18** (0.054 g, 0.043 mmol), donor **4** (0.109 g, 0.17 mmol) and TMSOTf in dry
9
10 CH₂Cl₂ (0.11 mL, *c* = 0.04 mmol/mL). Reaction time 75 min then Et₃N (0.02 mL, 0.11 mmol).
11
12 Eluent TLC (heptane/acetone 1:1) and flash column chromatography (heptane/acetone 2:1). The
13
14 product **19** was isolated as a white amorphous solid (0.100 g, 90%). *R*_f 0.41 (heptane/acetone
15
16 1:1). [α]²⁵_D -46.9 (*c* 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 8.18–8.16 (m, 2H), 8.09–7.87
17
18 (m, 28H), 7.83 (t, *J* = 7.4 Hz, 1H), 7.64–7.16 (m, 49H), 5.71 (t, *J*_{3D,4D} = 9.3 Hz, 1H, H^{3D}), 5.52
19
20 (d, *J* = 5.5 Hz, 1H), 5.49 (d, *J* = 5.3 Hz, 1H), 5.45 (d, *J* = 4.6 Hz, 1H), 5.43 (s, 1H), 5.37 (d, *J* =
21
22 1.0 Hz, 1H), 5.35–5.34 (m, 2H), 5.30–5.28 (m, 2H, H^{2D}), 5.26–5.24 (m, 2H, H^{2A}), 5.15 (td,
23
24 *J*_{3D,4D} = 9.3 Hz, *J*_{4D,5D} = 4.7 Hz, 1H, H^{4D}), 5.07–4.83 (m, 2H, H^{2B}, H^{2C}), 4.68 (d, *J* = 8.8 Hz,
25
26 1H, H^{1A}), 4.23 (d, *J* = 8.0 Hz, 1H, H^{1B}), 4.10 (d, *J* = 7.4 Hz, 1H, H^{1D}), 4.08–3.99 (m, 2H, H^{3A},
27
28 H^{5A}), 3.88–3.82 (m, 3H, H^{3B}, H^{3C}, H^{5D}), 3.76 (td, *J* = 8.8, 4.8 Hz, 1H, H^{4A}), 3.59–3.53 (m, 2H,
29
30 H^{1C}, H^{4C}), 3.33 (dd, *J* = 11.8, 5.1 Hz, 1H, H^{5C}), 3.29–3.19 (m, 2H, H^{4B}, H^{5'D}), 3.10–3.02 (m,
31
32 2H, H^{5'A}, H^{5B}), 2.67–2.56 (m, 2H, H^{5'B}, H^{5'C}). ¹³C {¹H} NMR (101 MHz, CDCl₃) δ 166.4,
33
34 166.4, 166.4, 166.1, 165.9, 165.8, 165.7, 165.1, 165.1 (×3), 165.0, 164.7, 164.4, 164.2, 133.9,
35
36 133.7, 133.5, 133.4, 133.3, 133.2, 133.2, 133.1, 133.0, 132.9, 132.5, 130.4, 130.2, 130.2, 130.1,
37
38 130.1, 130.0, 129.9, 129.9, 129.9, 129.9, 129.8, 129.8, 129.7, 129.7, 129.7, 129.6, 129.5, 129.3,
39
40 129.2, 129.1, 129.1, 129.0, 129.0, 129.0, 129.0, 128.9, 128.9, 128.6, 128.6, 128.5, 128.5, 128.4,
41
42 128.4, 128.3, 128.3, 128.2, 127.9, 106.1, 105.7, 105.6 (C^{1E}, C^{1F}, C^{1G}), 100.3 (C^{1B}), 100.1
43
44 (C^{1C}), 99.9 (C^{1B}), 86.7 (C^{1A}), 82.8, 82.7, 82.5, 82.1, 81.5, 80.8, 78.3, 78.2 (×2), 76.0 (C^{3A}),
45
46 75.3 (C^{3B}/C^{3C}), 75.2 (C^{3B}/C^{3C}), 74.5 (C^{4C}), 74.2 (C^{4B}), 73.7 (C^{4A}), 73.3 (C^{2B}), 73.0 (C^{2C}),
47
48 72.2 (C^{2A}/C^{3D}), 72.0 (C^{2A}/C^{3D}), 71.2 (C^{2D}), 69.8 (C^{4D}), 66.1 (C^{5A}), 64.0, 63.9, 63.8 (C^{5E}, C^{5F},

C5^G), 63.1, 63.1, 63.0 (C5^B, C5^C, C5^D). HRMS: m/z calcd for C₁₄₆H₁₂₂O₄₃SNa [M + Na]⁺ 2617.6973, found 2617.6976.

Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-2-O-benzoyl-3-O-levulinoyl-β-D-xylopyranosyl-(1→4)-2,3-di-O-levulinoyl-1-thio-β-D-xylopyranoside (20). General procedure B with crushed mol sieves (2.6 g), donor **14** (1.00 g, 1.12 mmol), AgOTf (0.579 g, 2.25 mmol) and *p*NO₂PhSCl (0.214 g, 1.12 mmol). Activation time 40 min then acceptor **9** (0.447 g, 1.02 mmol). Reaction time 50 min then Et₃N (0.47 mL, 3.37 mmol). Purification by flash column chromatography (hexane/toluene/EtOAc 3:1:4) gave **20** as a white amorphous solid (0.883 g, 71%). In addition, donor **14** was reisolated (0.119 g, 12%). *R*_f 0.48 (hexane/acetone 3:2). [α]_D²⁵ -60.8 (*c* 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 8.02–7.92 (m, 8H), 7.61–7.43 (m, 6H), 7.41–7.32 (m, 8H), 7.28–7.26 (m, 3H), 5.70 (t, *J* = 6.5 Hz, 1H, H3^C), 5.31 (t, *J* = 8.1 Hz, 1H, H3^B), 5.28–5.24 (m, 2H, H2^C, H4^C), 5.11 (t, *J*_{2A,3A} = 8.7 Hz, 1H, H3^A), 5.00 (dd, *J*_{2B,3B} = 8.3 Hz, *J*_{1B,2B} = 6.4 Hz, 1H, H2^B), 4.97 (d, *J*_{1C,2C} = 4.7 Hz, 1H, H1^C), 4.84 (t, *J*_{2A,3A} = 9.0 Hz, 1H, H2^A), 4.61–4.58 (m, 2H, H1^A, H1^B), 4.46 (dd, *J*_{5Ceq,5Cax} = 12.4 Hz, *J*_{5Ceq,4C} = 3.8 Hz, 1H, H5^C_{eq}), 4.06 (dd, *J*_{5Beq,5Bax} = 11.9 Hz, *J*_{5Beq,4B} = 4.8 Hz, 1H, H5^B_{eq}), 4.02–3.98 (m, 1H, H4^B), 3.94 (dd, *J*_{5Aeq,5Aax} = 11.9 Hz, *J*_{5Aeq,4A} = 5.2 Hz, 1H, H5^A_{eq}), 3.77–3.71 (m, 2H, H4^A, H5^C_{ax}), 3.38 (dd, *J*_{5Bax,5Beq} = 11.9 Hz, *J*_{5Bax,4B} = 8.1 Hz, 1H, H5^B_{ax}), 3.21 (dd, *J*_{5Aax,5Aeq} = 11.8 Hz, *J*_{5Aax,4A} = 9.8 Hz, 1H, H5^A_{ax}), 2.85–2.39 (m, 12H), 2.17 (s, 3H), 2.16 (s, 3H), 1.98 (s, 3H). ¹³C {¹H} NMR (101 MHz, CDCl₃) δ 206.5, 206.4, 206.0, 171.9, 171.7, 171.5, 165.6, 165.4, 165.2, 165.1, 133.5, 133.5, 133.5, 132.8 (×2), 132.3, 130.1 (×2), 130.0 (×2), 130.0 (×2), 129.9 (×2), 129.4, 129.4, 129.3, 129.2, 129.1 (×2), 128.7 (×2), 128.5 (×4), 128.2, 100.6 (C1^B), 99.3 (C1^C), 86.5 (C1^A), 75.2 (C4^A), 74.3 (C4^B), 73.4 (C3^A), 71.9 (C3^B), 71.4 (C2^B), 70.2, 70.0 (C2^A, C2^C), 69.7

(C3^C), 68.8 (C4^C), 66.6 (C5^A), 62.5 (C5^B), 61.1 (C5^C), 37.9, 37.9, 37.8, 30.0, 29.9, 29.7, 28.1, 28.0, 28.0. HRMS: m/z calcd for C₆₄H₆₄O₂₂SNa [M + Na]⁺ 1239.3502, found 1239.3489.

Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-2-O-benzoyl-3-O-levulinoyl-β-D-xylopyranosyl-(1→4)-2,3-di-O-levulinoyl-β-D-xylopyranosyl-(1→4)-2,3-di-O-benzoyl-1-thio-β-D-xylopyranoside (21). General procedure B with crushed mol sieves (2.1 g), donor **20** (1.21 g, 0.99 mmol), AgOTf (0.510 g, 1.98 mmol), *p*NO₂PhSCl (0.187 g, 0.99 mmol) and acceptor **6** (0.402 g, 0.89 mmol). Reaction time 40 min then Et₃N (0.40 mL, 2.97 mmol). Purification by flash column chromatography (heptane/toluene/acetone 3:2:2) gave **21** as a white amorphous solid (0.851 g, 61%). R_f 0.40 (hexane/acetone 3:2). $[\alpha]_D^{25}$ -38.2 (*c* 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 8.01–7.91 (m, 12H), 7.61 (t, $J = 7.4$ Hz, 1H), 7.56–7.45 (m, 9H), 7.41–7.32 (m, 10H), 7.29–7.24 (m, 2H), 7.18–7.16 (m, 1H), 5.70 (t, $J = 6.5$ Hz, 1H, H3^D), 5.53 (t, $J = 7.6$ Hz, 1H, H3^A), 5.30–5.24 (m, 4H, H2^A, H3^C, H2^D, H4^D), 5.03 (d, $J_{1A,2A} = 7.7$ Hz, 1H, H1^A), 4.99–4.91 (m, 3H, H3^B, H2^C, H1^D), 4.70 (dd, $J_{2B,3B} = 8.9$ Hz, $J_{1B,2B} = 7.0$ Hz, 1H, H2^B), 4.45 (dd, $J_{5Deq,5Dax} = 12.4$ Hz, $J_{5Deq,4D} = 3.8$ Hz, 1H, H5^D_{eq}), 4.41 (d, $J_{1B,2B} = 6.9$ Hz, 1H, H1^B), 4.36 (d, $J_{1C,2C} = 6.5$ Hz, 1H, H1^C), 4.30 (dd, $J_{5Aeq,5Aax} = 12.1$ Hz, $J_{5Aeq,4A} = 4.6$ Hz, 1H, H5^A_{eq}), 4.01–3.86 (m, 3H, H4^A, H4^C, H5^C), 3.74 (dd, $J_{5Dax,5Deq} = 12.4$ Hz, $J_{5Dax,4D} = 6.1$ Hz, 1H, H5^D_{ax}), 3.58 (dd, $J_{5Aax,5Aeq} = 12.1$ Hz, $J_{5Aax,4A} = 8.2$ Hz, 1H, H5^A_{ax}), 3.49–3.44 (m, 1H, H4^B), 3.34–3.28 (m, 2H, H5^B_{eq}, H5^C), 2.89 (dd, $J_{5Bax,5Beq} = 12.0$ Hz, $J_{5Bax,4B} = 9.4$ Hz, 1H, H5^B_{ax}), 2.71–2.35 (m, 12H), 2.14 (bs, 6H), 1.97 (s, 3H). ¹³C {¹H} NMR (101 MHz, CDCl₃) δ 206.4, 206.4, 205.9, 171.8, 171.7, 171.3, 165.6, 165.4, 165.3 (×2), 165.2, 165.0, 133.5, 133.5, 133.5, 133.5, 133.3, 133.2, 133.0, 132.6 (×2), 130.1 (×3), 130.0 (×2), 130.0 (×2), 129.9 (×2), 129.8 (×2), 129.6, 129.5, 129.4, 129.4, 129.2, 129.2, 129.1, 129.1 (×2), 128.6 (×2), 128.6 (×3), 128.5 (×4), 128.5

($\times 2$), 128.4 ($\times 2$), 128.3, 128.1, 100.9 (C1^B), 100.3 (C1^C), 99.2 (C1^D), 86.6 (C1^A), 75.4 (C4^A), 74.8 (C4^B), 74.4 (C4^C), 72.7 (C3^A), 72.1, 71.9 (C3^B, C2^A/C3^C/C2^D), 71.3, 71.2 (C2^B, C2^C), 70.4, 70.2 (C2^A/C3^C/C2^D), 69.6 (C3^D), 68.7 (C4^D), 65.5 (C5^A), 62.6, 62.5 (C5^B, C5^C), 61.1 (C5^D), 37.8, 37.8, 37.7, 30.0, 29.9, 29.6, 28.0, 27.9, 27.9. HRMS: m/z calcd for C₈₃H₈₀O₂₈SNa [M + Na]⁺ 1579.4449, found 1579.4430.

Phenyl 2,3,4-tri-O-benzoyl- β -D-xylopyranosyl-(1 \rightarrow 4)-2-O-benzoyl- β -D-xylopyranosyl-(1 \rightarrow 4)- β -D-xylopyranosyl-(1 \rightarrow 4)-2,3-di-O-benzoyl-1-thio- β -D-xylopyranoside (22). General procedure C with tetrasaccharide **21** (0.507 g, 0.33 mmol) and 50% solution of hydrazine hydrate (0.20 mL, 3.21 mmol). Reaction time 20 min. Eluent for TLC (hexane/toluene/acetone 2:1:2). The product was a white powder (0.400 g, 97%). R_f 0.24 (hexane/acetone 3:2). $[\alpha]_D^{25}$ -15.4 (c 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 8.04–7.92 (m, 12H), 7.63 (t, J = 7.4 Hz, 1H), 7.57–7.26 (m, 22H), 5.84 (t, $J_{2D,3D}$ = 8.8 Hz, 1H, H3^D), 5.57 (t, $J_{2A,3A}$ = 7.5 Hz, 1H, H3^A), 5.43 (dd, $J_{2D,3D}$ = 8.7 Hz, $J_{1D,2D}$ = 7.1 Hz, 1H, H2^D), 5.38–5.30 (m, 2H, H2^A, H4^D), 5.10 (d, $J_{1A,2A}$ = 7.4 Hz, 1H, H1^A), 5.05 (t, J = 8.2 Hz, 1H, H2^C), 4.88 (d, $J_{1D,2D}$ = 6.9 Hz, 1H, H1^D), 4.42–4.36 (m, 2H, H5^A, H5^D), 4.33 (d, J = 7.6 Hz, 1H, H1^C), 4.27 (d, J = 6.7 Hz, 1H, H1^B), 3.97–3.92 (m, 1H, H4^A), 3.89–3.81 (m, 3H, H3^C, H4^C, H5^C), 3.73–3.62 (m, 4H, H5^{'A}, H5^{'D}, OH), 3.45 (t, J = 8.1 Hz, 1H, H3^B), 3.33–3.19 (m, 4H, H2^B, H4^B, H5^B_{eq}, H5^{'C}), 3.05 (bs, 1H, OH) 2.89 (dd, $J_{5Bax,5Beq}$ = 11.5 Hz, $J_{5bax,4B}$ = 9.2 Hz, 1H, H5^B_{ax}). ¹³C {¹H} NMR (101 MHz, CDCl₃) δ 165.6, 165.6, 165.6, 165.6, 165.3, 165.2, 133.8, 133.7, 133.5 ($\times 2$), 133.4, 133.3, 133.0, 132.6 ($\times 3$), 130.0 ($\times 4$), 129.9 ($\times 4$), 129.9 ($\times 2$), 129.8 ($\times 2$), 129.6, 129.5, 129.5, 129.1 ($\times 2$), 129.0, 129.0, 128.9, 128.7 ($\times 2$), 128.7 ($\times 2$), 128.6 ($\times 2$), 128.5 ($\times 2$), 128.5 ($\times 2$), 128.4 ($\times 2$), 128.2, 102.1 (C1^B, J_{C-H} = 160 Hz), 101.8 (C1^D, J_{C-H} = 163 Hz), 101.4 (C1^C, J_{C-H} = 163 Hz), 86.6 (C1^A, J_{C-H} = 159 Hz), 80.2 (C4^C), 79.0

(C4^B), 74.1 (C4^A), 73.3, 73.1, 73.0 (C3^B, C2^C, C3^C), 72.5, 72.3 (C3^A, C2^B), 71.4, 71.2 (C2^D, C3^D), 70.2 (C2^A), 69.4 (C4^D), 65.1 (C5^a), 63.4 (C5^C), 62.8 (C5^D), 62.0 (C5^B). HRMS: m/z calcd for C₆₈H₆₂O₂₂SNa [M + Na]⁺ 1285.3345, found 1285.3334.

Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-[2,3,5-tri-O-benzoyl-α-L-arabinofuranosyl-(1→3)]-2-O-benzoyl-β-D-xylopyranosyl-(1→4)-[(2,3,5-tri-O-benzoyl-α-L-arabinofuranosyl-(1→2))-[2,3,5-tri-O-benzoyl-α-L-arabinofuranosyl-(1→3)]-β-D-xylopyranosyl-(1→4)-2,3-di-O-benzoyl-1-thio-β-D-xylopyranoside (23). General procedure D with acceptor **22** (0.201 g, 0.16 mmol), donor **4** (0.401 g, 0.63 mmol) and TMSOTf in dry CH₂Cl₂ (0.16 mL, $c = 0.10$ mmol/mL). Reaction time 1 h then Et₃N (0.02 mL, 0.11 mmol). Eluent TLC analysis (hexane/toluene/acetone 2:1:1) and eluent flash column chromatography (hexane/toluene/acetone 3:2:1→2:1:1). The product was a white amorphous solid (0.393 g, 95%). R_f 0.42 (hexane/toluene/acetone 2:1:1). $[\alpha]^{25}_D -52.5$ (c 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 8.18 (d, $J = 7.3$ Hz, 2H), 8.09–7.73 (m, 28H), 7.65 (d, $J = 7.3$ Hz, 2H), 7.58–7.05 (m, 48H), 5.76 (s, 1H, H1^F), 5.69 (t, $J = 9.1$ Hz, 1H, H3^D), 5.65 (s, 1H, H1^E), 5.54–5.47 (m, 6H, H3^A), 5.34–5.34 (m, 1H, H2^E), 5.25 (s, 1H, H1^G), 5.25–5.20 (m, 2H, H2^A, H2^D), 5.19–5.13 (m, 1H, H4^D), 5.10–5.06 (m, 2H), 5.04–4.96 (m, 3H, H2^C), 4.94–4.85 (m, 3H, H1^A), 4.79 (dd, $J = 11.4, 3.0$ Hz, 1H), 4.71–4.62 (m, 2H), 4.31 (d, $J_{1B,2B} = 6.9$ Hz, 1H, H1^B), 4.17–4.13 (m, 2H, H5^A, H1^D), 3.88–3.81 (m, 4H, H3^B, H1^C, H3^C, H5^D), 3.79–3.74 (m, 1H, H4^A), 3.65 (dd, $J_{2B,3B} = 9.5$ Hz, $J_{1B,2B} = 7.0$ Hz, 1H, H2^B), 3.53–3.47 (m, 2H, H5^{'A}, H4^C), 3.32–3.26 (m, 3H, H4^B, H5^B, H5^{'D}), 3.03 (dd, $J_{5C,5C'} = 11.8$ Hz, $J_{5C,4C} = 5.1$ Hz, 1H, H5^C), 2.73–2.67 (m, 1H, H5^{'B}), 2.53 (t, $J = 11.1$ Hz, 1H, H5^{'C}). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 166.5, 166.3, 166.3, 166.0, 165.8, 165.8, 165.7, 165.6, 165.3, 165.2 (×2), 165.0, 165.0, 164.5, 164.2, 133.6, 133.6, 133.6, 133.5,

1
2
3 133.4, 133.4, 133.3, 133.2, 133.1, 133.0, 132.9, 132.8, 132.2, 130.3, 130.1, 130.1, 130.0, 129.9,
4
5 129.9, 129.8, 129.8, 129.7, 129.7, 129.5, 129.2, 129.1, 129.1, 129.0, 129.0, 129.0, 128.9, 128.9,
6
7 128.6, 128.6, 128.4, 128.4, 128.3, 128.3, 128.2, 127.9, 106.9 (C1^E), 106.2 (C1^G), 105.4 (C1^F),
8
9 101.7 (C1^B), 99.8 (C1^D), 99.5 (C1^C), 86.2 (C1^A), 82.6 (C2^F), 82.3, 82.2, 81.8, 81.8, 80.7, 79.7
10
11 (C2^B), 78.6, 78.1, 77.8, 76.0 (C3^B/C3^C), 75.8 (C3^B/C3^C), 74.6 (C4^C), 73.8 (C4^B), 73.6 (C4^A),
12
13 73.1 (C2^C), 71.9 (C3^D), 71.8 (C3^A), 71.1 (C2^D), 70.2 (C2^A), 69.7 (C4^D), 64.6 (C5^A), 64.2, 64.0,
14
15 63.9 (C5^E, C5^F, C5^G), 63.0 (C5^C/C5^D), 62.8 (C5^C/C5^D), 62.3 (C5^B). HRMS: *m/z* calcd for
16
17 C₁₄₆H₁₂₂O₄₃S(Na⁺)₂ [M + 2Na]²⁺ 1320.3432; found *m/z* 1320.3463.
18
19
20
21
22
23

24 *Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-2,3-di-O-levulinoyl-1-thio-β-D-*
25
26 *xylopyranoside (24)*. General procedure B with crushed mol sieves (2.0 g), donor **5** (0.402 g,
27 0.72 mmol), AgOTf (0.378 g, 1.47 mmol), *p*NO₂PhSCl (0.141 g, 0.74 mmol) and acceptor **9**
28 (0.286 g, 0.65 mmol). Reaction time 25 min then Et₃N (0.30 mL, 2.15 mmol). Purification by
29 flash column chromatography (pentane/EtOAc 3:2) gave **24** as a white amorphous solid (0.547
30 g, 95%). *R*_f 0.42 (hexane/EtOAc 1:1). [α]_D²⁵ -28.1 (*c* 0.5, CHCl₃). ¹H NMR (400 MHz, CDCl₃)
31 δ 8.02 (d, *J* = 7.2 Hz, 2H), 7.94 (d, *J* = 7.4 Hz, 4H), 7.59–7.49 (m, 3H), 7.45–7.28 (m, 11H),
32 5.69 (t, *J* = 6.4 Hz, 1H, H3^B), 5.30 (td, *J* = 6.1 Hz, *J*_{5Beq,4B} = 4.0 Hz, 1H, H4^B), 5.25 (d, *J* = 8.8
33 Hz, 1H, H3^A), 5.22–5.19 (m, 1H, H2^B), 4.92–4.87 (m, 2H, H2^A, H1^B), 4.66 (d, *J*_{A1,A2} = 9.3 Hz,
34 1H, H1^A), 4.49 (dd, *J*_{5Beq,5Bax} = 12.4 Hz, *J*_{5Beq,4B} = 3.8 Hz, 1H, H5^B_{eq}), 4.09 (dd, *J*_{5Aeq,5Aax} = 11.9
35 Hz, *J*_{5Aeq,4A} = 5.2 Hz, 1H, H5^A_{eq}), 3.90 (td, *J* = 9.4 Hz, *J*_{5Aeq,4A} = 5.2 Hz, 1H, H4^A), 3.75 (dd,
36 *J*_{5Bax,5Beq} = 12.4 Hz, *J*_{5Bax,4B} = 6.0 Hz, 1H, H5^B_{ax}), 3.32 (dd, *J*_{5Aax,5Aeq} = 11.7 Hz, *J*_{5Aeq,4A} = 9.9 Hz,
37 1H, H5^A_{ax}), 2.88–2.53 (m, 8H), 2.19 (s, 3H), 2.08 (s, 3H). ¹³C {¹H} NMR (101 MHz, CDCl₃) δ
38 206.5, 206.4, 171.9, 171.5, 165.7, 165.4, 165.2, 133.6, 133.5, 133.5, 132.9 (×2), 132.2, 130.1
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

($\times 2$), 130.1 ($\times 2$), 129.9 ($\times 2$), 129.4, 129.2, 129.1, 129.1 ($\times 2$), 128.6 ($\times 4$), 128.3, 99.4 (C1^B), 86.6 (C1^A), 74.9 (C4^A), 73.5 (C3^A), 70.3 (C2^A), 70.2 (C2^B), 69.6 (C3^B), 68.7 (C4^B), 66.7 (C5^A), 61.1 (C5^B), 37.9, 37.8, 30.0, 29.8, 28.1, 28.0. HRMS: m/z calcd for C₄₇H₄₆O₁₅SNa [M + Na]⁺ 905.2449, found 905.2444.

Phenyl 2,3,4-tri-O-benzoyl- β -D-xylopyranosyl-(1 \rightarrow 4)-2,3-di-O-levulinoyl- β -D-xylopyranosyl-(1 \rightarrow 4)-2-O-benzoyl-3-O-levulinoyl-1-thio- β -D-xylopyranoside (25). General procedure B with crushed mol sieves (1.2 g), donor **24** (0.252 g, 0.29 mmol), AgOTf (0.148 g, 0.58 mmol), *p*NO₂PhSCl (0.054 g, 0.28 mmol) and acceptor **11** (0.114 g, 0.26 mmol). Reaction time 25 min then Et₃N (0.12 mL, 0.86 mmol). Purification by flash column chromatography (heptane/toluene/EtOAc 2:1:1) gave **25** as a light yellow amorphous solid (0.162 g, 52%). R_f 0.35 (hexane/EtOAc 1:1). $[\alpha]_D^{25}$ -53.8 (c 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 8.03–7.99 (m, 6H), 7.96–7.93 (m, 4H), 7.60–7.50 (m, 5H), 7.47–7.32 (m, 12H), 7.28–7.26 (m, 3H), 5.69 (t, $J = 6.6$ Hz, 1H, H3^C) 5.30–5.25 (m, 2H, H3^A, H4^C), 5.22 (dd, $J_{2C,3C} = 6.5$ Hz, $J_{1C,2C} = 4.8$ Hz, 1H, H2^C), 5.17–5.10 (m, 2H, H2^A, H3^B), 4.93 (d, $J_{1A,2A} = 8.1$ Hz, 1H, H1^A), 4.87 (d, $J_{1C,2C} = 4.7$ Hz, 1H, H1^C), 4.77 (dd, $J = 8.9, 7.0$ Hz, 1H, H2^B), 4.48–4.44 (m, 2H, H1^B, H5^C), 4.22 (dd, $J_{5Aeq,5Aax} = 12.0$ Hz, $J_{5Aeq,4A} = 4.7$ Hz, 1H, H5^A_{eq}), 3.95 (dd, $J_{5Beq,5Bax} = 12.0$ Hz, $J_{5Beq,4B} = 5.0$ Hz, 1H, H5^B_{eq}), 3.86–3.76 (m, 2H, H4^A, H4^B), 3.73 (dd, $J_{5C',5C} = 12.4$ Hz, $J_{5C',4C} = 6.2$ Hz, 1H, H5^{'C}), 3.50 (dd, $J_{5Aax,5Aeq} = 12.0$ Hz, $J_{5ax,4A} = 8.6$ Hz, 1H, H5^A_{ax}), 3.27 (dd, $J_{5Bax,5Beq} = 12.0$ Hz, $J_{5Bax,4B} = 9.1$ Hz, 1H, H5^B_{ax}), 2.75–2.69 (m, 4H), 2.62–2.36 (m, 8H), 2.17 (s, 3H), 2.09 (s, 3H), 2.04 (s, 3H). ¹³C {¹H} NMR (101 MHz, CDCl₃) δ 206.5, 206.4, 206.1, 172.0, 171.5, 171.3, 165.7, 165.4, 165.3, 165.2, 133.5, 133.5, 133.5, 133.4, 132.8, 132.7 ($\times 2$), 130.1 ($\times 2$), 130.1 ($\times 2$), 130.0 ($\times 2$), 129.9 ($\times 2$), 129.5, 129.4, 129.2, 129.1, 129.1 ($\times 2$), 128.6 ($\times 4$), 128.1, 100.8 (C1^B),

99.3 (C1^C), 86.5 (C1^A), 75.0 (C4^A), 74.7 (C4^B), 72.7 (C3^A), 72.2 (C3^B), 71.4 (C2^B), 70.4 (C2^A), 70.2 (C2^C), 69.7 (C3^C), 68.8 (C4^C), 65.8 (C5^A), 62.9 (C5^B), 61.1 (C5^C), 37.9, 37.8, 37.8, 29.9, 29.8, 29.8, 28.1, 28.0, 27.9. HRMS: m/z calcd for C₆₄H₆₄O₂₂SNa [M + Na]⁺ 1239.3502, found 1239.3483.

Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-2,3-di-O-levulinoyl-β-D-xylopyranosyl-

(1→4)-2-O-benzoyl-3-O-levulinoyl-β-D-xylopyranosyl-(1→4)-2,3-di-O-benzoyl-1-thio-β-D-

xylopyranoside (26). General procedure B with crushed mol sieves (1.2 g), donor **25** (0.414 g,

0.34 mmol), AgOTf (0.175 g, 0.68 mmol), *p*NO₂PhSCl (0.064 g, 0.34 mmol) and acceptor **6**

(0.139 g, 0.31 mmol). Reaction time 65 min then Et₃N (0.14 mL, 1.02 mmol). Purification by

flash column chromatography (hexane/toluene/acetone 3:2:1) gave **26** as a white amorphous

solid (0.358 g, 75%). In addition, donor **25** was reisolated (0.064 g, 15%). *R*_f 0.32

(hexane/EtOAc 1:1). [α]_D²⁵ -35.1 (*c* 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 8.01–7.92 (m,

12H), 7.58–7.50 (m, 6H), 7.45–7.31 (m, 14H), 7.28–7.24 (m, 3H), 5.69 (t, *J* = 6.6 Hz, 1H, H3^D),

5.60 (t, *J* = 8.0 Hz, 1H, H3^A), 5.32–5.26 (m, 2H, H2^A, H4^D), 5.21 (dd, *J*_{2D,3D} = 6.5 Hz, *J*_{1D,2D} =

4.8 Hz, 1H, H2^D), 5.13–5.07 (m, 2H, H3^B, H3^C), 4.98–4.95 (m, 2H, H1^A, H2^B), 4.84 (d, *J*_{1D,2D} =

4.7 Hz, 1H, H1^D), 4.73–4.68 (m, 2H, H1^B, H2^C), 4.46 (dd, *J*_{5Deq,5Dax} = 12.4 Hz, *J*_{5Deq,4D} = 3.9 Hz,

1H, H5^D_{eq}), 4.21–4.15 (m, 2H, H5^A, H1^C), 4.02–3.96 (m, 1H, H4^A), 3.88 (dd, *J*_{5C,5'C} = 11.9 Hz,

*J*_{5C,4C} = 5.1 Hz, 1H, H5^C), 3.83–3.77 (m, 1H, H4^C), 3.72 (dd, *J*_{5Dax,5Dax} = 12.4 Hz, *J*_{5Dax,4D} = 6.2

Hz, 1H, H5^D_{ax}), 3.56 (dd, *J*_{5Beq,5Bax} = 11.9 Hz, *J*_{5Beq,4B} = 4.6 Hz, 1H, H5^B_{eq}), 3.52–3.44 (m, 2H,

H5^A, H4^B), 3.20–3.14 (m, 2H, H5^C, H5^B_{ax}), 2.73–2.69 (m, 4H), 2.60–2.31 (m, 8H), 2.17 (s,

3H), 2.08 (s, 3H), 2.03 (s, 3H). ¹³C {¹H} NMR (101 MHz, CDCl₃) δ 206.4, 206.3, 206.1, 172.0,

171.6, 171.3, 165.6, 165.4, 165.4, 165.3, 165.1, 165.1, 133.6, 133.5, 133.5, 133.4, 133.4, 133.3,

1
2
3 132.8, 132.6 (×2), 130.1 (×2), 130.0 (×2), 130.0 (×2), 129.9 (×2), 129.9 (×2), 129.9 (×2), 129.6,
4
5 129.5, 129.4, 129.4, 129.2, 129.1, 129.1 (×2), 128.6 (×2), 128.6 (×4), 128.5 (×2), 128.1, 100.5
6
7 (C1^C, $J_{C-H} = 161$ Hz), 100.3 (C1^B, $J_{C-H} = 164$ Hz), 99.3 (C1^D, $J_{C-H} = 165$ Hz), 86.7 (C1^A, $J_{C-H} =$
8
9 159 Hz), 75.0 (C4^A), 74.7 (C4^C), 74.5 (C4^B), 73.0 (C3^A), 72.2 (C3^C), 71.4 (C3^B), 71.2 (C2^C),
10
11 71.0 (C2^B), 70.4, 70.2 (C2^D, C4^D), 69.7 (C3^D), 68.8 (C2^A), 65.8 (C5^A), 62.9 (C5^C), 62.0 (C5^B),
12
13 61.1 (C5^D), 37.8, 37.8 (×2), 29.9, 29.8, 29.8, 28.0, 27.9, 27.8. HRMS: m/z calcd for
14
15 C₈₃H₈₀O₂₈SNa [M + Na]⁺ 1579.4449, found 1579.4425.
16
17
18
19
20
21

22 *Phenyl 2,3,4-tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-β-D-xylopyranosyl-(1→4)-2-O-benzoyl-β-*
23 *D-xylopyranosyl-(1→4)-2,3-di-O-benzoyl-1-thio-β-D-xylopyranoside (27)*. General procedure C
24
25 with tetrasaccharide **26** (0.252 g, 0.16 mmol) and 50% solution of hydrazine hydrate (0.10 mL,
26
27 1.59 mmol). Reaction time 10 min. Eluent for TLC (hexane/toluene/acetone 2:1:1). The product
28
29 was a white powder (0.202 g, 99%). R_f 0.64 (hexane/acetone 1:1). $[\alpha]^{25}_D -9.7$ (c 1.0, CHCl₃).
30
31 ¹H NMR (400 MHz, CDCl₃) δ 8.03–7.91 (m, 12H), 7.59–7.33 (m, 20H), 7.28–7.25 (m, 3H),
32
33 5.80 (t, $J = 8.5$ Hz, 1H, H3^D), 5.58 (t, $J_{2A,3A} = 7.7$ Hz, 1H, H3^A), 5.40–5.34 (m, 2H, H2^D, H4^D),
34
35 5.31 (t, $J = 7.8$ Hz, 1H, H2^A), 4.99 (d, $J_{1A,2A} = 7.8$ Hz, 1H, H1^A), 4.92 (dd, $J_{2B,3B} = 8.6$ Hz, $J_{1B,2B}$
36
37 = 7.0 Hz, 1H, H2^B), 4.79 (d, $J_{1D,2D} = 6.6$ Hz, 1H, H1^D), 4.63 (d, $J_{1B,2B} = 6.9$ Hz, 1H, H1^B), 4.50
38
39 (dd, $J_{5Deq,5Dax} = 11.9$ Hz, $J_{5Deq,4D} = 4.9$ Hz, 1H, H5^D_{eq}), 4.18 (dd, $J_{5Aeq,5Aax} = 12.1$ Hz, $J_{5Aeq,4A} =$
40
41 4.6 Hz, 1H, H5^A_{eq}), 4.14 (d, $J_{1C,2C} = 7.4$ Hz, 1H, H1^C), 4.01–3.95 (m, 1H, H4^A), 3.75–3.42 (m,
42
43 9H, H5^A_{ax}, H3^B, H4^B, H5^B, H3^C, H4^C, H5^C, H5^D_{ax}), 3.37–3.33 (m, 1H, H2^C), 3.15–3.09 (m, 2H,
44
45 H5^B, H5^C). ¹³C {¹H} NMR (101 MHz, CDCl₃) δ 165.7, 165.6, 165.6, 165.5, 165.3, 165.2,
46
47 133.7, 133.6, 133.5, 133.4, 133.3, 133.3, 132.9, 132.5 (×2), 130.0 (×2), 130.0 (×2), 129.9 (×2),
48
49 129.9 (×3), 129.8 (×2), 129.6, 129.6, 129.4, 129.1 (×2), 129.0, 129.0, 128.9, 128.7 (×2), 128.6
50
51
52
53
54
55
56
57
58
59
60

($\times 3$), 128.5 ($\times 2$), 128.5 ($\times 3$), 128.1, 102.5 (C1^C), 101.4 (C1^B), 101.0 (C1^D), 86.6 (C1^A), 78.8 (C4^C), 77.4 (C4^B), 75.7 (C4^A), 74.0 (C3^C), 73.7 (C2^B), 72.8, 72.7 (C3^A, C3^B), 72.1 (C2^C), 71.1, 70.9 (C2^D, C3^D), 70.3 (C2^A), 69.3 (C4^D), 65.7 (C5^A), 63.1 (C5^B/C5^C), 63.0 (C5^B/C5^C), 62.5 (C5^D). HRMS: m/z calcd for C₆₈H₆₂O₂₂SNa [M + Na]⁺ 1285.3345, found 1285.3335.

*Phenyl 2,3,4-tri-O-benzoyl- β -D-xylopyranosyl-(1 \rightarrow 4)-[2,3,5-tri-O-benzoyl- α -L-arabinofuranosyl-(1 \rightarrow 2)]-[2,3,5-tri-O-benzoyl- α -L-arabinofuranosyl-(1 \rightarrow 3)]- β -D-xylopyranosyl-(1 \rightarrow 4)-[2,3,5-tri-O-benzoyl- α -L-arabinofuranosyl-(1 \rightarrow 3)]-2-O-benzoyl- β -D-xylopyranosyl-(1 \rightarrow 4)-2,3-di-O-benzoyl-1-thio- β -D-xylopyranoside (**28**). General procedure D with acceptor **27** (0.088 g, 0.069 mmol), donor **4** (0.175 g, 0.28 mmol) and TMSOTf in dry CH₂Cl₂ (0.17 mL, $c = 0.04$ mmol/mL). Reaction time 105 min then Et₃N (0.02 mL, 0.14 mmol). Purification by flash column chromatography (hexane/toluene/acetone 3:3:1) gave **28** as a white amorphous solid (0.168 g, 91%). R_f 0.26 (hexane/toluene/acetone 3:3:1). $[\alpha]_D^{25} -58.4$ (c 1.0, CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 8.22 (d, $J = 8.1$ Hz, 2H), 8.12–8.07 (m, 6H), 8.03–7.93 (m, 14H), 7.86–7.79 (m, 8H), 7.73 (d, $J = 7.3$ Hz, 2H), 7.58 (d, $J = 7.4$ Hz, 1H), 7.62–7.10 (m, 46H), 7.03 (t, $J = 7.8$ Hz, 1H), 5.69 (s, 1H, H1^F), 5.63 (t, $J = 9.6$ Hz, 1H, H4^D), 5.58 (s, 1H, H1^G), 5.53 (s, 1H), 5.51 (d, $J = 4.4$ Hz, 1H), 5.40–5.39 (m, 3H, H3^A), 5.37–5.34 (m, 2H), 5.21–5.16 (m, 4H, H2^A, H2^D, H1^E), 5.13–5.03 (m, 3H, H2^B, H4^D), 5.00–4.93 (m, 2H), 4.86–4.74 (m, 7H, H1^A), 4.31 (d, $J_{B1,B2} = 6.6$ Hz, 1H, H1^B), 4.02–3.96 (m, 2H, H5^A, H1^D), 3.90 (t, $J = 8.7$ Hz, 1H, H3^B), 3.83–3.78 (m, 2H, H1^C, H3^C), 3.58–3.52 (m, 2H, H4^B, H2^C), 3.45 (td, $J = 9.5, 5.0$ Hz, 1H, H4^C), 3.41–3.35 (m, 2H, H4^A, H5^D), 3.32–3.24 (m, 2H, H5^{'A}, H5^B_{eq}), 3.09–3.00 (m, 2H, H5^C, H5^{'D}), 2.92 (dd, $J_{5Bax,5Beq} = 12.2$ Hz, $J_{5Bax,4B} = 7.5$ Hz, 1H, H5^B_{ax}), 2.45 (t, $J = 10.8$ Hz, 1H, H5^{'C}). ¹³C {¹H} NMR (101 MHz, CDCl₃) δ 166.6 ($\times 2$), 166.4, 166.3, 165.9, 165.8, 165.8,*

1
2
3 165.6, 165.5, 165.3, 165.2, 164.9, 164.9, 164.9, 164.4, 134.1, 134.0, 133.5, 133.4, 133.4, 133.3,
4
5 133.3, 133.1, 133.0, 132.9, 132.8, 132.1, 130.3, 130.2, 130.1, 130.1, 130.0, 130.0, 130.0, 129.9,
6
7 129.9, 129.9, 129.8, 129.8, 129.7, 129.7, 129.6, 129.5, 129.3, 129.3, 129.2, 129.1, 129.0, 129.0,
8
9 129.0, 128.9, 128.6, 128.6, 128.5, 128.5, 128.4, 128.3, 128.3, 128.2, 128.2, 106.1 (C1^G), 105.7
10
11 (C1^F), 105.4 (C1^E), 100.1 (C1^B), 99.8 (C1^D), 99.6 (C1^C), 86.2 (C1^A), 82.0, 81.9, 81.9 (×2), 81.7,
12
13 81.6, 78.9, 78.7, 78.4, 77.4 (C2^C), 76 (C3^C), 75.5 (C3^B), 74.7 (C4^C), 73.6, 73.3, 73.1 (×2), (C3^A,
14
15 C4^A, C2^B, C4^B), 72.4 (C3^D), 71.1, 70.3 (C2^A, C2^D), 69.8 (C4^D), 66.1 (C5^A), 64.4, 64.1, 64.1
16
17 (C5^E, C5^F, C5^G), 63.1, 62.8 (C5^C, C5^D), 62.0 (C5^B). HRMS: *m/z* calcd for C₁₄₆H₁₂₂O₄₃SNa [M +
18
19 Na]⁺ 2617.6973, found 2617.6916.
20
21
22
23
24
25

26 *2,3,4-Tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-[2,3,5-tri-O-benzoyl-α-L-arabinofuranosyl-*
27
28 *(1→3)]-2-O-benzoyl-β-D-xylopyranosyl-(1→4)-[2,3,5-tri-O-benzoyl-α-L-arabinofuranosyl-*
29
30 *(1→3)]-2-O-benzoyl-β-D-xylopyranosyl-(1→4)-[2,3,5-tri-O-benzoyl-α-L-arabinofuranosyl-*
31
32 *(1→3)]-2-O-benzoyl-D-xylopyranose (29)*. General procedure E with thioglycoside **19** (0.081 g,
33
34 0.031 mmol), NBS (0.030 g, 0.17 mmol), additional NBS added after 2 h and 15 min (0.026 g,
35
36 0.15 mmol). Reaction time 3 h. Purification by flash column chromatography (heptane/acetone
37
38 3:2) gave **29** as a white amorphous solid (0.072 g, α/β ratio ~1:0.3, 92%). *R*_f 0.42
39
40
41 (heptane/EtOAc 1:1). ¹H NMR (400 MHz, CDCl₃) δ 8.18–7.80 (m, 39H), 7.64–7.19 (m, 58.5H),
42
43 5.78 (t, *J* = 8.9 Hz, 0.2H) 5.71 (t, *J* = 9.3 Hz, 1H, H3^D), 5.53–5.49 (m, 4H, H1^E/H1^F), 5.46–5.42
44
45 (m, 2H), 5.39–5.35 (m, 4H, H1^A), 5.30–5.28 (m, 2H, H2^D, H1^G), 5.26 (s, 1H, H1^E/H1^F), 5.16
46
47 (td, *J* = 9.3, 5.5 Hz, 1H, H4^D), 5.09–4.84 (m, 15H, H2^A, H2^B, H2^C), 4.48 (d, *J* = 7.8 Hz, 0.3H,
48
49 H1^A_β), 4.36 (t, *J* = 9.3 Hz, 1H, H3^A), 4.25 (dd, *J* = 8.0, 3.7 Hz, 1H, H1^B), 4.11–4.05 (m, 2H,
50
51 H1^D), 3.91–3.81 (m, 4H, H3^B, H3^C, H5^D), 3.76 (td, *J* = 9.5, 5.0 Hz, 1H, H4^A), 3.58–3.46 (m, 4H,
52
53
54
55
56
57
58
59
60

1
2
3 H5^A, H5'^A, H1^C, H4^C), 3.37–3.29 (m, 2H, H4^B, H5^C), 3.24–3.19 (m, 1H, H5'^D), 3.12 (dd, *J*=
4 11.9, 5.6 Hz, 1H, H5^B), 2.99–2.94 (m, 0.3H), 2.66–2.59 (m, 2H, H5'^B, H5'^C). ¹³C{¹H} NMR
5 (101 MHz, CDCl₃) δ 167.0, 166.4, 166.4, 166.4, 166.1, 165.9, 165.8, 165.8, 165.8, 165.6, 165.2,
6 165.1, 165.1, 165.0, 164.8, 164.5, 164.4, 164.2, 133.9, 133.9, 133.7, 133.7, 133.5, 133.4, 133.4,
7 133.3, 133.2, 133.2, 133.2, 133.0, 133.0, 132.9, 130.4, 130.4, 130.2, 130.2, 130.1, 130.1, 130.0,
8 129.9, 129.9, 129.8, 129.8, 129.8, 129.7, 129.6, 129.5, 129.5, 129.5, 129.3, 129.2, 129.2, 129.1,
9 129.1, 129.0, 129.0, 129.0, 128.9, 128.9, 128.9, 128.7, 128.6, 128.5, 128.5, 128.4, 128.4, 128.4,
10 128.3, 128.3, 128.3, 128.3, 106.1 (C1^G), 105.7, 105.6 (C1^E, C1^F), 100.4 (C_β), 100.3 (C1^B), 100.0
11 (C1^C), 100.0 (C_β), 99.9 (C1^D), 96.4 (C_β), 90.5 (C1^A), 82.8, 82.7, 82.7, 81.8 (C_β), 81.7, 81.5,
12 80.8, 78.3, 78.3, 78.2, 78.1 (C_β), 76.3 (C_β), 75.4 (C_β), 75.3, 73.3 (C3^B, C3^C), 74.7 (C_β), 74.5,
13 74.3, 74.3 (C4^A, C4^B, C4^C), 74.2 (C_β), 74.2 (C_β), 74.1 (C_β), 73.4 (C2^B), 73.0 (C2^C), 72.1, 72.0
14 (C2^A, C3^D), 71.2 (C2^D), 69.8 (C4^D), 64.0, 63.9, 63.9, 63.8 (C5^E, C5^F, C5^G), 63.3 (C_β), 63.1, 63.0,
15 63.0 (C5^B, C5^C, C5^D), 59.3 (C5^A). HRMS: *m/z* calcd for C₁₄₀H₁₁₈O₄₄SNa [M + Na]⁺ 2525.6888,
16 found 2525.6876.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 *2,3,4-Tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-[2,3,5-tri-O-benzoyl-α-L-arabinofuranosyl-*
40 *(1→3)]-2-O-benzoyl-β-D-xylopyranosyl-(1→4)-[2,3,5-tri-O-benzoyl-α-L-arabinofuranosyl-*
41 *(1→2)]-[2,3,5-tri-O-benzoyl-α-L-arabinofuranosyl-(1→3)]-α-D-xylopyranosyl-(1→4)-2,3-di-*
42 *O-benzoyl-D-xylopyranose (30)*. General procedure E with thioglycoside **23** (0.200 g, 0.077
43 mmol), acetone/H₂O 9:1 (1.23 mL), NBS (0.055 g, 0.31 mmol), additional NBS added after 30
44 min (0.030 g, 0.17 mmol). Reaction time 60 min. Eluent for TLC and flash column
45 chromatography (heptane/acetone 3:2). Product was isolated as a white amorphous solid (0.181
46 g, α/β ratio ~1:0.35, 93%). *R*_f 0.45 (heptane/EtOAc 1:1). ¹H NMR (400 MHz, CDCl₃) δ
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 8.19–8.18 (m, 2H), 8.10–7.80 (m, 27H), 7.78–7.74 (m, 1H), 7.69–7.64 (m, 2H), 7.59–7.20 (m,
4 39H), 7.18–7.07 (m, 4H), 5.86–5.82 (m, 1H, H3^A), 5.79–5.78 (m, 1H, H1^F), 5.69 (t, $J = 9.2$ Hz,
5 1H, H3^D), 5.63–5.61 (m, 1H, H1^E), 5.55–5.50 (m, 6H), 5.36–5.35 (m, 2H, H1^A), 5.26 (s, 1H,
6 H1^G), 5.23 (dd, $J_{2D,3D} = 9.3$ Hz, $J_{1D,2D} = 7.4$ Hz, 1H, H2^D), 5.17 (td, $J_{3D,4D} = 9.2$ Hz, $J_{4D,5D} = 5.5$
7 Hz, 1H, H4^D), 5.12–5.07 (m, 2H), 5.05–4.81 (m, 7H, H2^A, H2^C), 4.76–4.64 (m, 2H), 4.57 (d, $J =$
8 7.3 Hz, 0.33H, H1^{A_β}), 4.31 (t, $J_{1B,2B} = 7.5$ Hz, 1H, H1^B), 4.14 (d, $J_{1D,2D} = 7.3$ Hz, 1H, H1^D), 4.00
9 (dd, $J = 11.9, 5.3$ Hz, 0.40H), 3.93–3.82 (m, 6H, H4^A, H5^A, H3^B, H1^C, H3^C, H5^D), 3.74–3.69 (m,
10 1H, H5^{'A}), 3.66 (dd, $J_{2B,3B} = 9.6$ Hz, $J_{1B,2B} = 6.9$ Hz, 1H, H2^B), 3.54–3.38 (m, 3H, H4^A, H4^C),
11 3.36–3.24 (m, 2H, H5^B, H5^{'D}), 3.07–3.00 (m, 1H, H5^C), 2.74–2.63 (m, 1H, H5^{'B}), 2.53 (t, $J =$
12 11.1 Hz, 1H, H5^{'C}). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 166.9, 166.7, 166.5, 166.4, 166.1,
13 165.9, 165.9, 165.8, 165.8, 165.7, 165.7, 165.6, 165.6, 165.4, 165.4, 165.4, 165.1, 165.0, 164.6,
14 164.2, 133.7, 133.6, 133.6, 133.5, 133.4, 133.4, 133.3, 133.2, 133.1, 133.0, 132.9, 132.8, 130.3,
15 130.2, 130.2, 130.1, 130.0, 130.0, 129.9, 129.9, 129.9, 129.8, 129.8, 129.8, 129.6, 129.6, 129.3,
16 129.2, 129.2, 129.2, 129.1, 129.1, 129.1, 129.0, 128.9, 128.8, 128.7, 128.7, 128.6, 128.6, 128.5,
17 128.4, 128.4, 128.3, 128.2, 107.2 (C_β), 107.1 (C1^E), 106.2 (C1^G), 105.5 (C1^F), 101.3 (C_β), 100.9
18 (C1^B), 99.8 (C1^D), 99.6 (C1^C), 96.0 (C_β), 90.5 (C1^A), 82.7, 82.5 (C_β), 82.3 (C_β), 82.3, 82.0, 81.9,
19 81.8, 81.8 (C_β), 81.7 (C_β), 80.8, 80.3 (C_β) 80.0 (C2^B), 78.6, 78.2, 77.9, 77.8 (C_β), 76.0
20 (C3^B/C3^C), 75.9 (C_β), 75.7 (C3^B/C3^C), 74.6 (C4^C), 74.2, 74.0 (C4^A/C4^B), 74.0, 73.8 (C4^A/C4^B),
21 73.2 (C2^C), 72.4, 72.0 (2C, C2^A, C3^D), 71.2 (C2^D), 70.4 (C3^A), 69.7 (C4^D), 64.4 (C_β), 64.3, 64.1,
22 63.9 (C5^E, C5^F, C5^G), 63.1 (C_β), 63.0 (C5^C), 62.9 (C5^D), 62.4 (C5^B), 59.4 (C5^A). HRMS: m/z
23 calcd for C₁₄₀H₁₁₈O₄₄S(Na⁺)₂ [M + 2Na]²⁺ 1274.3390, found 1274.3381.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 *2,3,4-Tri-O-benzoyl-β-D-xylopyranosyl-(1→4)-[2,3,5-tri-O-benzoyl-α-L-arabinofuranosyl-*
5 *(1→2)]-[2,3,5-tri-O-benzoyl-α-L-arabinofuranosyl-(1→3)]-β-D-xylopyranosyl-(1→4)-[2,3,5-*
6 *tri-O-benzoyl-α-L-arabinofuranosyl-(1→3)]-2-O-benzoyl-β-D-xylopyranosyl-(1→4)-2,3-di-O-*
7 *benzoyl-D-xylopyranose (31).* General procedure E with thioglycoside **28** (0.190 g, 0.073 mmol),
8
9
10
11
12 acetone/H₂O 9:1 (1.1 mL), NBS (0.057 g, 0.32 mmol), additional NBS added after 30 min
13
14 (0.033 g, 0.19 mmol). Reaction time 60 min. Eluent for TLC and flash column chromatography
15
16 (heptane/acetone 3:2). Product was isolated as a white amorphous solid (0.169 g, α/β ratio
17
18 ~1:0.3, 92%). *R*_f 0.24 (heptane/EtOAc 1:1). ¹H NMR (400 MHz, CDCl₃) δ 8.22 (d, *J* = 8.1 Hz,
19
20 2H), 8.13–8.03 (m, 8H), 7.99–7.94 (m, 8H), 7.86–7.80 (m, 7H), 7.75–7.72 (m, 2H), 7.69 (d, *J* =
21
22 7.4 Hz, 2H), 7.65–7.59 (m, 5H), 7.56–7.25 (m, 31H), 7.20–7.11 (m, 8H), 7.08–7.03 (m, 2H),
23
24 5.79 (t, *J* = 9.5 Hz, 1H, H^{3A}), 5.71 (s, 1H, H^{1G}), 5.64 (t, *J* = 9.6 Hz, 1H, H^{3D}), 5.58 (d, *J* = 2.7
25
26 Hz, 1H, H^{1F}), 5.54 (s, 1H), 5.53–5.49 (m, 2H, H^{1A}), 5.45–5.41 (m, 2H), 5.37 (t, *J* = 4.8 Hz, 2H),
27
28 5.29 (s, 0.2H), 5.21–5.19 (m, 3H, H^{2D}, H^{1E}), 5.16–4.94 (m, 6H, H^{2A}, H^{2B}, H^{4D}), 4.87–4.83 (m,
29
30 3H), 4.81–4.75 (m, 2H), 4.71 (d, *J* = 7.7 Hz, 1H, H^{1Aβ}), 4.38 (d, *J*_{1B,2B} = 6.5 Hz, 1H, H^{1B}), 4.35
31
32 (d, *J* = 6.6 Hz, 0.3H), 3.98–3.92 (m, 2H, H^{3B}, H^{1D}), 3.84–3.79 (m, 2H, H^{1C}, H^{3C}), 3.76 (t, *J* =
33
34 11.0 Hz, 1H, H^{5A}), 3.62–3.30 (m, 7H, H^{4A}, H^{5'A}, H^{4B}, H^{5B}, H^{2C}, H^{4C}, H^{5D}), 3.23 (t, *J* = 11.2
35
36 Hz, 0.4H), 3.10–2.96 (m, 3H, H^{5'B}, H^{5C}, H^{5'D}), 2.49–2.44 (m, 1H, H^{5'C}). ¹³C{¹H} NMR (101
37
38 MHz, CDCl₃) δ 167.1, 166.6 (×2), 166.4, 166.3, 166.0, 165.9, 165.8, 165.8, 165.8, 165.8, 165.6,
39
40 165.5, 165.5, 165.4, 165.4, 165.0, 165.0, 164.9, 164.9, 164.9, 164.4, 164.4, 134.0, 133.5, 133.5,
41
42 133.5, 133.4, 133.4, 133.3, 133.3, 133.1, 133.0, 132.9, 132.9, 132.9, 130.3, 130.2, 130.1, 130.1,
43
44 130.0, 129.9, 129.9, 129.9, 129.8, 129.8, 129.7, 129.7, 129.6, 129.4, 129.3, 129.2, 129.1, 129.0,
45
46 129.0, 128.9, 128.9, 128.7, 128.6, 128.6, 128.5, 128.4, 128.3, 128.3, 128.3, 128.2, 106.1 (C^{1F}),
47
48 105.7 (C^{1G}), 105.4 (C^{1E}), 105.3 (C_β), 100.2 (C_β), 100.1 (C^{1B}), 99.8 (C^{1D}), 99.6 (C^{1C}), 96.3
49
50
51
52
53
54
55
56
57
58
59
60

(C_β), 90.5 (C1^A), 82.0, 82.0, 81.9, 81.8, 81.8 (C_β), 81.7, 81.5, 79.0, 78.6, 78.3, 77.4 (C2^C), 76.7 (C3^C), 75.4 (C3^B), 74.7, 74.5 (C4^A, C4^C), 74.3 (C_β), 74.2 (C_β), 73.1, 73.1 (C2^B, C4^B), 72.4, 72.3 (C2^A, C3^D), 71.1 (C2^B), 70.1 (C3^C), 69.7 (C4^D), 64.3, 64.1, 64.1 (C5^E, C5^F, C5^G), 63.5 (C_β), 63.0, 62.8 (C5^C, C5^D), 62.0 (C5^B), 59.3 (C5^A). HRMS: *m/z* calcd for C₁₄₀H₁₁₈O₄₄S(Na⁺)₂ [M + 2Na]²⁺ 1274.3390, found 1274.3378.

Phenyl β-D-xylopyranosyl-(1→4)-[α-L-arabinofuranosyl-(1→3)]-β-D-xylopyranosyl-(1→4)-[α-L-arabinofuranosyl-(1→3)]-β-D-xylopyranosyl-(1→4)-[α-L-arabinofuranosyl-(1→3)]-1-thio-β-D-xylopyranoside (32). General procedure F with heptasaccharide **19** (0.028 g, 0.011 mmol).

The product was obtained as a white powder (0.009 g, 77%). *R*_f 0.63

(EtOAc/MeOH/H₂O/AcOH 6:3:0.8:0.2). [α]²⁵_D -30 (*c* 0.23, H₂O). ¹H NMR (800 MHz, D₂O) δ 7.60–7.57 (m, 2H), 7.47–7.43 (m, 3H), 5.43 (s, 1H, H1^{arabinose}), 5.41 (s, 1H, H1^{arabinose}), 5.40 (s, 1H, H1^{arabinose}), 4.82 (d, *J* = 9.6 Hz, 1H, H1^A), 4.49–4.48 (m, 2H, H1^B, H1^C), 4.45 (d, *J* = 7.8 Hz, 1H, H1^D), 4.30–4.27 (m, 3H, H4^{arabinose}), 4.19–4.17 (m, 4H, H5^A, H2^{arabinose}), 4.10–4.06 (m, 2H, H5^B, H5^C), 3.93–3.91 (m, 4H, H5^D, H3^{arabinose}), 3.82–3.79 (m, 7H, H3^A, H4^A, H4^B, H4^C, H5^{arabinose}), 3.76–3.72 (m, 5H, H3^B, H3^C, H5^{arabinose}), 3.61 (td, *J* = 9.9, 5.5 Hz, 1H, H4^D), 3.53 (t, *J* = 8.8 Hz, 1H, H2^A), 3.46–3.41 (m, 4H, H5^A, H2^B, H2^C, H3^D), 3.37 (t, *J* = 11.1 Hz, 2H, H5^B, H5^C), 3.28 (t, *J* = 11.1 Hz, 1H, H5^D), 3.26–3.24 (m, 1H, H2^D). ¹³C {¹H} NMR (201 MHz, D₂O) δ 132.4 (×2), 131.2, 129.4 (×2), 128.4, 107.7, 107.6, 107.6 (C1^{arabinose}), 101.4 (C1^D), 101.2, 101.2 (C1^B, C1^C), 87.8 (C1^A), 84.8, 84.7, 84.7 (C4^{arabinose}), 80.7, 80.7, 80.7 (C2^{arabinose}), 78.7 (C3^A), 77.3, 77.2, 77.2, 77.2 (C3^B, C3^C, C3^{arabinose}), 75.6 (C3^D), 73.6, 73.6, 73.5, 73.5, 73.5 (C4^A, C2^B, C4^B, C2^C, C4^C), 72.9 (C2^D), 71.9 (C2^A), 69.2 (C4^D), 66.3 (C5^A), 65.1 (C5^D), 62.7, 62.7

(C5^B, C5^C), 61.3 (C5^{arabinose}). HRMS: m/z calcd for C₄₁H₆₃O₂₈S [M + H]⁺ 1035.3221, found 1035.3214.

Phenyl β-D-xylopyranosyl-(1→4)-[α-L-arabinofuranosyl-(1→3)]-β-D-xylopyranosyl-(1→4)-[α-L-arabinofuranosyl-(1→2)]-[α-L-arabinofuranosyl-(1→3)]-β-D-xylopyranosyl-(1→4)-1-thio-β-D-xylopyranoside (33). General procedure F with heptasaccharide **23** (0.071 g, 0.027 mmol).

The product was obtained as a white powder (0.023 g, 82%). R_f 0.53

(EtOAc/MeOH/H₂O/AcOH 6:3:0.8:0.2). $[\alpha]_D^{25}$ -36 (c 0.21, H₂O). ¹H NMR (800 MHz, D₂O) δ 7.59–7.58 (m, 2H), 7.46–7.42 (m, 3H), 5.42 (s, 1H, H1^G), 5.28 (s, 1H, H1^F), 5.23 (s, 1H, H1^E), 4.77 (d, J = 9.5 Hz, 1H, H1^A), 4.64 (d, J = 7.2 Hz, 1H, H1^B), 4.49 (d, J = 7.9 Hz, 1H, H1^C), 4.45 (d, J = 7.8 Hz, 1H, H1^D), 4.32 (q, J = 5.4 Hz, 1H, H4^F), 4.29 (q, J = 5.2 Hz, 1H, H4^G), 4.19–4.14 (m, 5H, H5^A, H5^B, H2^E, H2^F, H2^G), 4.12 (td, J = 5.8, 3.5 Hz, 1H, H4^E), 4.09 (dd, J = 11.8, 5.2 Hz, 1H, H5^C), 3.96–3.95 (m, 2H, H3^E, H3^F), 3.94–3.91 (m, 2H, H5^D, H3^G), 3.89 (td, J = 9.0, 5.1 Hz, 1H, H4^B), 3.84–3.77 (m, 6H, H4^A, H3^B, H4^C, H5^E, H5^F, H5^G), 3.77–3.70 (m, 4H, H3^C, H5^{'E}, H5^{'F}, H5^{'G}), 3.63–3.59 (m, 2H, H3^A, H4^D), 3.59–3.57 (m, 1H, H2^B), 3.47–3.36 (m, 6H, H2^A, H5^{'A}, H5^{'B}, H2^C, H5^{'C}, H3^D), 3.29 (t, J = 11.4 Hz, 1H, H5^{'D}), 3.26 (dd, J = 9.2, 8.0 Hz, 1H, H2^D). ¹³C {¹H} NMR (201 MHz, D₂O) δ 132.1 (×2), 131.4, 129.4 (×2), 128.3, 108.7 (C1^E), 108.1 (C1^F), 107.6 (C1^G), 101.4 (C1^D), 101.2 (C1^C), 99.8 (C1^B), 88.0 (C1^A), 84.8 (C4^G), 84.4 (C4^E), 84.2 (C4^F), 81.2, 81.0, 80.7 (C2^E, C2^F, C2^G), 78.6 (C2^B), 77.6 (C3^B), 77.2, 77.2, 77.2 (C3^C, C3^F, C3^G), 76.6 (C3^E), 75.6 (C3^D), 75.4 (C4^A), 75.2 (C3^A), 73.8, 73.7, 73.6 (C4^B, C2^C, C4^C), 72.9 (C2^D), 71.6 (C2^A), 69.2 (C4^D), 66.5 (C5^A), 65.1 (C5^D), 62.7 (C5^C), 62.5 (C5^B), 61.3, 61.2, 61.1 (C5^E, C5^F, C5^G). HRMS: m/z calcd for C₄₁H₆₃O₂₈S [M + H]⁺ 1035.3221, found 1035.3216.

1
2
3
4
5
6 Phenyl β -D-xylopyranosyl-(1 \rightarrow 4)-[α -L-arabinofuranosyl-(1 \rightarrow 2)]-[α -L-arabinofuranosyl-
7
8 (1 \rightarrow 3)]- β -D-xylopyranosyl-(1 \rightarrow 4)-[α -L-arabinofuranosyl-(1 \rightarrow 3)]- β -D-xylopyranosyl-(1 \rightarrow 4)-1-
9
10 thio- β -D-xylopyranoside (**34**). General procedure F with heptasaccharide **28** (0.061 g, 0.024
11
12 mmol). Product was obtained as a white powder (0.022 g, 91%). R_f 0.53
13
14 (EtOAc/MeOH/H₂O/AcOH 6:3:0.8:0.2). $[\alpha]^{25}_D$ -22 (c 0.28, H₂O). ¹H NMR (800 MHz, D₂O) δ
15
16 7.59–7.58 (m, 2H), 7.46–7.43 (m, 3H), 5.43 (s, 1H, H1^E), 5.29 (s, 1H, H1^G), 5.25 (s, 1H, H1^F),
17
18 4.79 (d, J = 9.2 Hz, 1H, H1^A), 4.61 (d, J = 7.5 Hz, 1H, H1^C), 4.50 (d, J = 7.7 Hz, 1H, H1^B), 4.44
19
20 (d, J = 7.8 Hz, 1H, H1^D), 4.34–4.31 (m, 2H, H4^E, H4^G), 4.18–4.13 (m, 6H, H5^A, H5^B, H2^E, H2^F,
21
22 H4^F, H2^G), 4.10 (dd, J = 11.9, 4.4 Hz, 1H, H5^C), 3.99 (dd, J = 5.8, 3.1 Hz, 1H, H3^F), 3.95–3.92
23
24 (m, 3H, H5^D, H3^E, H3^G), 3.86–3.73 (m, 11H, H4^A, H3^B, H4^B, H3^C, H4^C, H5^E, H5^{'E}, H5^F, H5^{'F},
25
26 H5^G, H5^{'G}), 3.63–3.60 (m, 2H, H3^A, H4^D), 3.57 (t, J = 7.8 Hz, 1H H2^C), 3.48–3.38 (m, 6H, H2^A,
27
28 H5^{'A}, H2^B, H5^{'B}, H5^{'C}, H3^D), 3.30–3.25 (m, 2H, H2^D, H5^{'D}). ¹³C{¹H} NMR (201 MHz, D₂O) δ
29
30 132.1 (\times 2), 131.4, 129.4 (\times 2), 128.4, 108.7 (C1^F), 108.0 (C1^G), 107.5 (C1^E), 101.6 (C1^B), 101.3
31
32 (C1^D), 99.8 (C1^C), 87.9 (C1^A), 85.0 (C4^E/C4^G), 84.3 (C4^F), 84.2 (C4^E/C4^G), 81.3 (C2^F), 80.9,
33
34 80.6 (C2^E, C2^G), 78.7 (C2^C), 77.7 (C3^C), 77.2, 77.2 (C3^E, C3^G), 77.0 (C3^B), 76.7 (C3^F), 76.1
35
36 (C4^A), 75.6 (C3^D), 75.2 (C3^A), 73.6, 73.3 (C4^B, C4^C), 73.3 (C2^B), 73.0 (C2^D), 71.6 (C2^A), 69.2
37
38 (C4^D), 66.5 (C5^A), 65.1 (C5^D), 62.8 (C5^B), 62.6 (C5^C), 61.3, 61.1, 61.1 (C5^E, C5^F, C5^G). HRMS:
39
40 m/z calcd for C₄₁H₆₃O₂₈S [M + H]⁺ 1035.3221, found 1035.3218.
41
42
43
44
45
46
47
48
49

50 β -D-Xylopyranosyl-(1 \rightarrow 4)-[α -L-arabinofuranosyl-(1 \rightarrow 3)]- β -D-xylopyranosyl-(1 \rightarrow 4)-[α -L-
51
52 arabinofuranosyl-(1 \rightarrow 3)]- β -D-xylopyranosyl-(1 \rightarrow 4)-[α -L-arabinofuranosyl-(1 \rightarrow 3)]-D-
53
54 xylopyranose (**1**). General procedure G with partially deprotected heptasaccharide **32** (10.0 mg,
55
56
57
58
59
60

0.097 mmol), 2,6-lutidine (3 μ L, 0.015 mmol) and NBS (5.3 mg, 0.030 mmol). Reaction time 1.5 h. The product was obtained as a white powder (8.8 mg, 97%). α/β ratio 0.7:1. R_f 0.45 (EtOAc/MeOH/H₂O/AcOH 6:3:0.8:0.2). ¹H NMR (800 MHz, D₂O) δ 5.41–5.41 (m, 4.01H, H1^F, H1^G), 5.36 (s, 0.73H, H1^{E α}), 5.19 (d, J = 3.5 Hz, 0.70H, H1^{A α}), 4.64 (d, J = 7.9 Hz, 1H, H1^{A β}), 4.51–4.49 (m, 3.35H, H1^B, H1^C), 4.45–4.44 (m, 1.76H, H1^D), 4.31–4.28 (m, 4.90H, H4^E, H4^F, H4^G), 4.19–4.17 (m, 4.94H, H2^E, H2^F, H2^G), 4.10–4.07 (m, 4.29H, H5^{A β} , H5^B, H5^C), 3.94–3.91 (m, 7.29H, H3^{A α} , H5^D, H3^E, H3^F, H3^G), 3.87–3.78 (m, 12.26H, H4^{A α} , H4^{A β} , H5^{A α} , H5^{A α} , H4^B, H5^E, H5^F, H5^G), 3.76–3.72 (m, 9.69H, H3^{A β} , H3^B, H3^C, H5^{'E}, H5^{'F}, H5^{'G}), 3.70 (dd, J = 9.2, 3.6 Hz, 1.06H, H2^{A α}), 3.63–3.59 (m, 2H, H4^D), 3.47–3.36 (m, 10.88H, H2^{A β} , H5^{A β} , H2^B, H5^{'B}, H2^C, H5^{'C}, H3^D), 3.30–3.24 (m, 3.99H, H2^D, H5^{'D}). ¹³C {¹H} NMR (201 MHz, D₂O) δ 107.8, 107.7, 107.7, 107.6 (C1^E, C1^F, C1^G), 101.4, 101.3, 101.3, 101.2 (C1^B, C1^C, C1^D), 96.4 (C1^{A β}), 92.2 (C1^{A α}), 84.8, 84.7, 84.7, 84.6 (C4^E, C4^F, C4^G), 80.7, 80.7, 80.7 (C2^E, C2^F, C2^G), 77.7 (C3^{A β}), 77.3, 77.3, 77.2, 77.2 (C3^B, C3^C, C3^E, C3^F, C3^G), 75.6 (C3^D), 75.2 (C3^{A α}), 74.5 (C2^{A β}), 73.8, 73.7, 73.6, 73.6, 73.6, 73.5, 73.5 (C4^{A α} , C4^{A β} , C2^B, C4^B, C2^C, C4^C), 72.9 (C2^D), 71.7 (C2^{A α}), 69.2 (C4^D), 65.1 (C5^D), 62.8, 62.7, 62.7, 62.7 (C5^{A β} , C5^B, C5^C), 61.3 (C5^E, C5^F, C5^G), 59.1 (C5^{A α}). HRMS: m/z calcd for C₃₅H₅₈O₂₉Na [M + Na]⁺ 965.2956, found 965.2962.

β -D-Xylopyranosyl-(1 \rightarrow 4)-[α -L-arabinofuranosyl-(1 \rightarrow 3)]- β -D-xylopyranosyl-(1 \rightarrow 4)-[α -L-arabinofuranosyl-(1 \rightarrow 2)]-[α -L-arabinofuranosyl-(1 \rightarrow 3)]- β -D-xylopyranosyl-(1 \rightarrow 4)-D-xylopyranose (2). General procedure G with partially deprotected heptasaccharide **33** (13.4 mg, 0.013 mmol), 2,6-lutidine (4 μ L, 0.019 mmol) and NBS (6.8 mg, 0.038 mmol). Reaction time 1.5 h. The product was obtained as a white powder (10.2 mg, 84%). α/β ratio 0.5:1. R_f 0.31

(EtOAc/MeOH/H₂O/AcOH 6:3:0.8:0.2). ¹H NMR (800 MHz, D₂O) δ 5.43 (s, 1.54H, H1^G), 5.29 (s, 1.40H, H1^F), 5.25 (s, 0.53H, H1^{E_{α/β}}), 5.25 (s, 0.86H, H1^{E_{α/β}}), 5.21 (d, *J* = 3.6 Hz, 0.53H, H1_α), 4.66 (d, *J* = 7.2 Hz, 1.37H, H1^B), 4.60 (d, *J* = 7.9 Hz, 1H, H1^{A_β}), 4.50 (d, *J* = 7.8 Hz, 1.43H, H1^C), 4.46 (d, *J* = 7.8 Hz, 1.49H, H1^D), 4.33 (td, *J* = 5.4, 3.8 Hz, 1.54H, H4^F), 4.30 (q, *J* = 5.3 Hz, 1.66H, H4^G), 4.19–4.09 (m, 11.09H, H5^A, H5^B, H5^C, H2^E, H4^E, H2^F, H2^G), 4.01–4.00 (m, 0.61H, H4^{A_α}), 3.98 (dd, *J* = 5.8, 3.2 Hz, 1.14H, H3^E), 3.96 (dd, *J* = 5.5, 2.4 Hz, 1.62H, H3^F), 3.94–3.92 (m, 3.49H, H5^D, H3^G), 3.89 (dd, *J* = 9.1, 4.9 Hz, 1.58H, H4^B), 3.86–3.73 (m, 19.09H, H3^{A_α}, H4^{A_β}, H5^{A_α}, H5^{A_α}, H3^B, H3^C, H4^C, H5^E, H5^E, H5^{F*}, H5^F, H5^G, H5^G), 3.63–3.57 (m, 4.95H, H2^{A_α}, H3^{A_β}, H2^B, H4^D), 3.47–3.42 (m, 6.09H, H5^{A_β}, H5^B, H2^C, H3^D), 3.40–3.37 (m, 1.75H, H5^C), 3.31–3.25 (m, 4.64H, H2^{A_β}, H2^D, H5^D). ¹³C{¹H} NMR (201 MHz, D₂O) δ 108.7, 108.7 (C1^E), 108.1 (C1^F), 107.6 (C1^G), 101.4 (C1^D), 101.2 (C1^C), 99.9, 99.8 (C1^B), 96.5 (C1^{A_β}), 92.0 (C1^{A_α}), 84.8 (C4^G), 84.4, 84.3 (C4^E), 84.3 (C4^F), 81.2, 81.2 (C2^E), 81.0 (C2^F), 80.7 (C2^G), 78.6, 78.6 (C2^B), 77.6, 77.5 (C3^B), 77.2, 77.2, 77.2 (C3^C, C3^F, C3^G), 76.6 (C3^E), 76.5 (C4^{A_α}), 75.9 (C4^{A_β}), 75.8, 75.6 (C3^D), 74.0, 73.9, 73.8 (C2^{A_β}, C3^{A_β}, C4^B), 73.7, 73.6 (C2^C, C4^C), 72.9 (C2^D), 71.4 (C2^{A_α}), 71.0 (C3^{A_α}), 69.2 (C4^D), 65.1 (C5^D), 62.9, 62.7 (C5^{A_β}, C5^C), 62.5 (C5^B), 62.5, 61.3, 61.2, 61.1, 61.0 (C5^E, C5^F, C5^G), 58.7 (C5^{A_α}). HRMS: *m/z* calcd for C₃₅H₅₈O₂₉Na [M + Na]⁺ 965.2956, found 965.2965.

β-D-Xylopyranosyl-(1→4)-[α-L-arabinofuranosyl-(1→2)]-[α-L-arabinofuranosyl-(1→3)]-β-D-xylopyranosyl-(1→4)-[L-arabinofuranosyl-(1→3)]-β-D-xylopyranosyl-(1→4)-D-xylopyranose (3). General procedure G with partially deprotected heptasaccharide **34** (10.6 mg, 0.010 mmol), 2,6-lutidine (3 μL, 0.015 mmol) and NBS (5.5 mg, 0.031 mmol). Reaction time 1.5 h. The

product was obtained as a white powder (8.5 mg, 88%). α/β ratio 0.6:1. R_f 0.33 (EtOAc/MeOH/H₂O/AcOH 6:3:0.8:0.2). ¹H NMR (800 MHz, D₂O) δ 5.44–5.44 (m, 1.57H, H1^E), 5.30 (s, 1.56H, H1^G), 5.25 (s, 1.55H, H1^F), 5.21 (d, J = 3.6 Hz, 1H, H1^{A α}), 4.62–4.60 (m, 2.70H, H1^{A β} , H1^C), 4.52–4.51 (m, 1.58H, H1^B), 4.45 (d, J = 7.8 Hz, 1.58H, H1^D), 4.34–4.31 (m, 3.29H, H4^E, H4^G), 4.19–4.18 (m, 3.20H, H2^E, H2^F), 4.17–4.15 (m, 5.36H, H5^B, H2^F, H4^F), 4.10 (dd, J = 11.9, 4.2 Hz, 1.60H, H5^C), 4.07 (dd, J = 11.8, 5.4 Hz, 1.09H, H5^{A β}), 4.00 (dd, J = 5.9, 3.1 Hz, 1.57H, H3^F), 3.96–3.93 (m, 5.17H, H5^D, H3^E, H3^G), 3.87–3.73 (m, 22.09H, H3^{A α} , H3^{A β} , H4^{A α} , H5^{A α} , H5^{A β} , H3^B, H4^B, H3^C, H4^C), 3.63–3.55 (m, 5.31H, H2^{A α} , H5^{A β} , H2^C, H4^D), 3.50–3.45 (m, 3.52H, H2^B, H5^B), 3.43 (t, J = 9.3 Hz, 1.94H, H3^D), 3.41–3.38 (m, 2.99H, H3^C, H5^C), 3.30–3.25 (m, 4.54H, H2^{A β} , H2^D, H5^D). ¹³C {¹H} NMR (201 MHz, D₂O) δ 108.7 (C1^F), 108.0 (C1^G), 107.5 (C1^E), 101.6 (C1^B), 101.3 (C1^D), 99.8 (C1^C), 96.5 (C1^{A β}), 92.0 (C1^{A α}), 85.0 (C4^E), 84.3, 84.2 (C4^F, C4^G), 81.3 (C2^F), 80.9 (C2^G), 80.5 (C2^E), 78.7 (C2^C), 77.7 (C3^C), 77.3, 77.2 (C3^E, C3^G), 77.0, 77.0 (C3^B), 76.7 (C3^F), 76.6 (C4^{A α}), 76.4 (C4^{A β}), 75.6 (C3^D), 74.0, 73.9 (C2^{A β} , C3^{A β}), 73.6 (C4^C), 73.3 (C2^B, C4^B), 73.0 (C2^D), 71.3 (C2^{A α}), 70.9 (C3^{A α}), 69.2 (C4^D), 65.1 (C5^D), 62.9, 62.8, 62.8, 62.6 (C3^C) (C5^{A β} , C5^B, C5^C), 61.4, 61.1, 61.1 (C5^E, C5^F, C5^G), 58.8 (C5^{A α}). HRMS: m/z calcd for C₃₅H₅₈O₂₉Na [M + Na]⁺ 965.2956, found 965.2966.

Supporting Information. Copies of ¹H and ¹³C {¹H} NMR spectra for the prepared compounds as well as 2D NMR spectra for compounds **1** – **3**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

ACKNOWLEDGMENT

We thank the Danish Council for Strategic Research for financial support (SET4Future project, grant 0603-00463B). In addition, The NMR Center • DTU and the Villum Foundation are acknowledged for access to the 800 MHz spectrometer.

References

- 1) Naidu, D. S.; Hlangothi, S. P.; John, M. J. Bio-based products from xylan: A review. *Carbohydr. Polym.* **2018**, *179*, 28–41.
- 2) Zhang, Z.; Smith, C.; Li, W. Extraction and modification technology of arabinoxylans from cereal by-products: A critical review. *Food Res. Int.* **2014**, *65*, 423–436.
- 3) (a) Chen, Z.; Li, S.; Fu, Y.; Li, C.; Chen, D.; Chen, H. Arabinoxylan structural characteristics, interaction with gut microbiota and potential health functions. *J. Funct. Foods* **2019**, *54*, 536–551. (b) Mendis, M.; Leclerc, E.; Simsek, S. Arabinoxylans, gut microbiota and immunity. *Carbohydr. Polym.* **2016**, *139*, 159–166.
- 4) (a) Stoklosa, R. J.; Latona, R. J.; Bonnaille, L. M.; Yadav, M. P. Evaluation of arabinoxylan isolated from sorghum bran, biomass and bagasse for film formation. *Carbohydr. Polym.* **2019**, *213*, 382–392. (b) Yu, L.; Yakubov, G. E.; Gilbert, E. P.; Sewell, K.; van de Meene, A. M. L.; Stokes, J. R. Multi-scale assembly of hydrogels formed by highly branched arabinoxylans from *Plantago ovata* seed mucilage studied by USANS/SANS and rheology. *Carbohydr. Polym.* **2019**, *207*, 333–342.
- 5) (a) Biely, P.; Singh, S.; Puchart, V. Towards enzymatic breakdown of complex plant xylan structures: State of the art. *Biotechnol. Adv.* **2016**, *34*, 1260–1274. (b) Lagaert, S.; Pollet, A.; Courtin, C. M.; Volckaert, G. β -Xylosidases and α -L-arabinofuranosidases: Accessory enzymes for arabinoxylan degradation. *Biotechnol. Adv.* **2014**, *32*, 316–332.

- 1
2
3
4
5
6
7
8
9
- 6) Wilkens, C.; Andersen, S.; Dumon, C.; Berrin, J.-G.; Svensson, B. GH62 arabinofuranosidases: Structure, function and applications. *Biotechnol. Adv.* **2017**, *35*, 792–804.
- 10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
- 7) (a) Mathew, S.; Karlsson, E. N.; Adlercreutz, P. Extraction of soluble arabinoxylan from enzymatically pretreated wheat bran and production of short xylo-oligosaccharides and arabinoxylooligosaccharides from arabinoxylan by glycoside hydrolase family 10 and 11 endoxylanases. *J. Biotechnol.* **2017**, *260*, 53–61. (b) Mechelke, M.; Koeck, D. E.; Broeker, J.; Roessler, B.; Krabichler, F.; Schwartz, W. H.; Zverlov, V. V.; Liebl, W. Characterization of the arabinoxylan-degrading machinery of the thermophilic bacterium *Herbinix hemicellulosilytica* – Six new xylanases, three arabinofuranosidases and one xylosidase. *J. Biotechnol.* **2017**, *257*, 122–130. (c) McCleary, B. V.; McKie, V. A.; Draga, A.; Rooney, E.; Mangan, D.; Larkin, J. Hydrolysis of wheat flour arabinoxylan, acid-debranched wheat flour arabinoxylan and arabino-xylo-oligosaccharides by β -xylanase, α -L-arabinofuranosidase and β -xylosidase. *Carbohydr. Res.* **2015**, *407*, 79–96.
- 36
37
38
39
40
41
42
43
44
45
46
- 8) (a) Utille, J.-P.; Jeacomine, I. Synthesis of a library of allyl α -L-arabinofuranosyl- α - or β -D-xylopyranosides; route to higher oligomers. *Carbohydr. Res.* **2007**, *342*, 2649–2656. (b) Hirsch, J.; Petráková, E.; Schraml, J. Stereoselective synthesis and ^{13}C -N.M.R. spectra of two isomeric methyl β -glycosides of trisaccharides related to arabinoxylan. *Carbohydr. Res.* **1984**, *131*, 219–226.
- 47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 9) (a) Senf, D.; Ruprecht, C.; de Kruijff, G. H. M.; Simonetti, S. O.; Schuhmacher, F.; Seeberger, P. H.; Pfrengle, F. Active Site Mapping of Xylan-Deconstructing Enzymes with Arabinoxylan Oligosaccharides Produced by Automated Glycan Assembly. *Chem. Eur. J.* **2017**, *23*, 3197–3205. (b) Schmidt, D.; Schuhmacher, F.; Geissner, A.; Seeberger, P. H.;

- Pfrenge, F. Automated Synthesis of Arabinoxylan-Oligosaccharides Enables Characterization of Antibodies that Recognize Plant Cell Wall Glycans. *Chem. Eur. J.* **2015**, *21*, 5709–5713.
- 10) Kinnaert, C.; Daugaard, M.; Nami, F.; Clausen, M. H. Chemical Synthesis of Oligosaccharides Related to the Cell Walls of Plants and Algae. *Chem. Rev.* **2017**, *117*, 11337–11405.
- 11) Takeo, K.; Ohguchi, Y.; Hasegawa, R.; Kitamura, S. Synthesis of (1→4)- β -D-xylo-oligosaccharides of dp 4-10 by a blockwise approach. *Carbohydr. Res.* **1995**, *278*, 301–313.
- 12) Pedersen, M. J.; Madsen, R.; Clausen, M. H. Iridium catalysis: reductive conversion of glucan to xylan. *Chem. Commun.* **2018**, *54*, 952–955.
- 13) Bonora, B.; Boos, I.; Clausen, M. H. Convergent strategy for the synthesis of *S*-linked oligoxylans. *Carbohydr. Res.* **2017**, *443-444*, 53–57.
- 14) (a) Zakharova, A. N.; Madsen, R.; Clausen, M. H. Synthesis of a Backbone Hexasaccharide Fragment of the Pectic Polysaccharide Rhamnogalacturonan I. *Org. Lett.* **2013**, *15*, 1826–1829. (b) Clausen, M. H.; Madsen, R. Synthesis of Hexasaccharide Fragments of Pectin. *Chem. Eur. J.* **2003**, *9*, 3821–3832. (c) Clausen, M. H.; Jørgensen, M. R.; Thorsen, J.; Madsen, R. A strategy for chemical synthesis of selectively methyl-esterified oligomers of galacturonic acid. *J. Chem. Soc., Perkin Trans. 1* **2001**, 543–551.
- 15) (a) Yang, B.; Yang, W.; Ramadan, S.; Huang, X. Pre-Activation-Based Stereoselective Glycosylations. *Eur. J. Org. Chem.* **2018**, 1075–1096. (b) Yang, W.; Yang, B.; Ramadan, S.; Huang, X. Preactivation-based chemoselective glycosylations: A powerful strategy for oligosaccharide assembly. *Beilstein J. Org. Chem.* **2017**, *13*, 2094–2114.

- 1
2
3 16) Huang, X.; Huang, L.; Wang, H.; Ye, X.-S. Iterative One-Pot Synthesis of
4 Oligosaccharides. *Angew. Chem. Int. Ed.* **2004**, *43*, 5221–5224.
5
6
7
8 17) For recent examples, see: (a) Zhang, H.; Shao, L.; Wang, X.; Zhang, Y.; Guo, Z.; Gao, J.
9 One-Pot Synthesis of the Repeating Unit of Type VII Group B *Streptococcus*
10 Polysaccharide and the Dimer. *Org. Lett.* **2019**, *21*, 2374–2377. (b) Zhang, Y.; Zhou, S.;
11 Wang, X.; Zhang, H.; Guo, Z.; Gao, J. A new method for α -specific glucosylation and its
12 application to the one-pot synthesis of a branched α -glucan. *Org. Chem. Front.* **2019**, *6*,
13 762–772. (c) Wang, P.; Lo Cascio F.; Gao, J.; Kayed, R.; Huang, X. Binding and
14 neurotoxicity mitigation of toxic tau oligomers by synthetic heparin like oligosaccharides.
15 *Chem. Commun.* **2018**, *54*, 10120–10123. (d) Xiong, C.; Feng, S.; Qiao, Y.; Guo, Z.; Gu, G.
16 Synthesis and Immunological Studies of Oligosaccharides that Consist of the Repeating
17 Unit of *Streptococcus pneumoniae* Serotype 3 Capsular Polysaccharide. *Chem. Eur. J.*
18 **2018**, *24*, 8205–8216.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33 18) Wu, Y.; Xiong, D.-C.; Chen, S.-C.; Wang, Y.-S.; Ye, X.-S. Total synthesis of
34 mycobacterial arabinogalactan containing 92 monosaccharide units. *Nat. Commun.* **2017**, *8*,
35 14851.
36
37
38
39
40 19) (a) Andersen, M. C. F.; Kračun, S. K.; Rydahl, M. G.; Willats, W. G. T.; Clausen, M. H.
41 Synthesis of β -1,4-Linked Galactan Side-Chains of Rhamnogalacturonan I. *Chem. Eur. J.*
42 **2016**, *22*, 11543–11548. (b) Dai, Y.; Yu, B. Total synthesis of astrosterioside A, an anti-
43 inflammatory asterosaponin. *Chem. Commun.* **2015**, *51*, 13826–13829.
44
45
46
47
48
49 20) López, R.; Fernández-Mayorales, A. Enzymatic β -Galactosidation of Modified
50 Monosaccharides: Study of the Enzyme Selectivity for the Acceptor and Its Application to
51 the Synthesis of Disaccharides. *J. Org. Chem.* **1994**, *59*, 737–745.
52
53
54
55
56
57
58
59
60

- 1
2
3
4 21) Thorsheim, K.; Siegbahn, A.; Johnsson, R. E.; Stålbrand, H.; Manner, S.; Widmalm, G.;
5 Ellervik, U. Chemistry of xylopyranosides. *Carbohydr. Res.* **2015**, *418*, 65–88.
6
7
8 22) Rio, S.; Beau, J.-M.; Jacquinet, J.-C. Synthesis of glycopeptides from the carbohydrate-
9 protein linkage region of proteoglycans. *Carbohydr. Res.* **1991**, *219*, 71–90.
10
11
12 23) Yang, B.; Yoshida, K.; Yin, Z.; Dai, H.; Kavunja, H.; El-Dakdouki, M. H.; Sungsuwan,
13 S.; Dulaney, S. B.; Huang, X. Chemical Synthesis of a Heparan Sulfate Glycopeptide:
14
15
16
17
18
19
20 24) Andersen, S. M.; Heuckendorff, M.; Jensen, H. J. 3-(Dimethylamino)-1-propylamine: A
21 Cheap and Versatile Reagent for Removal of Byproducts in Carbohydrate Chemistry. *Org.*
22
23
24
25
26
27 25) This procedure has previously been used for regioselective benzylation of ethyl 1-
28 thiohexopyranosides, see: Garegg, P. J.; Kvarström, I.; Niklasson, A.; Niklasson, G.;
29
30
31
32
33
34
35
36 26) Martichonok, V.; Whitesides, G. M. Stereoselective α -Sialylation with Sialyl Xanthate
37 and Phenylsulfenyl Triflate as a Promoter. *J. Org. Chem.* **1996**, *61*, 1702–1706.
38
39
40
41 27) Tatai, J.; Fügedi, P. A New, Powerful Glycosylation Method: Activation of
42
43
44
45 28) Crich, D.; Smith, M. 1-Benzenesulfinyl Piperidine/Trifluoromethanesulfonic Anhydride:
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 29) Crich, D.; Cai, F.; Yang, F. A stable, commercially available sulfenyl chloride for the
5 activation of thioglycosides in conjunction with silver trifluoromethanesulfonate.
6
7 *Carbohydr. Res.* **2008**, *343*, 1858–1862.
8
9
- 10 30) Zeng, Y.; Wang, Z.; Whitfield, D.; Huang, X. Installation of Electron-Donating Protective
11 Groups, a Strategy for Glycosylating Unreactive Thioglycosyl Acceptors using the
12 Preactivation-Based Glycosylation Method. *J. Org. Chem.* **2008**, *73*, 7952–7962.
13
14
- 15 31) Christensen, H. M.; Oscarson, S.; Jensen, H. H. Common side reactions of the glycosyl
16 donor in chemical glycosylation. *Carbohydr. Res.* **2015**, *408*, 51–95.
17
18
- 19 32) For a previous example of aglycon transfer during the pre-activation protocol, see: Huang,
20 L.; Wang, Z.; Li, X.; Ye, X.-s.; Huang, X. Iterative one-pot syntheses of chitotetroses.
21
22 *Carbohydr. Res.* **2006**, *341*, 1669–1679.
23
24
25
26
27
- 28 33) Crich, D.; Dai, Z.; Gastaldi, S. On the Role Neighboring Group Participation and Ortho
29 Esters in β -Xylosylation: ^{13}C NMR Observation of a Bridging 2-Phenyl-1,3-dioxalenium
30 Ion. *J. Org. Chem.* **1999**, *64*, 5224–5229.
31
32
33
34
- 35 34) Crich, D.; Smith, M.; Yao, Q.; Picione, J. 2,4,6-Tri-*tert*-butylpyrimidine (TTBP): A Cost
36 Effective, Readily Available Alternative to the Hindered Base 2,6-Di-*tert*-butylpyridine and
37 its 4-Substituted Derivatives in Glycosylation and Other Reactions. *Synthesis* **2001**,
38 323–326.
39
40
41
42
43
44
- 45 35) van Rijssel, E. R.; Goumans, T. P. M.; Lodder, G.; Overkleeft, H. S.; van der Marel, G.
46 A.; Codée, J. D. C. Chiral Pyrroline-Based Ugi-Three-Component Reactions Are under
47 Kinetic Control. *Org. Lett.* **2013**, *15*, 3026–3029.
48
49
50
- 51 36) Bock, K.; Pedersen, C. A Study of ^{13}C H Coupling Constants in Pentopyranoses and Some
52 of their Derivatives. *Acta Chem. Scand.* **1975**, *B29*, 258–264.
53
54
55
56
57
58
59
60

- 1
2
3
4 37) Mizutani, K.; Kasai, R.; Nakamura, M.; Tanaka, O. N.M.R. Spectral Study of α - and β -L-
5
6 Arabinofuranosides. *Carbohydr. Res.* **1989**, *185*, 27–38.
7
8 38) Motawia, M. S.; Marcussen, J.; Møller, B. L. A General Method Based on the Use of *N*-
9
10 Bromosuccinimide for Removal of the Thiophenyl Group at the Anomeric Position to
11
12 Generate A Reducing Sugar with the Original Protecting Groups Still Present. *J.*
13
14 *Carbohydr. Chem.* **1995**, *14*, 1279–1294.
15
16
17 39) For Zemplén deprotection of perbenzoylated gluco-, manno- and galactopyranose in
18
19 quantitative yields, see: Ren, B.; Wang, M.; Liu, J.; Ge, J.; Zhang, X.; Dong, H. Zemplén
20
21 transesterification: a name reaction that has misled us for 90 years. *Green Chem.* **2015**, *17*,
22
23 1390–1394.
24
25
26 40) The term peeling originates from the observed degradation of polysaccharides during the
27
28 alkaline pulping and bleaching of wood, see: (a) Berglund, J.; Azhar, S.; Lawoko, M.;
29
30 Lindström, M.; Vilaplana, F.; Wohler, J.; Henriksson, G. The structure of
31
32 galactoglucomannan impacts the degradation under alkaline conditions. *Cellulose* **2019**, *26*,
33
34 2155–2175. (b) Wang, Y.; Azhar, S.; Lindström, M. E.; Henriksson, G. Stabilization of
35
36 Polysaccharides During Alkaline Pre-Treatment of Wood Combined with Enzyme-
37
38 Supported Extractions in a Biorefinery. *J. Wood Chem. Technol.* **2015**, *35*, 91–101.
39
40
41 41) Andersen, M. C. F.; Boos, I.; Ruprecht, C.; Willats, W. G. T.; Pfrengle, F.; Clausen, M. H.
42
43 Synthesis and Application of Branched Type II Arabinogalactans. *J. Org. Chem.* **2017**, *82*,
44
45 12066–12084.
46
47
48 42) Pedersen, D. S.; Rosenbohm, C. Dry Column Vacuum Chromatography. *Synthesis* **2001**,
49
50 2431–2434.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 43) Kongkathip, B.; Kongkathip, N.; Rujirawanich, J. New Strategy for Synthesis of the
4
5 Disaccharide Moiety of the Highly Potent Anticancer Natural Product OSW-1. *Synth.*
6
7 *Commun.* **2014**, *44*, 2248–2255.
8
9
10 44) Crich, D.; Dai, Z. Direct Synthesis of β -Mannosides. Synthesis of β -D-Xyl-(1 \rightarrow 2)- β -D-
11
12 Man-(1 \rightarrow 4)- α -D-Glc-OMe: a Trisaccharide Component of the *Hyriopsis schlegelii*
13
14 Glycosphingolipid. Formation of an Orthoester from a Xylopyranosyl Sulfoxide.
15
16 *Tetrahedron* **1999**, *55*, 1569–1580.
17
18
19 45) Phanumartwiwath, A.; Hornsby, T. W.; Jamalis, J.; Bailey, C. D.; Willis, C. L. Silyl
20
21 Migrations in D-Xylose Derivatives: Total Synthesis of a Marine Quinoline Alkaloid. *Org.*
22
23 *Lett.* **2013**, *15*, 5734–5737.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60