



Agglomeration in Fluidized Bed Gasification of Biomass

Zhao, Liyan; Lin, Weigang; Wu, Hao; Dam-Johansen, Kim

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Zhao, L., Lin, W., Wu, H., & Dam-Johansen, K. (2019). *Agglomeration in Fluidized Bed Gasification of Biomass*. Abstract from Fluidization XVI, Guilin, China.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Agglomeration in Fluidized Bed Gasification of Biomass

Liyan Zhao, Weigang Lin, Hao Wu and Kim Dam-Johansen

Department of Chemical and Biochemical Engineering, Technical University of Denmark,
Kongens Lyngby, 2800, Denmark

Abstract

Fluidized beds (FB) are widely applied to gasification of biomass due to its advantages of high fuel flexibility and conversion. However, bed agglomeration induced by the high content of alkali compounds can have strong impact on the operation of a FB gasifier and may lead to defluidization and unscheduled plant shutdown in severe cases. Agglomeration at high temperature, such as combustion and gasification, is mainly attributed to the presence of molten phase, i.e. the low melting point alkali containing compounds originated from biomass. During gasification, the alkali species in biomass can be transformed into gaseous phase, such as KCl and KOH vapor, and condensed phase, such as K_2SO_4 aerosols and K-silicates particles, depending on the local conditions. The interactions of the alkali species with bed materials (mainly SiO_2) may result in the formation of molten phase, causing agglomeration. However, the agglomeration tendencies and mechanisms during biomass gasification with different gasification agents are still not fully understood.

In this work, the agglomeration behaviors of different types of biomass during gasification are studied in a lab-scale fluidized bed reactor. The impact of the composition of biomass, e.g. types of biomass, and the operational conditions, such as temperature and gasification agents (steam and CO_2), are studied. The different agglomeration mechanisms are revealed based on SEM-EDX analyses of the agglomerate samples and chemical equilibrium calculations.