



INTEGRAL detection of the neutron-star X-ray transient Swift J1734.5-3027

Kuulkers, E.; Eckert, D.; Ferrigno, C.; Chenevez, J.; Alfonso-Garzon, J.; Beckmann, V.; Bird, A. J.; Brandt, S.; Del Santo, M.; Domingo, A.

Total number of authors:
19

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Kuulkers, E., Eckert, D., Ferrigno, C., Chenevez, J., Alfonso-Garzon, J., Beckmann, V., Bird, A. J., Brandt, S., Del Santo, M., Domingo, A., Ebisawa, K., Jonker, P. G., Kretschmar, P., Markwardt, C. B., Oosterbroek, T., Paizis, A., Pottschmidt, K., Sanchez-Fernandez, C., & Wijnands, R. (2013, Sept 4). INTEGRAL detection of the neutron-star X-ray transient Swift J1734.5-3027. The Astronomer's telegram No. ATel #5361 <http://www.astronomerstelegam.org/?read=5361>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Outside
[GCN](#)
[IAUCs](#)


Other
 MacOS: [Dashboard Widget](#)
 Follow ATel on [Twitter](#)
[ATELstream](#)
[ATel Community Site](#)

The Astronomer's Telegram

[Post a New Telegram](#) | [Search](#) | [Information](#)
[Telegram Index](#)
[Obtain Credential To Post](#) | [RSS Feeds](#) | [Email Settings](#)

Present Time: 19 Dec 2013; 10:25 UT

This space for free for your conference.



The Structure and Signals of Neutron Stars, from Birth to Death

[[Previous](#) | [Next](#) | [ADS](#)]

INTEGRAL detection of the neutron-star X-ray transient Swift J1734.5-3027

ATel #5361; [E. Kuulkers \(ESA/ESAC, Spain\)](#), [D. Eckert, C. Ferrigno \(ISDC, University of Geneva\)](#), [J. Chenevez \(DTU Space, Denmark\)](#), [J. Alfonso-Garzon \(CAB/INTA-CSIC, Spain\)](#), [V. Beckmann \(APC, France\)](#), [A. J. Bird \(Southampton, UK\)](#), [S. Brandt \(DTU Space, Denmark\)](#), [M. Del Santo \(INAF/IAPS Roma, Italy\)](#), [A. Domingo \(CAB/INTA-CSIC, Spain\)](#), [K. Ebisawa \(U of Tokyo, JAXA/ISAS, Japan\)](#), [P. G. Jonker \(SRON/CfA/RU\)](#), [P. Kretschmar \(ESA/ESAC, Spain\)](#), [C. B. Markwardt \(NASA/GSFC, USA\)](#), [T. Oosterbroek \(ESA/ESTEC, The Netherlands\)](#), [A. Paizis \(INAF-IASF Milano, Italy\)](#), [K. Pottschmidt \(CRESST/UMBC, NASA/GSFC, USA\)](#), [C. Sanchez-Fernandez \(ESA/ESAC, Spain\)](#) & [R. Wijnands \(UvA, The Netherlands\)](#)

on 4 Sep 2013; 16:12 UT

Credential Certification: Erik Kuulkers (Erik.Kuulkers@sciops.esa.int)

Subjects: X-ray, Binary, Neutron Star, Transient, Variables

Referred to by ATel #: [5448](#), [5646](#)

Tweet

Recommend

INTEGRAL Galactic bulge monitoring (see ATel #[438](#)) observations performed between UT 2013 September 3 20:43 and September 4 00:25 clearly reveal the newly detected neutron star X-ray transient Swift J1734.5-3027 (ATel #[5354](#), GCN #[15157](#), #[15172](#)). The JEM-X and IBIS/ISGRI flux averages during the observations are 14 ± 3 mCrab (3-10 keV), 19 ± 6 mCrab (10-25 keV), 23 ± 2 mCrab (18-40 keV), and 17 ± 2 mCrab (40-100 keV). The average IBIS/ISGRI spectrum, with an effective exposure of about 9 ksec, can be well (reduced $\chi^2=0.8$ for 9 degrees of freedom) fitted by a power-law with index 2.1 ± 0.4 , with a 20-100 keV flux of about 3.7×10^{-10} erg/s/cm².

A re-analysis of the previous monitoring observations, performed between August 31 20:38 and September 1 00:19, shows that the source was already marginally detected by JEM-X and IBIS/ISGRI with fluxes of 7 ± 2 mCrab (3-10 keV; 4 sigma significance) and 14 ± 2 mCrab (18-40 keV; 6.6 sigma significance), respectively. The 10-25 keV JEM-X 6-sigma upper limit is about 4 mCrab, while the 40-100 keV IBIS/ISGRI 3-sigma upper limit is about 8 mCrab. The average IBIS/ISGRI spectrum for these observations, with an effective exposure of about 6.5 ksec, can be described by a power-law with index 2.3 ± 0.8 and a 20-100 keV flux of about 2.3×10^{-10} erg/s/cm² (reduced $\chi^2=1.5$ for 7 degrees of freedom).

We encourage further observations of this new X-ray transient, which showed a long Type I X-ray burst, possibly a superburst (ATel #[5354](#)), near the start of its current outburst, i.e., about 9 hours after the end of the INTEGRAL Galactic bulge observations on September 1.

[[Telegram Index](#)]

R. E. Rutledge, Editor-in-Chief

Derek Fox, Editor

Mansi M. Kasliwal, Co-Editor

rrutledge@astronomerstelegam.org

dfox@astronomerstelegam.org

mansi@astronomerstelegam.org

Related

5646 [Swift J1734.5-3027: a bright burst in March 2013 detected in the long term BAT light curve.](#)

5448 [IGR J17344-3023 = Swift J1734.5-3027](#)

5447 [IGR J17344-3023: a new X-ray transient detected by INTEGRAL/JEM-X](#)

5361 [INTEGRAL detection of the neutron-star X-ray transient Swift J1734.5-3027](#)

5354 [Swift J1734.5-3027: Swift discovery of a possible new superbursting transient](#)