



A new spectrum model for the atmospheric crosswind component applicable from mesoscales to microscales

Larsén, Xiaoli G.; Larsen, Søren E.; L. Petersen, Erik; Mikkelsen, Torben K.

Publication date:
2020

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Larsén, X. G., Larsen, S. E., L. Petersen, E., & Mikkelsen, T. K. (2020). A new spectrum model for the atmospheric crosswind component applicable from mesoscales to microscales. Sound/Visual production (digital)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A new spectrum model for the atmospheric crosswind component applicable from mesoscales to microscales

by **Xiaoli G. Larsén**, Søren E. Larsen, Erik L. Petersen and Torben K. Mikkelsen
Technical University of Denmark, Wind Energy Department, xgal@dtu.dk

The issue:



Fig. 1. Overview of wind farms (4coffshore.com)

Meandering effect is described through classical boundary-layer turbulence theories that are valid for $t < 1$ h, $l < a$ few km.

How is meandering effect over modern wind farm clusters?

The observation:




Fig. 2. Mean spectrum of u and v at 80 m from two years of data of stationary days from Høvsøre. Sonic data with 1-day length were used for calculation of each spectrum.

The red curves are the new spectral models for u and v that cover meso and micro scales.

This new model has also shown to be valid over 5 years of data from Østerild, from 37 m to 241 m.

The approach:

- Spectral model development for u and v over micro and mesoscale.
- u -spectrum from Larsén et al. (2016, 2019)

$$fS_u(f) = \text{Eq. 1a (or Eq. 2a)} + \text{Eq. 3}$$

Note:

Eq. 1a $fS_u(f) = \frac{51u_*^2 n}{(1 + 33n)^{5/3}}$

Eq. 1b $fS_v(f) = \frac{8.5u_*^2 n}{(1 + 9.5n)^{5/3}}$

Boundary-layer model by Kaimal & Finnigan 1994

Eq. 2a $fS_u(f) = \frac{0.5a_u u_*^2 n / n_{l,u}}{(1 + n/n_{l,u})(1 + n/n_{u,u})^{2/3}}$

Eq. 2b $fS_v(f) = \frac{0.5a_v u_*^2 n / n_{l,v}}{(1 + n/n_{l,v})(1 + n/n_{u,v})^{2/3}}$

Boundary-layer model by Mikkelsen et al. 2017

Eq. 3 $fS_{lu}(f) = a_1 f^{-2/3} + a_2 f^{-2}$

Mesoscale model by Larsén et al. 2013

- v -spectrum derived here:

$$fS_v(f) = \begin{cases} \text{Eq. 1b (or Eq. 2b)} & \text{for } f > f_1 \\ \text{Constant} & \text{for } f_2 \leq f \leq f_1 \\ \text{Eq. 3} & \text{for } f < f_2 \end{cases}$$

Refer to Fig. 2

The approach:

- Time series (1-day and 5-day for a year) are generated for u and v using the spectral models:
 - Boundary-layer model Eq. 1 & 2
 - The new model combining meso and boundary-layer models
- Statistics of modelled direction are compared with observations

The results:




Fig. 3. The distribution of direction deviation with groups of observation data that satisfy: (left) the largest direction difference is 15° and (right) 60° , with data length (DL) from 1 h to 1 day.

The boundary-layer model (Eq. 1 and 2, dashed curves) seems to capture well this distribution when the largest direction change is less than 15° (stationary), but significantly underestimate the directional change for non-stationary situations.

The results:




Fig. 4. Deviation of direction distribution, from measurements (gray curves) and from models. The new spectral model (blue curve) provides much improved estimation for non-stationary condition (an example of data length DL= 5 days) in comparison with the boundary-layer model Eq. 2 (red dashed curve).

The conclusion:

- We defined a mesoscale version of the lateral, v -spectrum, extending the description of v -variability to mesoscale: the meso-BL-model.
- The new meso-BL-models of u and v are used to simulate the climatology of wind direction.
- This method pertains at least to the western Denmark and North Sea regions, but can be applied in other regions with calibrations for the spectral models using measurements.
- At the scale of modern wind farm clusters, stationary conditions are challenging to meet. Meandering calculation needs to take large scale variability of u and v into account.

References & Acknowledgements:

Kaimal J, Finnigan J (1994) Atmospheric boundary layer flows. Oxford University Press, New York, 289 pp

Larsén XG, Vincent CL, Larsen SE (2013) Spectral structure of the mesoscale winds over the water. Q J R Meteorol Soc 139:685700, DOI DOI:10.1002/qj.2003

Larsén XG, Larsen SE, Petersen EL (2016) Full-scale spectrum of boundary-layer winds. Boundary-Layer Meteorol 159:349371

Larsén XG, Larsen SE, Petersen EL, Mikkelsen TK (2019a) Turbulence Characteristics of Wind-Speed Fluctuations in the Presence of Open Cells: A Case Study. Boundary-Layer Meteorol 171:191212

Mikkelsen TK, Larsen SE, Jørgensen HE, Astrup P, Larsén XG (2017) Scaling of turbulence spectra measured in strong shear flow near the earth surface. Phys Scr 92:124,002

The authors acknowledge part of the support from the ForskEL project OffshoreWake (PSO-12521).