



INTEGRAL and Swift detection of high energy emission from Swift J1749.4-2807

Lucia, Pavan; Chenevez, J.; Brandt, S.; Bozzo, E.; Kuulkers, E.; Alfonso-Garzon, J.; Beckmann, V.; Bird, T.; Courvoisier, T.; Domingo, A.

Total number of authors:
18

Publication date:
2010

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Lucia, P., Chenevez, J., Brandt, S., Bozzo, E., Kuulkers, E., Alfonso-Garzon, J., Beckmann, V., Bird, T., Courvoisier, T., Domingo, A., Ebisawa, K., Jonker, P., Kretschmar, P., Markwardt, C., Oosterbroek, T., Paizis, A., Sanchez-Fernandez, C., & Wijnands, R. (2010, Apr 12). INTEGRAL and Swift detection of high energy emission from Swift J1749.4-2807. The Astronomer's telegram No. ATel #2548
<http://www.astronomerstelegam.org/?read=2548>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

INTEGRAL and Swift detection of high energy emission from Swift J1749.4-2807

ATel #2548; [L. Pavan \(ISDC, Switzerland\)](#), [J. Chenevez \(DTU space, Denmark\)](#), [E. Bozzo \(ISDC, Switzerland\)](#), [E. Kuulkers \(ESA/ESAC, Spain\)](#), [J. Alfonso-Garzon \(LAEX-CAB/INTA-CSIC, Spain\)](#), [V. Beckmann \(APC, France\)](#), [T. Bird \(Southampton, UK\)](#), [S. Brandt \(DTU/NSI, Denmark\)](#), [T. Courvoisier \(ISDC, Switzerland\)](#), [A. Domingo \(LAEX-CAB/INTA-CSIC, Spain\)](#), [K. Ebisawa \(ISAS, Japan\)](#), [P. Jonker \(SRON, The Netherlands\)](#), [P. Kretschmar \(ESAC/ESA, Spain\)](#), [C. Markwardt \(GSFC, USA\)](#), [T. Oosterbroek \(ESTEC/ESA, The Netherlands\)](#), [A. Paizis \(INAF-IASF, Italy\)](#), [C. Sanchez-Fernandez \(ESAC/ESA, Spain\)](#), [R. Wijnands \(UvA, The Netherlands\)](#)

on 12 Apr 2010; 21:41 UT

Distributed as an Instant Email Notice Request For Observations

Password Certification: [Nami Mowlavi \(Nami.Mowlavi@obs.unige.ch\)](mailto:Nami.Mowlavi@obs.unige.ch)

Subjects: X-ray, Gamma Ray, Request for Observations, Neutron Star, Transient
Referred to by ATel #: [2561](#), [2565](#), [2569](#), [2576](#), [2579](#), [2585](#)

During the observations of the Galactic bulge, performed from 2010-04-10 22:00:00 to 2010-04-11 01:41:36 (UTC), INTEGRAL detected high energy emission from a position coincident with the burst-only source Swift J1749.4-2807 (GRB060602B).

The source is detected at 8 sigma in the IBIS/ISGRI mosaic in the energy band 20-40 keV. We used all data of this observation to extract the ISGRI spectrum. The spectrum could be equally well fitted by using a power law (photon index 2.5 ± 0.9) or a blackbody model with $kT = 7.9 (+2.6, -1.9)$ keV. The estimated flux in the 20-40 keV energy band was of $1.3E-10$ erg/cm²/s (corresponding to 17mCrab, effective exposure time 8.4 ks).

The ISGRI lightcurve did not show any clear evidence for a short burst with a fast rise and decay time, similar to that observed previously from this source (Wijnands et al. 2009, MNRAS, 393, 126; Campana, 2009, ApJ 699 1144).

The source was also detected by both JEMX1 and JEMX2 at a significance of 4.5 and 3.3 sigma in the energy bands 3-10 keV and 10-25 keV, respectively (effective exposure time 6.5 ks). Given the low significance of the detection, we roughly estimated a source X-ray flux of ~10-20 mCrab.

Previous observations of the region of Swift J1749.4-2807 were performed by INTEGRAL from 2010-04-07 13:31:59 to 2010-04-07 17:13:34 for an exposure time of 15 ks (IBIS) and 9 ks (JEM-X). On that occasion the source was not detected. We estimated a 5 sigma upper limit on the source X-ray flux of 10(17) mCrab in the 20-40(40-100) keV energy bands and 5(2) mCrab in the 3-10(11-25) keV energy bands.

The source activity has been confirmed using a Swift ToO observation performed from 2010-04-11 23:41:10 to 2010-04-12 04:48:48 (UTC). The Swift lightcurve does not present any evidence for a fast rise and decay. After clearing of the data for pile-up, the spectrum is well fitted by a powerlaw model with photon index 2.6 ± 0.2 and $NH = 3.4 \pm 0.4 \times 10^{22} \text{ cm}^{-2}$. The inferred model flux in the 0.5-10 keV band, not corrected for absorption, is $3.0 \times 10^{-10} \text{ erg/cm}^2/\text{s}$. These values are in agreement with those reported in Campana, 2009, soon after the burst (powerlaw index 2.7, $NH = 4.0 \times 10^{22} \text{ cm}^{-2}$).

Further INTEGRAL observations of the Galactic bulge will be performed from 2010-04-13 15:00:00 to 2010-04-13 18:41:45 (UTC).

We are grateful to the Swift PI, Neil Gehrels, and the Swift mission operations team for their prompt scheduling of the Swift observations.