



## Meat consumption, stratified by dietary quality, and risk of heart disease

Møller, Sanne Pagh; Mejborn, Heddie; Biloft-Jensen, Anja Pia; Christensen, Anne Illemann; Thygesen, Lau Caspar

*Published in:*  
British Journal of Nutrition

*Link to article, DOI:*  
[10.1017/S0007114521000623](https://doi.org/10.1017/S0007114521000623)

*Publication date:*  
2021

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Møller, S. P., Mejborn, H., Biloft-Jensen, A. P., Christensen, A. I., & Thygesen, L. C. (Accepted/In press). Meat consumption, stratified by dietary quality, and risk of heart disease. *British Journal of Nutrition*. <https://doi.org/10.1017/S0007114521000623>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Meat consumption, stratified by dietary quality, and risk of heart disease

Sanne Pagh Møller<sup>1</sup>, Heddie Mejborn<sup>2</sup>, Anja Biloft-Jensen<sup>2</sup>, Anne Illemann Christensen<sup>1</sup>, Lau Caspar Thygesen<sup>1</sup>

<sup>1</sup> National Institute of Public Health, University of Southern Denmark

<sup>2</sup> National Food Institute, Technical University of Denmark

Short title: Meat intake, diet quality and heart disease

Corresponding author:

Sanne Pagh Møller  
sapm@sdu.dk

National Institute of Public Health  
University of Southern Denmark  
Studivstræde 6  
1455 Copenhagen  
Denmark

Keywords: Meat; dietary quality; ischemic heart disease; longitudinal studies; nutrition surveys



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114521000623

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

**Abstract**

Associations between meat consumption and heart disease have been assessed in several studies but it has been suggested that other dietary factors influence these associations. The aim of this study was to assess whether meat consumption is associated with ischemic heart disease (IHD), and if the association is modified by dietary quality. The analyses were based on the cohort of adult participants in The Danish National Survey on Diet and Physical Activity in 2000-2002, 2003-2008, and 2011-2013. From these surveys, information on meat consumption and dietary quality were extracted. The cohort was followed in national registers to identify incident IHD. Associations were estimated using Cox regression analyses adjusting for sociodemographic and lifestyle factors. Analyses of associations between meat consumption and IHD stratified by dietary quality were subsequently evaluated. Among the 8,007 participants, the median follow-up was 9.8 years and 439 cases of IHD were recorded. The results suggested a trend between consumption increments of 100g/day of red meat (HR=1.23; 95%CI:0.99-1.53) or of 50 g/day of processed meat (HR=1.09; 95%CI:0.93-1.29) and higher risk of IHD. The trends were, however, not statistically significant. Stratification by dietary quality did not suggest that associations between meat consumption and risk of IHD were modified by dietary quality. This population-based cohort study with detailed dietary information, suggested a trend with higher meat consumption being associated with higher risk of IHD, but the association was not statistically significant. Results did not indicate that dietary quality modifies such associations.

## Introduction

The associations between red and processed meat consumption and heart disease incidence and mortality have been assessed in several studies and meta-analyses (1, 2). Two meta-analyses on red meat consumption and risk of coronary heart disease or ischemic heart disease (IHD) found no associations (1, 2). However, two prospective studies published after the meta-analyses were performed found an 18-19% increased risk for every consumption increment of 100 g/day (3, 4). A meta-analysis studying heart disease mortality and not just disease found that increments of 100 g/day in red meat consumption was associated with 24% increased risk of cardiovascular mortality (2).

A meta-analysis on processed meat consumption and cardiovascular events found that every 50 g/day increment in consumption was associated with 44 % increased risk of cardiovascular events (1). Another meta-analysis found that increments of 50 g/day in consumption of processed meat was associated with 24% increased risk of cardiovascular mortality. However, this analysis did not find associations with IHD mortality (2).

Confounding factors such as sex, age, smoking, physical activity, and disease history are routinely included in analyses on associations between meat consumption and risk of heart disease. However, it has been suggested that other dietary factors including fruit, vegetable, and fibre intake may influence these associations (5). Therefore, other dietary factors should be considered when estimating the causal association between meat consumption and heart disease risk. Previous studies have accommodated for this confounding by adjusting analyses for other dietary factors. Yet, in order to understand the influence of other dietary factors on associations between meat consumption and heart disease risk, dietary confounding is better assessed when stratifying the association by other dietary factors. The present study will allow for the identification of different associations between meat consumption and heart disease in participants with a healthy diet compared to participants with an unhealthy diet. A study conducted in Sweden investigated if an association between red meat consumption and cardiovascular mortality differed between those with high and low consumption of fruit and vegetables. In this study, the associations did not differ between those with high and low consumption of fruit and vegetables (6). Dietary quality is, however, composed of a range of

factors beyond fruit and vegetable consumption, and detailed dietary information is required to estimate the dietary quality. In the present study, detailed information on a range of dietary factors was obtained from a cohort of more than 10,000 individuals and information on IHD was obtained from national health registers. Therefore, the aim of the present study was to assess whether meat consumption is associated with IHD, and subsequently assess if the association is modified by dietary quality.

## Methods

### *The Danish National Survey on Diet and Physical Activity (DANSDA)*

The analyses were based on information from participants in DANSDA 2000-2002, 2003-2008, and 2011-2013. The study sample was a random sample drawn from the Danish Civil Registration System and comprised of non-institutionalised Danish citizens. The response proportions in DANSDA 2000-2002, 2003-2008, and 2011-2013 were 53.3 %, 53.3 %, and 54.4 %, respectively. Only participants aged 15-75 years at baseline were eligible for inclusion in the study (n=9,848). In order to study incident disease, individuals were excluded if they had been diagnosed with IHD any time before baseline. The same strategy was applied for analyses on acute coronary syndrome (ACS). Thus 8,007 participants were included in analyses on IHD, and 8,198 were included in analyses on ACS.

Information on food intake was obtained via self-administered, quantitative seven-day pre-coded food diaries. The data collection has previously been described (7). The detailed dietary registration allowed for a subdivision into red and processed meat and for constructing a diet quality index based on the Danish dietary guidelines. Information on meat consumption, dietary quality, energy intake, alcohol energy intake, BMI (<18.5; 18.5-24.9; 25-29.9;  $\geq 30$ ), smoking (never; former; current), and leisure physical activity (none; light; moderate/hard) were extracted or generated from the surveys. Red meat was defined as unprocessed muscle tissue from mammals such as beef, veal, pork, and lamb. In this population red meat mainly consisted of beef, veal, and pork. A small intake of unprocessed edible offal, e.g. liver and heart, was included. The meat could be minced and/or frozen. Processed meat was defined as red meat or poultry that undergo a

transformation and contain approved ingredients and may be subject to some form of preservation including smoking, drying, curing, or fermentation.

To assess the dietary quality, the diet of each participant was given a dietary quality score (8, 9). This was based on how well their diet complied with five of the official Danish dietary guidelines on fruit and vegetables, fish, whole grain, saturated fatty acids, and added sugars. For each dietary guideline, a score between 0 and 1 was given by dividing the dietary content of the actual component with the recommended dietary content. For example, the recommended intake of fruit and vegetables is 600 g per day. Thus, if a participant ate 200 g of fruit and vegetables per day, the score was  $200/600=0.33$ . For saturated fatty acids and added sugars, the scores were reverse. For example, for a participant with an energy intake of 15E% from saturated fatty acids where the recommendation is <10E%, the score was  $1-(15-9.999)/9.999=0.50$ . Scores >1 were truncated at 1 and scores < 0 were truncated at 0. The scores were added to yield the total score between 0 and 5. For those participants who had answered more than one survey (89 persons), information from their first survey was included.

### ***Register-based information***

Events of IHD and ACS were identified through linkage with information from registers using the unique personal identification number (10). The outcome IHD was based on information from the National Patient Register (11) on primary and secondary diagnoses and from the Register of Causes of Death (12) on underlying cause of death (ICD-8: 410-414; ICD-10: I20-I25). Information on ACS was also based on information from the National Patient Register on primary and secondary diagnoses and from the Register of Causes of Death on underlying cause of death (ICD-8: 410 and 427.27; ICD-10: I20.0 and I21).

Information on age, sex, ethnicity (Danish, Western non-Danish, Non-Western), and emigration were obtained from the Danish Civil Registration System (10). Educational attainment was based on information from the Population's Educational Register (13) and classified into short (primary school), medium (high school or vocational school), and long education (higher education). Information on diabetes up to 10 years before baseline was obtained from the National Patient Register (ICD-8: 249-250; ICD-10: E10-E14).

## *Analyses*

The cohort was followed from baseline (date of survey interview) or from age 37 whichever came last. Delayed entry into the analyses was applied because the outcomes were almost absent among participants younger than 37 years. Follow-up ended at first diagnosis or death due to the studied outcome or at emigration, death, or end of follow-up (31 December 2017), whichever came first.

Missing information on country of origin (0.01%) was imputed as Danish origin, missing educational level (1.5%) was imputed as short education, missing BMI (1.0%) was imputed as normal BMI (18.5-24.9), missing smoking status (1.1%) was imputed as never smoker, and missing information on physical activity (0.4%) was imputed as moderate/hard. Apart from educational level, these imputed values were chosen as they were the most prevalent among the non-missing values. For educational level, short education was chosen, because missing information on this is more common among persons with short education (13).

## *Associations between meat consumption and outcomes*

The categorisations of meat consumption were based on the observed quartiles. For analyses of IHD, the measures of meat consumption were categorised in three groups (lower quartile; the two middle quartiles; upper quartile). Due to the small number of cases with ACS, the measures of meat consumption were categorised in two groups (below median; median and above) in these analyses. The same methods were applied for categorisation of dietary quality. For the continuous measures of meat consumption, red meat was included as increment of 100 g/day, and processed meat was included as increment of 50 g/day.

The associations between meat consumption and the studied outcomes were estimated using Cox regression analyses with age as the underlying timescale. The different types of meat consumption were included in different regression models. The analyses were adjusted for sex, educational attainment (the year before baseline), ethnicity, smoking, physical activity, alcohol, BMI, total energy intake, and history of diabetes. To test if the associations between meat consumption and the outcomes were best represented by non-linear effects, quadratic, and cubic terms were included in the models. However, non-linear effects were non-significant, so meat consumption was included linearly.

To evaluate if the assumption of proportional hazards was fulfilled, the Schoenfeld Residuals were estimated. A linear regression model tested whether these residuals were correlated with the underlying timescale (age). These analyses indicated that the assumption was fulfilled. Further, the log-negative-log survival curves for each of the exposure and outcome variables were visually inspected and did not indicate serious violation of the assumption.

### *Associations between meat consumption and outcomes stratified by dietary quality*

The statistical significance of a modifying effect of dietary quality was tested by including both meat consumption and dietary quality as separate main effects and as an interaction term with each other. These tests were performed both on analyses with meat consumption as a categorical and a continuous variable. The tests on continuous variables were considered the main tests, as they require the least statistical power.

All analyses were performed using SAS 9.4 (SAS Institute Inc, Cary, North Carolina, USA).

### *Ethics statement*

Ethical permission for scientific studies carried out in Denmark is necessary only when they include biological samples, such as blood or tissue, or information from medical records. Therefore, ethical permission was not necessary for this study.

## **Results**

Among the 8,007 participants included in the analyses on IHD, the median follow-up was 9.8 years, during which 439 cases of IHD were recorded. Among the 8,198 participants included in the analyses on ACS, the median follow-up was 10 years, during which 242 cases of ACS were recorded.

The baseline characteristics of the study population stratified by red meat consumption and dietary quality are shown in table 1. The characteristics varied greatly with both red meat consumption and dietary quality. In supplement table 1, the same characteristics are shown distributed on processed meat consumption. The



distribution of meat consumption and dietary quality in the population showed a median consumption of 65 g/day for red meat and of 35 g/day for processed meat (supplement table 2).

The hazard ratio for IHD for increments of 100 g/day of red meat was 1.23 (95% CI 0.99;1.53). For increments of 50 g/day of processed meat the hazard ratio for IHD was 1.09 (95% CI 0.93;1.29) (table 3).

The hazard ratio for IHD for high consumption compared with low consumption of red meat was 1.26 (95% CI 0.93;1.69). For processed meat, the hazard ratio for IHD for high consumption compared with low consumption was 1.10 (95% CI 0.81;1.50). The results on ACS were similar, though estimates were lower (supplement table 3).

Associations between meat consumption and risk of IHD stratified by dietary quality are presented in [table 3](#).

For the continuous measure of red meat consumption, the hazard ratio was strongest among those with a healthy diet (1.93 [95% CI 1.18;3.17]) followed by those with an unhealthy diet (1.30 [95% CI 0.89;1.89]), and a medium diet (0.98 [95% CI 0.71;1.36]). The test for interaction showed that these estimates were not statistically different ( $p=0.10$ ). The same was true for the categorical measure of red meat consumption ( $p=0.22$ ). Dietary quality did not modify the associations between processed meat consumption and risk of IHD either.

Results on associations between red and processed meat consumption and risk of ACS stratified by dietary quality showed the same trends indicating that dietary quality did not modify the associations (supplement table 4). Analyses on poultry consumption are shown in supplement tables 5-9. The results showed no significant associations between poultry consumption and risk of IHD or ACS.

Separate analyses on the three cohorts from DANSDA 2000-2002, 2003-2008, and 2011-2013, respectively were also performed. These showed some differences in effect estimates, but none of the results were statistically significant, and we do not interpret these as different effects dependent on survey year.

## Discussion

The results suggested a trend with higher consumption of red or processed meat being associated with higher risk of IHD. The effect estimate was highest for consumption of red meat. However, neither of these trends were statistically significant. Stratification by dietary quality did not indicate that associations between meat consumption and risk of IHD was modified by dietary quality.

The trend for red meat consumption is similar to those found in some previous studies (2-4, 14), though these studied cardiovascular mortality. Further, a meta-analysis did not find an association with IHD mortality (2) and another meta-analysis did not find an association with risk of coronary heart disease (1). The trend for processed meat is somewhat weaker than the associations previously reported in systematic reviews (1, 2) though one of them only found an association with the broader cardiovascular mortality and not with IHD mortality (2). A cohort study conducted in America (14) on heart mortality with detailed dietary information found associations closer in magnitude to the trends observed in our study.

Meat consumption and especially red and processed meat consumption has been heavily debated in relation to possible health effects. Associations between meat consumption and heart disease have mainly been attributed to saturated fat and cholesterol content, but in regards to processed meat also to ingredients such as sodium or other preservatives (15). No agreement has been reached on recommended daily intake. However, The World Cancer Research Fund recommends that consumption of red meat should be limited to no more than 350-500 g/week, and that consumption of processed meat should be as low as possible (16). This recommendation is however based on possible carcinogenic effects of meat.

Identified associations and differences between study findings are influenced by differences in study populations and exposure and outcome definitions. For example, associations may differ between regions or even countries. In a meta-analysis on cardiovascular mortality, an association was found for processed meat consumption, but this association could only be observed in studies conducted in US populations and not in studies conducted in European or Asian population (17). Also, both red meat and processed meat are broader categories of meat that include different types of meat. What characterises dietary red meat or processed

meat may therefore differ between countries. For example, pork may be a bigger contributor to red meat consumption in some countries compared to others, where beef could be a bigger contributor to red meat consumption. Previous analyses on the data included in this study showed that red meat consumption is equally represented by beef/veal and pork and that processed meat consumption is mainly represented by “other processed meat”, some salamis and sausages and very little bacon (18). As few cohort studies have the power to analyse meat consumption in this level of detail, it is often not known what characterises red meat or processed meat in the diet of the particular study population. Other differences between countries could be in cooking methods. Examples include differences in how much salt is used or how hard the meat is fried, but also in the meat’s fat contents, which is believed to be part of the biological pathway between meat consumption and risk of heart disease.

Other factors that will influence the estimates are the abilities to include influential confounders in the analyses. In the present study, both behavioural factors, socioeconomic factors, BMI and disease history were included as confounders. BMI is commonly included in nutritional epidemiology. Yet, BMI is also influenced by diet and could be viewed as a mediator instead of a confounder in the associations between meat consumption and the outcomes. However, information on meat consumption, dietary quality, and BMI is collected cross-sectionally, and it is not possible to deduce the temporality between these measures. No adjustment was undertaken for other meat types, but consumption of different types of meat are correlated. It is therefore possible that such adjustments would have been appropriate to disentangle the effects of the different types of meat from each other. For example, other published analyses on the same data showed that high consumption of red meat was associated with low consumption of poultry (18).

We found a statistically significant association between red meat consumption measured continuously and IHD risk among participants with a healthy diet. Yet, we consider this a chance finding, which should be interpreted with caution. However, if this association is in fact true, a possible explanation could be if some of those with a healthy diet have made healthy dietary choices because of being diagnosed with high cholesterol or high blood pressure, which are risk factors for IHD. The results did not support that dietary quality modified associations between meat consumption and risk of heart disease. This may be surprising as

it could be expected that an overall healthy diet could counterbalance some of the potential harmful health effects of meat consumption. In particular, previous studies have shown that dietary quality is inversely associated with outcomes such as myocardial infarction (19) and cardiovascular events (20). The results are however also influenced by the applied measure of dietary quality, which does not include all possible aspects of a healthy diet. A study conducted in Sweden found a statistically significant association between consumption of red meat and cardiovascular mortality. This association did not differ between those with high and low consumption of fruit and vegetables (6). Though only one aspect of a healthy diet was included in the Swedish analyses, the findings, in line with the findings from the present analyses, do not support the hypothesis that other dietary factors heavily influence the associations between meat consumption and disease development. It is however possible that more detailed information on dietary quality assessed in a larger dataset would suggest different conclusions.

Even though the results indicate trends in the association between meat consumption and risk of IHD, neither of the results were statistically significant. This highlights that studies on dietary factors' association with disease outcomes require large cohorts with long follow-up and a sufficient number of disease outcomes in the population. The study population was followed for 5-18 years, but a major limitation of the study is that the population could be as young as 15 years at baseline. Hence, only a limited number of disease outcomes were observed in the population. This has deteriorated the power of the study and therefore limited the ability to reach statistical significance for observed associations. This is especially true for analyses on interactions between meat consumption and dietary quality.

Other limitations include that even though the DANSDA surveys are representative regarding gender and age, in the latest surveys, participants with short education were underrepresented (21), which may limit the generalisability of the findings. In addition, the study only included a baseline dietary registration for each individual. Therefore, it is assumed that the dietary quality does not change during follow-up, but if the population has large variations in dietary quality during follow-up, this would influence the estimated associations. A study has previously shown that study participants' diet does change substantially over time (22). This limitation is however very common in nutritional epidemiology, as repeated information on

detailed dietary intake is rarely obtained. Another common limitation in nutritional epidemiology is misreporting of dietary intake, which could influence the associations if misreporting is biased by meat consumption, dietary quality, or other factors that influence the risk of IHD.

Our study also has several strengths. It was possible to ensure complete follow-up of the study cohort, since identification of IHD was based on linking national registers with high validity and completeness and we included complete information on migration and death. This linkage also enabled us to include only incident cases of IHD, thereby minimising the risk of reverse causality as we excluded those with disease before baseline. The study also included information on a range of dietary components, which made it possible to evaluate if associations were modified by the overall dietary quality. This included registration of the total diet for seven days, including weekend days, and the data collection process covering all seasons to allow for seasonal variations in dietary quality.

In conclusion, we found trends that higher consumption of red or processed meat was associated with higher risk of IHD, but neither of the trends were statistically significant. Associations between meat consumption and risk of IHD was not modified by dietary quality. Further studies that are able to explore modifying effects of dietary on associations between meat consumption and risk of IHD are needed.

**Funding:** The study was financially supported by “*Promilleafgiftsfonden for Landbrug*”, Denmark. *Promilleafgiftsfonden for Landbrug* had no influence on the study design, presentation and interpretation of results nor preparation of the manuscript.

**Conflict of interest:** The authors have nothing to declare.

**Author contributions:** All authors planned and designed the study. SPM was responsible for data management and analyses and drafted the manuscript. All authors contributed to interpretation of study results, critical revision of the paper and approval of final version.

1. Micha R, Michas G, Mozaffarian D. Unprocessed red and processed meats and risk of coronary artery disease and type 2 diabetes--an updated review of the evidence. *Current atherosclerosis reports*. 2012;14(6):515-24.
2. Abete I, Romaguera D, Vieira AR, Lopez de Munain A, Norat T. Association between total, processed, red and white meat consumption and all-cause, CVD and IHD mortality: a meta-analysis of cohort studies. *The British journal of nutrition*. 2014;112(5):762-75.
3. Bernstein AM, Sun Q, Hu FB, Stampfer MJ, Manson JE, Willett WC. Major dietary protein sources and risk of coronary heart disease in women. *Circulation*. 2010;122(9):876-83.
4. Pan A, Sun Q, Bernstein AM, Schulze MB, Manson JE, Stampfer MJ, et al. Red meat consumption and mortality: results from 2 prospective cohort studies. *Archives of internal medicine*. 2012;172(7):555-63.
5. Fogelholm M, Kanerva N, Mannisto S. Association between red and processed meat consumption and chronic diseases: the confounding role of other dietary factors. *European journal of clinical nutrition*. 2015;69(9):1060-5.
6. Bellavia A, Stilling F, Wolk A. High red meat intake and all-cause cardiovascular and cancer mortality: is the risk modified by fruit and vegetable intake? *The American journal of clinical nutrition*. 2016;104(4):1137-43.
7. Biltoft-Jensen A, Matthiessen J, Rasmussen LB, Fagt S, Groth MV, Hels O. Validation of the Danish 7-day pre-coded food diary among adults: energy intake v. energy expenditure and recording length. *The British journal of nutrition*. 2009;102(12):1838-46.
8. Tetens I, Andersen LB, Astrup A, Gondolf UH, Hermansen K, Jakobsen MU, et al. Evidensgrundlaget for danske råd om kost og fysisk aktivitet: Danmarks Tekniske Universitet, Fødevareinstituttet; 2013.
9. Knudsen V, Fagt S, Trolle E, Matthiessen J, Groth M, Biltoft-Jensen A, et al. Evaluation of dietary intake in Danish adults by means of an index based on food-based dietary guidelines. *Food & Nutrition Research*. 2012;56(1):17129.

10. Pedersen CB. The Danish Civil Registration System. *Scandinavian journal of public health*. 2011;39(7 Suppl):22-5.
11. Lynge E, Sandegaard JL, Rebolj M. The Danish National Patient Register. *Scandinavian journal of public health*. 2011;39(7 Suppl):30-3.
12. Helweg-Larsen K. The Danish Register of Causes of Death. *Scandinavian journal of public health*. 2011;39(7 Suppl):26-9.
13. Jensen VM, Rasmussen AW. Danish Education Registers. *Scandinavian journal of public health*. 2011;39(7 Suppl):91-4.
14. Etemadi A, Sinha R, Ward MH, Graubard BI, Inoue-Choi M, Dawsey SM, et al. Mortality from different causes associated with meat, heme iron, nitrates, and nitrites in the NIH-AARP Diet and Health Study: population based cohort study. *Bmj*. 2017;357:j1957.
15. Bronzato S, Durante A. A Contemporary Review of the Relationship between Red Meat Consumption and Cardiovascular Risk. *International journal of preventive medicine*. 2017;8:40.
16. World Cancer Research Fund/American Institute for Cancer Research. Continuous Project Expert Report 2018. Recommendations and public health and policy implications. 2018.
17. Wang X, Lin X, Ouyang YY, Liu J, Zhao G, Pan A, et al. Red and processed meat consumption and mortality: dose-response meta-analysis of prospective cohort studies. *Public health nutrition*. 2016;19(5):893-905.
18. Mejbørn H, Møller SP, Thygesen LC, Køup K, Biloft-Jensen A. Dietary patterns, meat intake and health. National Food Institute, Technical University of Denmark, Kgs. Lyngby; 2019.
19. Hansen CP, Overvad K, Tetens I, Tjønneland A, Parner ET, Jakobsen MU, et al. Adherence to the Danish food-based dietary guidelines and risk of myocardial infarction: a cohort study. *Public health nutrition*. 2018;21(7):1286-96.

20. Hlebowicz J, Drake I, Gullberg B, Sonestedt E, Wallstrom P, Persson M, et al. A high diet quality is associated with lower incidence of cardiovascular events in the Malmo diet and cancer cohort. *PloS one*. 2013;8(8):e71095.
21. Pedersen AN, Christensen T, Matthiessen J, Knudsen VK, Sørensen MR, Biloft-Jensen AP, et al. Danskernes kostvaner 2011-2013: DTU Fødevareinstituttet, Danmarks Tekniske Universitet; 2015.
22. Dekker LH, Boer JM, Stricker MD, Busschers WB, Snijder MB, Nicolaou M, et al. Dietary patterns within a population are more reproducible than those of individuals. *The Journal of nutrition*. 2013;143(11):1728-35.



Table 1. Baseline characteristics of study population stratified by consumption of red meat and dietary quality. n=8,007

	Consumption of red meat <sup>1</sup>			Dietary score <sup>2</sup>		
	Low consumption of red meat	Medium consumption of red meat	High consumption of red meat	Unhealthy diet	Medium diet	Healthy diet
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
<b>Age, mean (sd)</b>	47 (14)	48 (13)	48 (13)	43 (14)	48 (13)	51 (13)
<b>Sex</b>						
Men	570 (29.0)	1752 (43.2)	1421 (71.4)	1130 (59.7)	1918 (47.4)	695 (33.7)
Women	1394 (71.0)	2302 (56.8)	568 (28.6)	763 (40.3)	2131 (52.6)	1370 (66.3)
<b>Ethnicity</b>						
Danish	1902 (96.8)	3968 (97.9)	1948 (97.9)	1865 (98.5)	3947 (97.5)	2006 (97.1)
Western	30 (1.5)	37 (0.9)	15 (0.8)	12 (0.6)	45 (1.1)	25 (1.2)
Non-western	32 (1.6)	49 (1.2)	26 (1.3)	16 (0.8)	57 (1.4)	34 (1.6)
<b>Educational level</b>						
Long	694 (35.3)	1335 (32.9)	569 (28.6)	371 (19.6)	1352 (33.4)	875 (42.4)
Medium	815 (41.5)	1757 (43.3)	960 (48.3)	999 (52.8)	1734 (42.8)	799 (38.7)
Short	455 (23.2)	962 (23.7)	460 (23.1)	523 (27.6)	963 (23.8)	391 (18.9)
<b>BMI</b>						
<18.5	44 (2.2)	73 (1.8)	19 (1.0)	38 (2.0)	70 (1.7)	28 (1.4)
18.5-24.9	1187 (60.4)	2200 (54.3)	842 (42.3)	906 (47.9)	2189 (54.1)	1134 (54.9)
25-29.9	545 (27.7)	1311 (32.3)	824 (41.4)	678 (35.8)	1326 (32.7)	676 (32.7)
≥30	188 (9.6)	470 (11.6)	304 (15.3)	271 (14.3)	464 (11.5)	227 (11.0)
<b>Smoking</b>						
Never	927 (47.2)	1846 (45.5)	861 (43.3)	716 (37.8)	1823 (45.0)	1095 (53.0)
Former	494 (25.2)	1141 (28.1)	562 (28.3)	363 (19.2)	1181 (29.2)	653 (31.6)
Current	543 (27.6)	1067 (26.3)	566 (28.5)	814 (43.0)	1045 (25.8)	317 (15.4)
<b>Physical activity</b>						
None	161 (8.2)	351 (8.7)	186 (9.4)	281 (14.8)	306 (7.6)	111 (5.4)
Light	784 (39.9)	1648 (40.7)	784 (39.4)	809 (42.7)	1662 (41.0)	745 (36.1)
Moderate/hard	1019 (51.9)	2055 (50.7)	1019 (51.2)	803 (42.4)	2081 (51.4)	1209 (58.5)
<b>Meat consumption g/day, mean (sd)</b>						
Red meat	25 (11)	67 (16)	141 (46)	90 (58)	75 (47)	61 (40)
Processed meat	32 (27)	41 (31)	59 (41)	55 (40)	44 (33)	31 (25)
<b>Dietary score, mean (sd)</b>	3.3 (0.9)	3.1 (0.8)	2.8 (0.8)	1.9 (0.4)	3.1 (0.4)	4.1 (0.3)

<sup>1</sup>Low consumption of red meat: < 41 g/day; Medium consumption of red meat: 41-97 g/day; High consumption of red meat: > 97 g/day.

<sup>2</sup>Unhealthy diet: < 2.4 on the dietary quality score; Medium diet: 2.4-3.7 on the dietary score; Healthy diet: > 3.7 on the dietary quality score.

Abbreviations: n, number of participants; sd, standard deviation

Table 2. Association between red meat consumption, processed meat consumption and risk of ischemic heart disease, n=8,007

	<b>No. of cases</b>	<b>IR<sup>1</sup></b>	<b>HR [95%CI]<sup>2</sup></b>	<b>HR [95%CI]<sup>3</sup></b>
Red meat, low (< 41 g/day)	96	492	1.00	1.00
Red meat, medium (41-97 g/day)	226	562	1.04 [0.82;1.32]	1.07 [0.84;1.37]
Red meat, high (> 97 g/day)	117	669	1.17 [0.88;1.55]	1.26 [0.93;1.69]
Red meat, per 100 g/day			1.14 [0.93;1.40]	1.23 [0.99;1.53]
Processed meat, low (< 19 g/day)	114	610	1.00	1.00
Processed meat, medium (19-58 g/day)	205	515	0.83 [0.65;1.05]	0.89 [0.70;1.13]
Processed meat high (> 58 g/day)	120	641	1.02 [0.77;1.35]	1.10 [0.81;1.50]
Processed meat, per 50 g/day			1.05 [0.91;1.22]	1.09 [0.93;1.29]

<sup>1</sup>Per 100,000 person-years.

<sup>2</sup>Adjusted by sex.

<sup>3</sup>Adjusted by sex, educational attainment (the year before baseline), ethnicity, smoking, physical activity, alcohol, BMI, history of diabetes (10 years before baseline), and total energy intake.

Abbreviations: n, number of participants; IR, incidence rates; HR, hazard ratios; CI, confidence interval.

Table 3. Association between red meat consumption, processed meat consumption and risk of ischemic heart disease. Stratified by dietary quality<sup>3</sup>, n=8,007

	Unhealthy diet no. of cases	Medium diet no. of cases	Healthy diet no. of cases	Unhealthy diet IR <sup>1</sup>	Medium diet IR <sup>1</sup>	Healthy diet IR <sup>1</sup>	Unhealthy diet HR [95%CI] <sup>2</sup>	Medium diet HR [95%CI] <sup>2</sup>	Healthy diet HR [95%CI] <sup>2</sup>	p-value for interaction
Red meat, low (< 41 g/day)	9	57	30	277	604	441	1.00	1.00	1.00	
Red meat, medium (41-97 g/day)	48	122	56	522	558	610	2.00 [0.96;4.15]	0.80 [0.58;1.11]	1.34 [0.85;2.11]	0.22
Red meat, high (> 97 g/day)	37	63	17	583	716	726	2.13 [0.99;4.57]	0.97 [0.65;1.45]	1.66 [0.86;3.21]	
Red meat, per 100 g/day							1.30 [0.89;1.89]	0.98 [0.71;1.36]	1.93 [1.18;3.17]	0.10
Processed meat, low (< 19 g/day)	19	51	44	674	566	641	1.00	1.00	1.00	
Processed meat, medium (19-58 g/day)	35	128	42	389	597	448	0.64 [0.36;1.14]	1.04 [0.74;1.46]	0.75 [0.48;1.17]	0.52
Processed meat high (> 58 g/day)	40	63	17	573	653	814	0.83 [0.44;1.59]	1.19 [0.78;1.83]	1.36 [0.69;2.66]	
Processed meat, per 50 g/day							1.03 [0.77;1.38]	1.13 [0.90;1.41]	1.15 [0.73;1.80]	0.99

<sup>1</sup>Per 100,000 person-years<sup>2</sup>Adjusted by sex, educational attainment (the year before baseline), ethnicity, smoking, physical activity, alcohol, BMI, history of diabetes (10 years before baseline), and total energy intake.<sup>3</sup>Unhealthy diet: < 2.4 on the dietary quality score; Medium diet: 2.4-3.7 on the dietary score; Healthy diet: > 3.7 on the dietary quality score  
Abbreviations: n, number of participants; IR, incidence rates; HR, hazard ratios; CI, confidence interval