



Co-localization of carbapenem (*bla*OXA-162) and colistin (*mcr-1*) resistance genes on a transferable IncHI2 plasmid in *Escherichia coli* of chicken origin

Bortolaia, Valeria; Ronco, Troels; Romascu, Luminita; Nicorescu, Isabela; Milita, Nicoleta M.; Vaduva, Angela M.; Leekitcharoenphon, Pimlapas; Kjeldgaard, Jette S.; Hansen, Inge M.; Svendsen, Christina A.

Total number of authors:
15

Published in:
Journal of Antimicrobial Chemotherapy

Link to article, DOI:
[10.1093/jac/dkab285](https://doi.org/10.1093/jac/dkab285)

Publication date:
2021

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Bortolaia, V., Ronco, T., Romascu, L., Nicorescu, I., Milita, N. M., Vaduva, A. M., Leekitcharoenphon, P., Kjeldgaard, J. S., Hansen, I. M., Svendsen, C. A., Mordhorst, H., Guerra, B., Beloeil, P.-A., Hoffmann, M., & Hendriksen, R. S. (2021). Co-localization of carbapenem (*bla*_{OXA-162}) and colistin (*mcr-1*) resistance genes on a transferable IncHI2 plasmid in *Escherichia coli* of chicken origin. *Journal of Antimicrobial Chemotherapy*, 76(11), 3063-3065. [dkab285]. <https://doi.org/10.1093/jac/dkab285>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

J Antimicrob Chemother 2021; **76**: 3063–3065
doi:10.1093/jac/dkab285
Advance Access publication 15 August 2021

Co-localization of carbapenem (*bla*_{OXA-162}) and colistin (*mcr-1*) resistance genes on a transferable IncHI2 plasmid in *Escherichia coli* of chicken origin

Valeria Bortolaia^{1†}, Troels Ronco^{1*†},
Luminita Romascu^{2,3}, Isabela Nicorescu⁴,
 Nicoleta M. Milita², Angela M. Vaduva⁴,
Pimplapas Leekitcharoenphon¹, Jette S. Kjeldgaard¹,
Inge M. Hansen¹, Christina A. Svendsen¹,
Hanne Mordhorst¹, Beatriz Guerra⁵,
Pierre-Alexandre Beloeil⁵, Maria Hoffmann⁶ and
René S. Hendriksen¹

¹Technical University of Denmark, National Food Institute, European Union Reference Laboratory for Antimicrobial Resistance (EURL-AMR), WHO Collaborating Centre for Antimicrobial Resistance in Foodborne Pathogens and Genomics, FAO Reference Laboratory for Antimicrobial Resistance (FAO RL), Kgs Lyngby, Denmark; ²Institute for Diagnosis and Animal Health, University of Bucharest, Bucharest, Romania; ³Faculty of Veterinary Medicine, University of Agronomic Sciences and Veterinary Medicine of Bucharest, Bucharest, Romania; ⁴Institute for Hygiene and Veterinary Public Health, University of Bucharest, Bucharest, Romania; ⁵European Food Safety Authority, Parma, Italy; ⁶Center for Food Safety and Applied Nutrition, U.S. Food and Drug Administration, College Park, MD, USA

*Corresponding author. E-mail: troro@food.dtu.dk
†These authors contributed equally to this work.

Carbapenems are critically important antimicrobials for the treatment of clinical infections caused by MDR Enterobacteriaceae, with colistin reserved for those cases in which even carbapenems are ineffective. Therefore, co-localization of carbapenem and colistin resistance genes on the same plasmid is worrisome and, to date, has been described only in China in an *Escherichia coli* from a diseased chicken (*bla*_{NDM-4} and *mcr-1* on an IncHI2/ST3 plasmid)¹ and in an *E. coli* from a diseased pet cat (*bla*_{NDM-5} and *mcr-1* on an IncX3-X4 hybrid plasmid).² Here, we report a novel case of co-localization of *bla*_{OXA-162} carbapenemase and *mcr-1* on a transferable IncHI2 plasmid in *E. coli* from a healthy broiler chicken in Europe.

Within the framework of the specific monitoring of ESBL-, AmpC- or carbapenemase-producing indicator commensal *E. coli*

according to Commission Implementing Decision 2013/652/EU ‘on the monitoring and reporting of antimicrobial resistance in zoonotic and commensal bacteria’, four presumptive carbapenemase-producing *E. coli* from chicken caeca and meat were collected in Romania in 2016 (Table S1, available as [Supplementary data](#) at JAC Online). The four isolates were tested for antimicrobial susceptibility by broth microdilution using EUVSEC and EUVSEC2 Sensititre panels (Thermo Fisher Scientific, MA, USA) according to the manufacturer’s instructions and obtained MIC values were interpreted according to EUCAST epidemiological cut-off values (<http://eucast.org/>). Confirmed carbapenemase-producing isolates were sequenced (WGS) using Illumina MiSeq (CA, USA) as previously described³ and PacBio (Menlo Park, CA, USA) technologies. PacBio sequences were *de novo* assembled using the Hierarchical Genome Assembly Process version 4.0 (HGAPv4) with the ‘aggressive option’ and the final polished sequence was obtained using the Resequencing analysis (default settings). *De novo* assembly of Illumina reads and analyses of assembled sequences were performed using the CGE tools with default settings (<http://www.genomicepidemiology.org/>). Accession numbers are in Table S2. BLASTn analyses, alignments and mapping were carried out using CLC Genomics workbench v20.0.4 (Qiagen, Aarhus, Denmark) and plasmids were annotated using RAST.⁴ Previously described IncHI2 plasmids, including the IncHI2 prototype plasmid R478 and representative IncHI2 plasmids with and without *mcr-1* (Table S3), were used for comparison, which was visualized using BRIG.⁵ Plasmid transferability was assessed by filter-mating experiments. Briefly, strain CFSAN083827 was used as the donor and *E. coli* MT102 and *Salmonella* Typhimurium SL5324 were used as recipients in independent experiments. Transconjugants were selected on selective medium containing colistin (2 mg/L), cefotaxime (0.5 mg/L) and rifampicin (50 mg/L) and verified using PCR for *mcr-1*⁶ and *bla*_{OXA-48},⁷ followed by WGS (Illumina MiSeq) and bioinformatics analyses as described above.

The carbapenemase phenotype (meropenem MIC >0.12 mg/L) was confirmed for three isolates that also exhibited resistance to antimicrobials other than β-lactams and one of them also showed colistin resistance. See Table S1. Genomic characterization showed that the three isolates had different serotypes, STs, plasmid contents, antimicrobial resistance genes and *parC* mutations mediating fluoroquinolone resistance. The carbapenemase phenotype was mediated by *bla*_{OXA-162}, which was carried on IncHI2/pST4 plasmids. These IncHI2 plasmids, here designated pCFSAN083827_1, pCFSAN083828_1 and pCFSAN083829_1, had a GC content of 46.8%, 47.1% and 46.3%, respectively, and differed by size (Table S1). *bla*_{OXA-162} was carried on the 21.9 kb transposon Tn6237 (Figure 1a). In all three plasmids, Tn6237 was almost entirely conserved (>99.5% nucleotide identity and 100% sequence length) compared with pRA35.⁸ Sequence alignment showed that Tn6237 was identical across the three plasmids, with the exception of one SNP in *parB*. In pCFSAN083828_1 and pCFSAN083829_1 a C was found at position 341, which is similar to pRA35, whereas pCFSAN083827_1 had a T. This leads to a L (leucine) in pCFSAN083827_1 and P (proline) in pCFSAN083828_1 and pCFSAN083829_1 at amino acid position 114. The *parB* sequence of

© The Author(s) 2021. Published by Oxford University Press on behalf of the British Society for Antimicrobial Chemotherapy. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com




Figure 1. BLAST atlases of IncHI2 plasmids. (a) IncHI2 plasmids from this study. pCFSAN083828_1 (p828, due to layout limitations) and pCFSAN083829_1 (p829) were compared with pCFSAN083827_1 (p827; reference sequence). Mobile element (*meb*), insertion sequence (IS) and transposase (*tnp*) genes are shown in purple, whereas IncHI2 backbone genes are shown in grey. Antimicrobial resistance genes are shown in black and it can be seen that only pCFSAN083829_1 is missing *aph(3'')-Ib*, *aph(6)-Id*, *floR* and *sul2*. Notably, only pCFSAN083827_1 carries *mcr-1* located on Tn6330, whereas all three plasmids carry *bla_{OXA-162}* located on Tn6237. (b) Comparative plasmid analysis. The three IncHI2/ST4 plasmids from this study were compared with other IncHI2 plasmids of ST1, 3 and 4 (Table S1) using pCFSAN083827_1 as the reference sequence. Tn6237 is only found in the plasmids from this study, whereas replication (*rep*), transfer (*tra*) and tellurium resistance (*ter*) genes were nearly entirely conserved among all analysed plasmids. This figure appears in colour in the online version of JAC and in black and white in the print version of JAC.

pCFSAN083827_1 has no 100% identical sequence in GenBank, whereas that of pCFSAN083828_1 and pCFSAN083829_1 has.

Notably, plasmid pCFSAN083827_1 carried the colistin resistance gene *mcr-1* on a 4.7 kb transposon, i.e. Tn6330,⁹ flanked by IS*Apl1* (Figure 1a). In pCFSAN083827_1, Tn6330 was entirely conserved (100% nucleotide identity and 100% sequence length) and was located next to a region of conjugative transfer genes (Figure 1a). Comparison of the three plasmids from this study with selected IncHI2 plasmids showed that Tn6237 (carrying *bla*_{OXA-162}) was only present in the plasmids from this study, whereas Tn6330 was also found on other IncHI2 plasmids inserted at the same position (Figure 1b). The replication, transfer and tellurium resistance genes were nearly entirely conserved among all analysed IncHI2 plasmids (Figure 1b). The filter-mating experiment confirmed that pCFSAN083827_1 was transferable from isolate CFSAN083827 to the *E. coli* and *Salmonella* sp. recipients and transferred all of its antimicrobial resistance genes (Table S1).

Previously, *mcr-1*-carrying IncHI2 plasmids have been widely observed among isolates from chickens, other production animals and food products from Europe¹⁰ and, indeed, pCFSAN083827_1 shared similarities with those plasmids, including, among others, the Tn6330 insertion site (Figure 1b). However, the presence of Tn6237 with *bla*_{OXA-162} on an *mcr-1*-carrying IncHI2 plasmid is unique. This study highlights the importance of monitoring antimicrobial resistance in a One Health perspective by leveraging on WGS technologies that allow fine resolution of mobilizable elements mediating resistance.

Funding

We acknowledge funding from the European Commission to the European Union Reference Laboratory for Antimicrobial Resistance (EURL-AR).

Transparency declarations

None to declare.

Disclaimer

B. Guerra and P.-A. Beloeil are currently employed by the European Food Safety Authority (EFSA) in its BIOCONTAM Unit that provides scientific and administrative support to EFSA's scientific activities in the area of Microbial Risk Assessment. The positions and opinions presented in this article are those of the authors alone and are not intended to represent the views or scientific works of EFSA.

Supplementary data

Tables S1, S2 and S3 are available as [Supplementary data](#) at JAC Online.

References

- Liu BT, Song FJ, Zou M *et al.* High incidence of *Escherichia coli* strains cohabiting *mcr-1* and *bla*_{NDM} from chickens. *Antimicrob Agents Chemother* 2017; **61**: e02347–16.
- Sun J, Yang RS, Zhang Q *et al.* Co-transfer of *bla*_{NDM-5} and *mcr-1* by an IncX3-X4 hybrid plasmid in *Escherichia coli*. *Nat Microbiol* 2016; **1**: 16176.
- Hendriksen RS, Leekitcharoenphon P, Lukjancenko O *et al.* Genomic signature of multidrug-resistant *Salmonella enterica* serovar Typhi isolates related to a massive outbreak in Zambia between 2010 and 2012. *J Clin Microbiol* 2015; **53**: 262–72.
- Brettin T, Davis JJ, Disz T *et al.* RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 2015; **5**: 8365.
- Alikhan NF, Petty NK, Ben Zakour NL *et al.* BLAST Ring Image Generator (BRIG): simple prokaryote genome comparisons. *BMC Genomics* 2011; **12**: 402.
- Rebelo AR, Bortolaia V, Kjeldgaard JS *et al.* Multiplex PCR for detection of plasmid-mediated colistin resistance determinants, *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4* and *mcr-5* for surveillance purposes. *Euro Surveill* 2018; **23**: pii=17-00672.
- Poirel L, Walsh TR, Cuvillier V *et al.* Multiplex PCR for detection of acquired carbapenemase genes. *Diagn Microbiol Infect Dis* 2011; **70**: 119–23.
- Beyrouthy R, Robin F, Delmas J *et al.* IS1R-mediated plasticity of IncL/M plasmids leads to the insertion of *bla*_{OXA-48} into the *Escherichia coli* chromosome. *Antimicrob Agents Chemother* 2014; **58**: 3785–90.
- Li R, Xie M, Zhang J *et al.* Genetic characterization of *mcr-1*-bearing plasmids to depict molecular mechanisms underlying dissemination of the colistin resistance determinant. *J Antimicrob Chemother* 2017; **72**: 393–401.
- Matamoros S, Van Hattem JM, Arcilla MS *et al.* Global phylogenetic analysis of *Escherichia coli* and plasmids carrying the *mcr-1* gene indicates bacterial diversity but plasmid restriction. *Sci Rep* 2017; **7**: 15364.