



Patients' assessment of care for type 2 diabetes: Results of the Patient Assessment of Chronic Illness Care scale in a Danish population

Frølich, Anne; Nielsen, Ann; Glümer, Charlotte; Eriksen, Christian U.; Maindal, Helle Terkildsen; Kleist, Bodil Helbech; Birke, Hanne; Stockmarr, Anders

Published in:
BMC Health Services Research

Link to article, DOI:
[10.1186/s12913-021-07051-6](https://doi.org/10.1186/s12913-021-07051-6)

Publication date:
2021

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Frølich, A., Nielsen, A., Glümer, C., Eriksen, C. U., Maindal, H. T., Kleist, B. H., Birke, H., & Stockmarr, A. (2021). Patients' assessment of care for type 2 diabetes: Results of the Patient Assessment of Chronic Illness Care scale in a Danish population. *BMC Health Services Research*, 21(1), Article 1069. <https://doi.org/10.1186/s12913-021-07051-6>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

RESEARCH ARTICLE

Open Access



Patients' assessment of care for type 2 diabetes: Results of the Patient Assessment of Chronic Illness Care scale in a Danish population

Anne Frølich^{1,2*} , Ann Nielsen³, Charlotte Glümer⁴, Christian U Eriksen⁵ , Helle Terkildsen Maimdal⁶ ,
Bodil Helbech Kleist⁷, Hanne Birke⁸  and Anders Stockmarr⁹ 

Abstract

Background: The Patient Assessment of Chronic Illness Care (PACIC) scale is the most appropriate for assessing self-reported experience in chronic care. We aimed to validate the PACIC questionnaire by (1) assess patients' perception of the quality of care for Danish patients with type 2 diabetes, (2) identify which factors are most important to the quality of care designated by the five subscales in PACIC, and (3) the validity of the questionnaire.

Methods: A survey of 7,745 individuals randomly selected from the National Diabetes Registry. Descriptive statistics inter-item and item-rest correlations and factor analysis assessed the PACIC properties. Quality of care was analysed with descriptive statistics; linear and multiple regression assessed the effect of forty-nine covariates on total and subscale scores.

Results: In total, 2,696 individuals with type 2 diabetes completed $\geq 50\%$ of items. The floor effect for individual items was 8.5–74.5 %; the ceiling effect was 4.1–47.8 %. Cronbach's alpha was 0.73–0.86 for the five subscales. The comparative fit index (CFI) and the Tucker–Lewis index (TLI) were 0,87, and 0,84, respectively. Mean PACIC score was 2.44 (± 0.04). Respondents, who receive diabetes care primarily at general practice and outpatient clinics had higher scores compared to those receiving care at a private specialist. Receiving rehabilitation was followed by higher scores in all subscales. Those 70 years or older had lower mean total and subscale scores compared to younger patient groups.

A higher number of diabetes visits were associated with higher total scores; a higher number of emergency department visits were associated with lower total scores. The effects of healthcare utilisation on subscale scores varied.

* Correspondence: anfrosund@sun.ku.dk

¹Section of General Practice, Department of Public Health, University of Copenhagen, 1356 K, Copenhagen, Denmark

²Innovation and Research Centre for Multimorbidity, Slagelse Hospital, 4200 Slagelse, Denmark

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Conclusions: These results provide insight into variations in the quality of provided care and can be used for targeting initiatives towards improving diabetes care. Factors important to the quality of perceived care are having a GP or hospital outpatient clinic as the primary organization. Also having a higher number of visits to the two organizations are perceived as higher quality of care as well as participating in a rehabilitation program. Floor and ceiling effects were comparable to an evaluation of the PACIC questionnaire in a Danish population. Yet, floor effects suggest a need for further evaluation and possible improvement of the PACIC questionnaire in a Danish setting. Total PACIC scores were lower than in other healthcare systems, possible being a result of different contexts and cultures, and of a need for improving diabetes care in Denmark.

Keywords: PACIC, Validation, Diabetes Mellitus, Type 2, Patient-Centred Care

Background

Disease management programs for type 2 diabetes patients have been implemented in the Capital Region of Denmark to support evidence-based care. The programs are founded based on the Chronic Care Model (CCM) that is one approach to improving chronic illness care [1]. Programs cover diagnosis, treatment, rehabilitation, and follow-up. The effectiveness of diabetes care increases by improving patient activation, information and self-management associated with behaviours, such as treatment adherence, a healthier diet and physical activity [2–4].

The twenty-item Patient Assessment of Chronic Illness Care (PACIC) questionnaire considers whether patients receive patient-centred care in alignment with the CCM. The PACIC is grouped into five subscales representing five care components of the CCM that is central to quality of care [5, 6]. The five care components are: patient activation, delivery system design/decision support, goal setting/tailoring care, problem-solving/contextual counselling, and follow-up/coordination [1, 7]. The PACIC has been translated into Danish and applied in a primary care setting in Denmark to assess data quality, internal consistency, and factorial structure [8]. The psychometric properties in that study were satisfactory, apart from floor and ceiling effects. Also, the subscales showed good model fit and could be used for separate sum scores. Later, the PACIC was applied in another Danish diabetes population to assess perceptions of care among asymptomatic diabetes individuals identified by a screening program and receiving intensive multifactorial treatment. Although the individuals were early in their disease trajectory, and thus the effect of treatment could exceed the disease burden, the intensive diabetes care did not worsen the perceptions of care compared to individuals receiving routine diabetes care [7]. Yet the overall low PACIC score indicated room for improvement in diabetes care [7]. Conversely, a German study comparing patients receiving usual care to those enrolled in a diabetes disease management program found that patients receiving structured evidence-based care scored significantly higher on all PACIC subscales [9]. A Dutch study

reviewing 37 instruments to measure self-reported experience in chronic care concluded PACIC to be the most appropriate [10]. Our objective of the study was to (1) assess the quality of care for Danish patients with type 2 diabetes, and (2) identify which factors that are most important to perceived quality of care designated by the five subscales in PACIC, and (3) the validity of the questionnaire.

Methods

Population

In 2013, we randomly selected 8,000 individuals aged 40 years or older with diabetes living in the Capital Region from the National Diabetes Registry [11]. Of these, 255 had died or moved out of the region, and the PACIC was subsequently mailed to 7,745 individuals, along with a link to an identical electronic version. Three weeks later, we sent a reminder to non-respondents. In total, 4,190 patients completed the PACIC, yielding a response rate of 54.1%. The inclusion criteria were persons 40 years or older with type 2 diabetes, living in the Capital region. The exclusion criteria were persons that had not diabetes and persons with type 1 diabetes. Of the responders, 2,810 had type 2 diabetes, while 1,380 patients (20%) were excluded.

PACIC scoring and additional variables

Each item was scored on a five-point scale ranging from one (no/never) to five (yes/always). Each subscale was scored by averaging scores for completed items, and the total score was calculated by averaging scores across all twenty items. In the Danish version, the wording of a few items was updated in collaboration with Judith Schaefer, developer of the PACIC [8]. The Danish version also includes items about diabetes type, the primary location of diabetes care and rehabilitation, and type(s) of received rehabilitation: diabetes education, smoking cessation, physical training, and dietician counselling. The Danish version can be provided by request to the authors.

PACIC scores were linked to national registers and health registers to obtain information on sex, age,

education, occupation, cohabitation status, socioeconomic area, multimorbidity status, number of visits with a general practitioner (GP), annual diabetes control visits, planned and acute hospital admissions and bed days, and readmissions. The socioeconomic area was identified by proximity to one of four hospital planning areas that differ socioeconomically. Chronic conditions included allergy, depression, stroke, dementia, chronic heart disease, hypertension, chronic pulmonary disease, high cholesterol, cancer, arthritis, osteoporosis, chronic back pain, and schizophrenia. Comorbidity status was based on algorithms as described in Robinson [12]. Multimorbidity was defined as the presence of two or more chronic conditions.

Statistical analyses

Of all 4,190 respondents, 1,380 (32.9%) did not have diabetes or had type 1 diabetes and were excluded. Of the remaining 2,810 respondents, 114 respondents completed fewer than ten items and were excluded. Thus, 2,696 respondents, who completed ten items or more, were included in the analysis. We used the expectation-maximisation (EM) algorithm to impute item values for 259 respondents who answered 50% or more but less than 100% of the items [13]. Comparative analyses without imputation were carried out.

Data quality was evaluated with standard summary statistics. Floor and ceiling effects were presented and analysed. Internal consistency was evaluated with correlational analyses within each subscale and between the five subscales using Cronbach's alpha.

Additionally, confirmatory factor analysis was carried out, investigating the fit of a latent class for each of the 5 PACIC subscores to data. This factor structure was assessed by the model χ^2 test, the Root Mean Square Error of Approximation (RMSEA), the Standardized Root Mean Square Residual (SRMR), the Tucker-Lewis Index (TLI) and the Comparative Fit Index (CFI). We considered a value of Cronbach's alpha of 0.7 or above as satisfactory [14]. Descriptive analyses included means, standard deviations, floor and ceiling effects for the total score, for each of the five subscales, and means and standard deviations for the twenty individual items.

To identify factors affecting total PACIC scores, we used linear regression analysis; the nature of the PACIC score as an average of twenty scores across five subscales invites continuous modelling. Multiple regression analysis was used to examine the total PACIC score and each subscale score for relationships with the variables listed above, a total of forty-nine different factors. The factors education, occupation, and socioeconomic area were multi-level factors, while the remaining factors

were binary or numeric. We used the following formula for modelling:

$$Y_i = \alpha_{education(i)} + \alpha_{occupation(i)} + \alpha_{area(i)} + \sum_{j=1}^{46} \beta_j X_{ij} + \epsilon_i, i = 1, \dots, n. \quad (1)$$

In formula (1), Y_i denotes either the total PACIC score or one of the five subscale scores, while X_{ij} denotes the value of the j^{th} of the forty-six binary or numeric factors for the i^{th} patient. The model was applied separately for the total PACIC score and each of the five subscale scores. The model reduction was done with the Likelihood Ratio method with a significance level of 5%. Subsequently, a forward selection procedure was applied to detect statistical significances obscured by the top-down model reduction. The forward selection procedure did not result in any alteration of the top-down reduced models. Analyses were performed with the software R, version 3.1.0 [15].

Results

Respondent characteristics

Of the 2,810 respondents, 59.1% were men, 78.4% were older than 60 years, 65.4% were retired, 80.7% of respondents had no education or education of short duration. Most (70.5%) received diabetes care from their GP, one fifth (21.8%) received care at an outpatient clinic, and a minority of 0.4% received care from a private specialist (Table 1).

Quality of PACIC data

In total, 2,434 respondents (86.7%) answered all twenty questions. Only 74 respondents (2.6%) did not answer any of the questions. Response rates were similar for all items, and the distributions of answered and unanswered questions were similar for the five subscales. The rate of answered questions decreased with increasing age, and women had a lower rate of answered questions than men.

The floor effect was 8.5–74.5%, with nineteen items higher than 15%. The ceiling effect was 4.1–47.8%, with eleven items of more than 15%. Cronbach's alpha varied between 0.73 and 0.86 for the 5 subscales. The mean inter-item correlation in the five subscales varied between 0.45 and 0.60. The inter-item correlation range was 0.34–0.73, and the item-rest correlation range was 0.47–0.79. Confirmatory factor analysis showed factor loadings for the twenty items ranging from 0.51 to 0.72. (Table 2)

The factor structure assessment characteristics are listed in Table 3.

Table 1 Characteristics of 2,810 respondents with type 2 diabetes

Sex	N (%)
Men	1,662 (59.1)
Women	1,148 (40.9)
Age in years	
40–49	135 (4.8)
50–59	475 (16.9)
60–69	1,064 (37.9)
70–79	811 (28.0)
≥ 80	325 (11.6)
Education level (total years of education)	
None (≤ 10)	890 (31.7)
Short (11–14)	1,378 (49.0)
Medium (15–16)	299 (10.6)
Long (≥ 17)	160 (5.7)
Not reported	83 (3.0)
Occupation	
Working or in school	827 (29.4)
Unemployed	30 (1.1)
Long-term disability	72 (2.6)
Early retirement	237 (8.4)
Retirement	1,603 (57.0)
Not reported	41 (1.5)
Cohabitation status	
Living alone	1,099 (39.1)
Living with spouse	1,711 (60.9)
Primary location of diabetes care	
General practitioner	1,980 (70.5)
Outpatient clinic	612 (21.8)
Private specialist	12 (0.4)
Not reported	49 (1.7)

PACIC score

The overall mean PACIC score was 2.44 (± 0.04). Subscale mean scores varied with the highest value of 3.26 (± 0.04) for delivery system design/decision support and the lowest value of 1.99 (± 0.04) for follow-up/coordination (Table 4).

The effect parameters for variables with a statistically significant effect on total and subscale scores are shown in Tables 5 and 6.

Age and sex

Total PACIC scores and subscale scores were inversely correlated with age and declined with increasing age from the age of approximately 70 years (Fig. 1). The effect was nonlinear for the total score and all subscales. A similar pattern was found for the five subscales.

Sex did not affect the total PACIC score. Men tended to have a lower score than women on the patient activation subscale, although the p-value of 0.02 did not indicate strong statistical significance.

Sociodemographic variables

Educational level, occupation, and socioeconomic area did not affect the total PACIC score (Table 6). Educational level affected subscale scores to a significantly negative degree for goal setting and follow-up/coordination ($p = 0.009$ for both). For both subscales, patients with the most education scored significantly lower than those with less education (Table 5). Occupation did not affect any subscales. Socioeconomic area affected patient activation ($p = 0.04$) and delivery system design/decision support ($p = 0.008$). For the latter, respondents living in planning area three had markedly lower scores than respondents living in the other areas; compared to a composite of all other areas, the estimated mean subscale score for respondents in planning area three was 0.16 lower ($p = 0.005$).

Diabetes treatment and rehabilitation

Respondents whose primary setting for diabetes care was a hospital outpatient clinic scored significantly and consistently higher for the total PACIC score on all subscales than respondents who did not attend a primary setting for care (Tables 4 and 5). Only twelve respondents reported a private specialist as the primary place for their diabetes care; their total PACIC scores were significantly higher than respondents who did not report a primary setting for diabetes care, but the effect was inconsistent across the subscales. Patients who reported receiving diabetes care at the hospital clinic had total PACIC scores that were an estimated 0.29 higher than patients who received diabetes care at their GP or a private specialist. However, the difference is borderline significant ($p = 0.05$). Patients who received diabetes care primarily at their GP or a specialist had an estimated 0.64 higher total PACIC scores than respondents who replied no primary setting for diabetes care or indicated “other” ($p < 0.0001$).

Participation in rehabilitation programs was associated with increased total and subscale scores except for smoking cessation (Tables 4 and 5). However, the effect varied according to the program location. Diabetes education with a GP showed the highest effect on the total PACIC score of all rehabilitation types, with an estimated rise in the total PACIC score of 0.62 (± 0.10). The effect on the total PACIC score from physical training was highest in the outpatient clinic, while dietary counselling yielded the highest increase in total PACIC score when it occurred in the municipality (Table 4).

Table 2 Results of the confirmatory factor analysis showing the standardized factor loadings and residuals for each item when modelled with its own scale

Over the past 12 months, when I received care for my chronic conditions, I was:	Standardized factor loading	Residual variance
Patient activation		
Item 1	0.629	0.027
Item 2	0.634	0.027
Item 3	0.626	0.027
Delivery system design/decision support		
Item 4	0.520	0.027
Item 5	0.560	0.027
Item 6	0.613	0.027
Goal setting/tailoring		
Item 7	0.666	0.025
Item 8	0.692	0.025
Item 9	0.510	0.025
Item 10	0.517	0.025
Item 11	0.623	0.025
Problem solving/contextual		
Item 12	0.613	0.025
Item 13	0.697	0.025
Item 14	0.724	0.025
Item 15	0.690	0.025
Follow-up/coordination		
Item 16	0.523	0.025
Item 17	0.564	0.025
Item 18	0.571	0.025
Item 19	0.586	0.025
Item 20	0.555	0.025

Contact with the healthcare system

The total PACIC score increased with the number of annual GP visits ($p = 0.02$) and decreased with the number of emergency department visits ($p = 0.005$, Tables 3 and 4). No consistent pattern occurred for the subscales. The patient activation subscale score decreased with the number of readmissions ($p = 0.009$) and for patients with a stroke diagnosis ($p = 0.04$) but increased with the number of annual GP visits ($p = 0.01$). Delivery system design/decision support subscale scores were lower for patients with a stroke diagnosis ($p = 0.005$) or a chronic heart condition ($p = 0.02$) and decreased with the number of outpatient contacts ($p = 0.04$) and planned hospitalisations ($p = 0.04$). Goal setting subscale scores decreased with the number of acute hospitalisations ($p = 0.007$) and increased with multimorbidity status ($p = 0.04$). Problem-solving/contextual counselling subscale scores decreased with the number of emergency department visits ($p = 0.01$) and increased with multimorbidity status ($p = 0.04$). Follow-up/coordination subscale scores increased with the number of GP visits ($p = 0.009$) and decreased as emergency department visits increased.

Discussion

The study showed that PACIC is reliable for assessing the quality of care that type 2 diabetes patients receive in Denmark. Central factors important to the quality of perceived care are having a hospital outpatient clinic as the primary organization for diabetes care, participating in a rehabilitation program, and having higher number of visits to the GP and outpatient visits. Compared to other healthcare systems, the low total PACIC score on 2.44 suggests a potentially lower quality of diabetes care in Denmark from the patients' perspective. The validation of the PACIC scale in a Danish context showed an acceptable ceiling effect and inter-item correlation, but floor effects persist for several items.

The study showed that PACIC is reliable for assessing the quality of provided care for type 2 diabetes in a Danish setting. In relation to the five subscales, we identified several factors important to the quality of care (Table 6). Patients perceived that they received better care if they were younger, were women, received annual control visits with their GP, and had primary location of diabetes in an outpatient clinic. Patients experiencing planned hospitalisations, emergency department visits

Table 3 The factor structure assessment characteristics

Statistic	χ^2	df	p	CFI	TLI	RMSEA	SRMR
Value for model fit with 5 subscales	3908	160	< 0.001	0.866	0.841	0.098	0.062

Table 4 Total PACIC score, five subscales, and item scores among respondents with type 2 diabetes ($N = 2,696^{**}$)

	n	Mean (SD)	Median	Missing (%)	Floor (%)	Ceiling (%)
Total score	2,696*	2.44 (0.02) *				
Patient activation scale	2,696*	2.66 (0.03) *				
Item 1	2,680	3.21 (0.03)	4	4.6	27.6	33.1
Item 2	2,647	2.19 (0.03)	1	5.8	52.4	13.1
Item 3	2,663	2.69 (0.03)	2	5.2	40.7	23.3
Delivery system design/decision support scale	2,696*	3.26 (0.02) *				
Item 4	2,670	2.45 (0.03)	2	5.0	45.0	16.4
Item 5	2,695	4.03 (0.03)	4	4.1	8.5	47.8
Item 6	2,678	3.43 (0.03)	4	4.7	19.7	35.0
Goal setting scale	2,696*	2.29 (0.02) *				
Item 7	2,673	2.87 (0.03)	3	4.9	32.2	22.4
Item 8	2,672	2.63 (0.03)	3	4.9	37.6	16.1
Item 9	2,661	1.83 (0.03)	1	5.3	69.9	12.0
Item 10	2,671	1.61 (0.02)	1	4.9	74.4	5.8
Item 11	2,695	2.63 (0.03)	3	4.1	36.2	15.6
Problem solving/contextual counselling scale	2,696*	2.40 (0.02) *				
Item 12	2,667	3.16 (0.03)	4	5.1	27.4	28.1
Item 13	2,671	2.05 (0.03)	1	4.9	59.0	11.0
Item 14	2,663	2.17 (0.03)	1	5.2	54.0	12.6
Item 15	2,677	2.33 (0.03)	2	4.7	48.1	14.3
Follow-up/ coordination scale	2,696*	1.99 (0.02) *				
Item 16	2,682	1.62 (0.02)	1	4.6	74.5	6.9
Item 17	2,675	1.57 (0.02)	1	4.8	74.0	4.1
Item 18	2,673	1.78 (0.03)	1	4.9	66.5	6.6
Item 19	2,677	2.82 (0.03)	3	4.7	37.8	25.4
Item 20	2,671	2.28 (0.03)	1	4.9	53.9	16.5

* Based on imputed data

** Out of 2,810 respondents, 114 respondents were excluded due to missing completed items, resulting in 2,696 respondents, who completed more than ten items

and readmissions perceived that the quality of care was lower. The results provided insight into variations in patients' perception of provided care and can be used for targeting initiatives towards improved care.

Study population

The sex distribution in our study population is consistent with earlier reports that men represent a higher proportion of the population with type 2 diabetes, with the standardised proportion ratio ranging from 1.1 to 1.4 in 1999–2011 [16]. The comparable ratio in our study population was 1.4, which differed significantly from 1.0 ($p < 0.0001$). The study populations mean age of 67.2 (40–97) years is slightly older than the mean age of 64 years in the Danish National Diabetes Register [17]. The proportion of individuals with only ten years of primary school (31.7%) [18] is slightly different, yet comparable, to another Danish diabetes study population, in which

28% had no more than 11 years of education [19]. The comparable levels of education provides acceptable representativeness with respect to education.

Quality of diabetes care and factors important to the quality of care

The lower total PACIC scores in all five subscales from patients receiving care primarily from a private specialist, compared to those receiving it at their GP or at the hospital outpatient clinic, may result from patients perceiving outpatient clinics to provide a higher level of specialist care. Also, patients with severe complications primarily receive hospital care and thus may receive more attention to their health status, which may result in a higher total PACIC score.

Participation in a rehabilitation program was, in general, associated with higher both total and subscale PACIC scores. The organisations receiving the highest

Table 5 Significant effect parameter estimates for total PACIC score in linear regression models

	Estimate (SD)	P value
Intercept	0.20 (0.56)	0.71
Age	0.042 (0.017)	0.01
Age squared	-3.8e-4 (1.3e-4)	0.002
Annual control visits	0.058 (0.024)	0.02
Emergency department visits	-0.083 (0.030)	0.005
Primary diabetes care location		
General practitioner	0.64 (0.09)	< 0.001
Outpatient clinic	0.93 (0.10)	< 0.001
Private specialist	0.65 (0.27)	0.02
Rehabilitation program location		
Patient education		
Municipality	0.15 (0.07)	0.04
General practitioner	0.62 (0.05)	< 0.001
Outpatient clinic	0.34 (0.05)	< 0.001
Physical training		
Municipality	0.18 (0.06)	0.004
General practitioner	0.24 (0.08)	0.004
Outpatient clinic	0.40 (0.08)	< 0.001
Dietary advice		
Municipality	0.50 (0.07)	< 0.001
General practitioner	0.36 (0.05)	< 0.001
Outpatient clinic	0.31 (0.05)	< 0.001

mean total PACIC scores varied with the type of diabetes rehabilitation provided. GP-generated patient education scored highest, 0.47 higher than patient education in the municipality. Patients receiving physical training in the hospital scored 0.22 higher than patients training in the municipality. Dietician advice in the municipality scored 0.14 higher than in the GP setting. The varying total scores for sites and types of rehabilitation may point to opportunities for inter-organisational learning and improvement.

The PACIC score decreased significantly with an increasing number of emergency department visits, which has not been previously reported. Experiencing an acute event in and of itself may cause a negative perception of the healthcare system. An increasing number of readmissions is followed by a decreasing mean score for the patient activation subscale and might be related to respondents interpreting readmissions as low-quality care.

Longer education negatively impacted the mean score for the goal setting and subscales, possibly due to well-educated patients not perceiving goal setting as relevant. The mean score for the delivery system design/decision support subscale was significantly lower for respondents living in planning area three, but with no differences

between mean subscale scores for patients living in the other planning areas after covariate adjustments. This may be explained by the characteristics of planning area three, which includes some of the most affluent communities. The healthcare utilisation pattern in area three differs from the other areas in the region; there is greater use of specialists in private practice and less use of outpatient visits and bed days. The impact of age on the total PACIC score was nonlinear, with no significant effect up to 60–70 years, where a steep decline occurred, and the estimated score dropped to approximately 2.2 at the age of 85 years (Fig. 1). To our knowledge, this effect has not previously been reported [7, 8]. Potential explanations include the possibility of treatment being perceived as less effective and failing to meet expectations due to age-related increases in disease severity and comorbidity. Also, as patients develop additional age-related conditions, they encounter more healthcare professionals. All subscales except follow-up/coordination were significantly related to age.

Assessment of PACIC questionnaire – imputation ceiling and floor effect and CFI and TLI estimates

With an increase in the study population of 10.6 % after imputation, the relevance of imputation is debatable. However, analyses comparing PACIC scores as a whole and stratified by sociodemographic variables revealed a consistent tendency towards lower PACIC scores for respondents with imputed values, compared to respondents completing all items. This introduces an upward bias in the population consisting of respondents answering all twenty items. A potential explanation is that the number of items answered decreased with age, which was also associated with lower total PACIC scores beginning around age 70 (Fig. 1). However, other characteristics of partial non-respondents may affect the PACIC scores. Thus, using imputed values was considered an appropriate improvement in the representativeness of the dataset.

The levels of floor and ceiling effects call for further investigation on specific items, particularly for those with high floor effects. The inter-item correlation is acceptable according to a lower limit of 0.15 [20], just as the item-rest correlation for fifteen of twenty items was higher than the maximum correlations with other scales, indicating item convergence within subscales. The values of Cronbach's alpha met acceptability criteria according to McDowell [21] and Sitzia [22]. Factor loadings from the confirmatory factor analysis were high. The model χ^2 test was strongly significant (Table 5). While undesirable, it is commonly seen in studies with large sample sizes (as ours and lower), even if the model is appropriate [23]. CFI and TLI estimates were slightly below the lower limit of 0.90 for acceptable fits [24], while RMSEA


Table 6 Effect parameters for PACIC subscale scores in linear regression models (N = 2,696*)

	Patient activation		Delivery system design/ decision support		Goal setting		Problem solving/ contextual counselling		Follow-up/ coordination	
	Estimate	P value	Estimate	P value	Estimate	P value	Estimate	P value	Estimate	P value
Intercept	-0.13	0.88	-0.17	0.81	1.52	< 0.001	1.37	< 0.001	-0.17	0.77
Age	0.065	0.007	0.068	< 0.001					0.048	0.006
Age squared	-5.7e-4	0.002	-5.5e-4	< 0.001	-8.2e-5	< 0.001	7.5e-5	< 0.001	-4.1e-4	0.002
Men	-0.12	0.02								
Living alone	-0.12	0.02								
Planning area		0.04		0.008						
2	-0.069	0.32	-0.11	0.07						
3	-0.19	0.02	-0.24	< 0.001						
4	-0.12	0.09	-0.13	0.02						
5	-0.33	0.02	-0.13	0.28						
Education										
Short					-0.066	0.009			-0.10	0.009
Medium					-0.079	0.35			-0.077	0.33
Long					-0.24	< 0.001			-0.21	< 0.001
Comorbidities										
Stroke	-0.23	0.04	-0.26	0.005						
Heart disease			-0.13	0.02						
Multimorbidity					0.22	0.04	0.29	0.03		
Utilization										
Annual control visits	0.09	0.01								
Planned hospitalization										
Emergency department visits			-0.013	0.04			-0.10	0.01	-0.09	0.002
GP visits									0.0045	0.009
Unplanned hospitalizations					0.062	0.007				
Out patient visits										
Readmissions	-0.14	.009	-0.0048	0.04						
Primary location of diabetes care										
General practice	0.66	< 0.001	1.13	< 0.001	0.48	< 0.001	0.58	< 0.001	0.45	< 0.001
Out patient Clinic	1.06	< 0.001	1.30	< 0.001	0.83	< 0.001	0.86	< 0.001	0.67	< 0.001
Private specialist			0.77	0.02	0.72	< 0.001	0.63	0.05		

Table 6 Effect parameters for PACIC subscale scores in linear regression models (N = 2,696*) (Continued)

	Patient activation		Delivery system design/ decision support		Goal setting		Problem solving/ contextual counselling		Follow-up/ coordination	
	Estimate	P value	Estimate	P value	Estimate	P value	Estimate	P value	Estimate	P value
Location of rehabilitation program										
Education										
Municipality			0.23	0.005					0.19	0.01
General practice	0.61	< 0.001	0.59	< 0.001	0.62	< 0.001	0.80	< 0.001	0.50	< 0.001
Out patient clinic	0.43	< 0.001	0.34	< 0.001	0.37	< 0.001	0.35	< 0.001	0.28	< 0.001
Physical training										
Municipality	0.32	< 0.001	0.17	0.02	0.17	0.01	0.21	0.008	0.15	0.02
General practice	0.34	0.004	0.24	0.02	0.31	< 0.001	0.24	0.03		
Out patient clinic	0.42	< 0.001	0.43	< 0.001	0.43	< 0.001	0.50	< 0.001	0.29	< 0.001
Dietary advice										
Municipality	0.42	< 0.001	0.56	< 0.001	0.57	< 0.001	0.51	< 0.001	0.53	< 0.001
General practice	0.34	< 0.001	0.46	< 0.001	0.38	< 0.001	0.44	< 0.001	0.26	< 0.001
Out patient clinic	0.37	< 0.001	0.32	< 0.001	0.27	< 0.001	0.31	< 0.001	0.30	< 0.001

* Out of 2,810 respondents, 114 respondents were excluded due to missing completed items, resulting in 2,696 respondents, who completed more than ten items



and SRMR appeared below upper the upper limits, 0.10 and 0.08, respectively, for acceptable fits [24, 25]. The evaluation of the PACIC questionnaire in this setting is thus consistent with a similar evaluation in Maindal [8] and surpasses it with respect to inter-item and item-rest correlation. In general, it is rather difficult to fit CFA models to PACIC evaluations. None of the studies we have encountered in our literature research fulfilled the criteria for a good fit i.e. non-significant χ^2 , CFI, TLI > 0.95, and RMSEA < 0.05 (e.g. [1, 8, 26, 27].) In summary, while challenges persist in the form of floor effects for several items, the data characteristics indicate that PACIC is a relevant tool for assessing patient-centred chronic care in Danish settings.

Limitations

The statistical significance should be interpreted cautiously due to many investigated effects; the significance levels of $p < 0.01$ possibly arose from numerous comparisons. However, some of these findings with significant but high p -values are supported by patterns across the subscales, e.g. the effect of physical training in the municipality. Statistical significances with $p < 0.001$ are not questionable as this corresponds to a full Bonferroni correction. In the present study, there is a low response rate of 54.1%. In our analyses of non-responders vs. responders we found that the non-responder group present a picture of people who, on average, are more

socio-economically deprived in their lives than the responders and therefore more unlikely to complete the questionnaire. With the high rate of non-responders this may cause the results to be biased towards a more socio-economically well-functioning population.

Conclusions

In conclusion, the PACIC questionnaire has potential as a tool measuring regular assessment of patient's perspectives on the quality of care in a diabetes population. The PACIC questionnaire can document areas that need to be improved in the three healthcare organizations; hospitals, communities, and general practice offices. Factors important to a high quality of perceived care are age lower than 60 years, being a woman, having a hospital outpatient clinic as the primary organization. Also participating in a rehabilitation program are associated with perceived quality of care. Results indicate that PACIC is a relevant tool for assessing patient-centred chronic care in a Danish setting. Although, the validation of the PACIC scale in a Danish context showed an acceptable ceiling effect and inter-item correlation, but floor effects persist for several items.

Abbreviations

PACIC: The Patient Assessment of Chronic Illness Care.; CCM: The Chronic Care Model; GP: General practitioner; EM: The expectation-maximisation; RMSEA: The Root Mean Square Error of Approximation; SRMR: The Standardized Root Mean Square Residual; TLI: The Tucker-Lewis Index; CFI: The Comparative Fit Index

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12913-021-07051-6>.

Additional file 1: Supplementary figure S1. The twenty-item Patient Assessment of Chronic Illness Care (PACIC) questionnaire

Acknowledgements

We would like to thank Judith Schaefer, a developer of the PACIC for assisting in updating the wording of items in the Danish version of PACIC.

Authors' contributions

AF and AS conceptualized, designed, and led the study. AS designed and analysed the data. AF, AN, AS, CG, HB, CUE, HTM, and BH interpreted the results, wrote, revised, and critically reviewed the manuscript. CG and AF drafted and performed the data collection by sending emails to the selected patients. All authors have read and approved the manuscript.

Authors' information

AF is an MD, Ph.D., Professor in Multimorbidity and chronic diseases at the Faculty of Health and Medical Sciences, Research Unit for General Practice, The University of Copenhagen, and Innovation and Research Centre for Multimorbidity, Slagelse hospital, Region Zealand, Denmark. Anne Frølich is the center manager at the Center for Multimorbidity and Chronic Illness located at Slagelse Hospital, Region Zealand, Denmark. Anne Frølich's research has a special focus on improving the organisation of treatment for patients with multimorbidity. AN is an MScPH, Ph.D. at Danish Dementia Research Centre, Rigshospitalet, University of Copenhagen, Denmark. AN is a project manager and senior researcher with organizational and professional insight into the healthcare system. CG is an MD, Ph.D., Professor, Head of Center for Diabetes, Copenhagen, Denmark. The focus of much of Charlotte Glümer's research is public health and epidemiology. HB, MPH, Ph.D., employed as a postdoc at Center for Clinical Research and Prevention, Frederiksberg Hospital, Copenhagen, Denmark. Her research area has focused on multimorbidity, health care organisation, and rehabilitation for patients with Cardiac diseases. CUE, MSc, employed at Center for Clinical Research and Prevention, Frederiksberg Hospital, Copenhagen, Denmark. CUE has authored several systematic reviews and reports about polypharmacy and multimorbidity. HTM is a Master of Public Health, Ph.D., and professor in health promotion, at the Department of Public Health, Aarhus University, Denmark. Her research area is, health promotion, prevention and behavioral medicine, primarily in diabetes, CVD risk, and other non-communicable diseases. She has been responsible or involved for the Danish translation of four questionnaires (PACIC, PAM, HLQ and HLS-EU). BH is a Master of Public Health Science, University of Copenhagen, Denmark. Her research experiences are analyses, monitoring and evaluation of health measurements based on data from questionnaires and registers. AS is an associate professor and statistician at the Technical University of Denmark, Department of Applied Mathematics and Computer Science. He has, among others, research experiences in mathematical modelling, epidemiology, biostatistics, and risk analysis.

Funding

The research was funded by the 50 Million Crown Foundation from the Danish Capital Region. The funders had no role in study design, data collection, analysis, interpretation, the decision to publish, or preparation of the manuscript.

Availability of data and materials

The data used to support the findings of this study are available from the corresponding author upon request.

Declarations

Ethics approval and consent to participate

According to The Danish National Committee on Health Research Ethics, (Sec. 14(2) of the Committee Act), no formal ethics approval is needed when using questionnaires that do not involve human biological material [28]. According to the Helsinki Declaration [29], the participants received information about the study. They were informed about the possibility to withdraw from the study; that their participation was voluntary; and that their anonymity would be maintained. After agreeing to participate, the participants provided written informed consent.

Consent for publication

Not applicable for this study.

Competing interests

The authors have no conflicts of interest to declare.

Author details

¹Section of General Practice, Department of Public Health, University of Copenhagen, 1356 K, Copenhagen, Denmark. ²Innovation and Research Centre for Multimorbidity, Slagelse Hospital, 4200 Slagelse, Denmark. ³Danish Dementia Research Centre, Rigshospitalet, University of Copenhagen, 2100 Copenhagen, Denmark. ⁴Center for Diabetes, KBH V, 1620 Copenhagen, Denmark. ⁵Danish Health Authority, Copenhagen, Denmark. ⁶Department of Public Health, Aarhus University, Aarhus, Denmark. ⁷Local Government Denmark (KL), Copenhagen, Denmark. ⁸Center for clinical Research and Prevention, Frederiksberg Hospital, Frederiksberg, Denmark. ⁹Department of Applied Mathematics and Computer Science, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark.

Received: 21 August 2020 Accepted: 17 September 2021

Published online: 09 October 2021

References

- Glasgow RE, Wagner EH, Schaefer J, Mahoney LD, Reid RJ, Greene SM. Development and Validation of the Patient Assessment of Chronic Illness Care (PACIC). *Med Care*. 2005;43:436–44.
- Avery L, Flynn D, Van Wersch A, Sniehotta FF, Trenell MI. Changing physical activity behavior in type 2 diabetes: A systematic review and meta-analysis of behavioral interventions. *Diabetes Care*. 2012;35:2681–9.
- Gregg EW, Chen H, Wagenknecht LE, Clark JM, Delahanty LM, Bantle J, et al. Association of an Intensive Lifestyle Intervention With Remission of Type 2 Diabetes. *JAMA*. 2012;308:2489.
- Asche CV, Bode B, Busk AK, Nair SR. The economic and clinical benefits of adequate insulin initiation and intensification in people with type 2 diabetes mellitus. *Diabetes Obes Metab*. 2012;14:47–57.
- Bodenheimer T, Wagner EH, Grumbach K. Improving primary care for patients with chronic illness: the chronic care model, Part 2. *JAMA*. 2002;288:1909–14.
- Bodenheimer T, Wagner EH, Grumbach K. Improving primary care for patients with chronic illness. *JAMA*. 2002;288:1775–9.
- Kuznetsov L, Simmons RK, Sandbaek A, Maingal HT. The impact of intensive multifactorial treatment on perceptions of chronic care among individuals with screen-detected diabetes: Results from the ADDITION-Denmark trial. *Int J Clin Pract*. 2015;69:466–73.
- Maingal HT, Sokolowski I, Vedsted P. Adaptation, data quality and confirmatory factor analysis of the Danish version of the PACIC questionnaire. *Eur J Public Health*. 2012;22:31–6.
- Szecsényi J, Rosemann T. German Diabetes Disease Management Programs Are Appropriate for Restructuring Care According to the Chronic Care Model. *Diabetes Care*. 2008;31:1150–4.
- Vrijhoef HJM, Berbee R, Wagner EH, Steuten LMG. Quality of integrated chronic care measured by patient survey: Identification, selection and application of most appropriate instruments. *Heal Expect*. 2009;12:417–29.
- Carstensen B, Kristensen JK, Marcussen MM, Borch-Johnsen K. The national diabetes register. *Scand J Public Health*. 2011;39:58–61.
- Robinson KM, Lau CJ, Jeppesen M, Vind AB, Glümer C. Kroniske sygdomme: hvordan opgøres kroniske sygdomme [Chronic Conditions - How to Assess Chronic Conditions]. Glostrup, Forskningscenter for Forebyggelse og Sundhed; 2011.
- Dempster AP, Laird NM, Rubin DB. Maximum Likelihood from Incomplete Data Via the EM Algorithm. 1977.
- Taber KS. The Use of Cronbach's Alpha When Developing and Reporting Research Instruments in Science Education. *Res Sci Educ*. 2018;48:1273–96.
- R: A language and environment for statistical computing. 2014. <http://www.r-project.org/>. Accessed 6 May 2019.
- Green A, Sortso C, Jensen PB, Emneus M. Incidence, morbidity, mortality, and prevalence of diabetes in Denmark, 2000–2011: Results from the Diabetes Impact Study 2013. *Clin Epidemiol*. 2015;7:421–30.
- Regionernes Kliniske Kvalitetsudviklingsprogram RKKP. 2019. <https://www.rkkp.dk/kvalitetsdatabaser/databaser/voksendiabetes/>. Accessed 12 Feb 2019.

18. Jensen VM, Rasmussen AW. Danish education registers. *Scand J Public Health*. 2011;39(7 Suppl):91–4.
19. Juul L, Rowlands G, Maindal HT. Relationships between health literacy, motivation and diet and physical activity in people with type 2 diabetes participating in peer-led support groups. *Prim Care Diabetes*. 2018;12:331–7. <https://doi.org/10.1016/j.pcd.2018.02.005>.
20. Clark LA, Watson D. Constructing Validity: Basic Issues in Objective Scale Development. *Psychol Assess*. 1995;7:309–19.
21. McDowell I. *Measuring Health: A guide to rating scales and questionnaires*. Oxford University Press; 2006.
22. Sitzia J. How valid and reliable are patient satisfaction data? An analysis of 195 studies. *Int J Qual Heal Care*. 1999;11:319–28.
23. Gatignon H. Confirmatory Factor Analysis. In: *Statistical Analysis of Management Data*. New York: Springer; 2014. pp. 55–122.
24. Hu LT, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Struct Equ Model*. 1999;6:1–55.
25. MacCallum RC, Browne MW, Sugawara HM. Power analysis and determination of sample size for covariance structure modeling. *Psychol Methods*. 1996;1:130–49.
26. Lambert S, McCusker J, Belzile E, Yaffe M, Ihejirika C, Richardson J, et al. Using confirmatory factor analysis and Rasch analysis to examine the dimensionality of The Patient Assessment of Care for Chronic Illness Care (PACIC). *Qual Life Res*. 2021;30:1503–12. doi:<https://doi.org/10.1007/s11136-020-02750-9>.
27. Fan J, McCoy RG, Ziegenfuss JY, Smith SA, Borah BJ, Deming JR, et al. Evaluating the Structure of the Patient Assessment of Chronic Illness Care (PACIC) Survey from the Patient's Perspective. *Ann Behav Med*. 2015;49:104–11.
28. The Danish National Committee on Health Research Ethics. The Danish National Committee on Health Research Ethics. 2019. <https://en.nvk.dk/how-to-notify/what-to-notify>. Accessed 19 Feb 2019.
29. Williams JR. The Declaration of Helsinki and public health. *Bull World Health Organ*. 2008;86:650–2.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

