



Single-photon superradiance: effects of the dielectric environment and accuracy of the rotating-wave approximation

Jørgensen, Mads Anders; Pandey, Devashish; Xiao, Sanshui; Leitherer-Stenger, Nicolas; Wubs, Martijn

Publication date:
2022

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Jørgensen, M. A., Pandey, D., Xiao, S., Leitherer-Stenger, N., & Wubs, M. (2022). *Single-photon superradiance: effects of the dielectric environment and accuracy of the rotating-wave approximation*. Poster session presented at 5th Nanophotonics and Micro/Nano Optics International Conference , Paris, France.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Single-photon superradiance: effects of the dielectric environment and accuracy of the rotating-wave approximation

Thursday, 27th October - 13:30: Poster Session - Poster

*Mr. Mads A. Jørgensen*¹, *Dr. Devashish Pandey*¹, *Dr. Sanshui Xiao*¹, *Dr. Nicolas Stenger*¹,
*Dr. Martijn Wubs*¹

1. Department of Electrical and Photonics Engineering, Technical University of Denmark

Introduction:

The electromagnetic environment influences the spontaneous-emission rate of a quantum emitter, but also the collective decay (superradiance and subradiance) of a collection of emitters. In a solid, dephasing due to phonons often but not always spoils the collective decay. The present study of collective decay in a 2D material close to a plasmonic surface is especially relevant for a remarkable class of lifetime-limited defect emitters in hexagonal boron nitride.

Methods:

We employ a multiple scattering theory where collective emission is described by the Green function of the macroscopic Maxwell wave equation. We calculate the propagator between various quantum emitters positioned in a multilayer medium [in preparation, 2022]. We also determine how much the propagator differs from the Green function when making the common rotating-wave approximation (RWA) [J. Phys. B 55, in press (2022)].

Results:

We consider emitters embedded in a thin high-index dielectric layer with either air or metal as a substrate. Both configurations allow for guided modes. The guided and surface-plasmon modes are shown to play a significant role in the collective decay (super- and subradiance) of emitters at all lateral separations considered. The relative orientation of the emitters w.r.t. each other and to the interfaces of the medium determine whether the guided modes enhance or inhibit collective decay. The largest effect is found for dipoles oriented along the plane. Making the RWA only affects the real part of the induced inter-emitter interactions. For free space, the RWA may lead to induced interatomic interactions that are wrong by up to a factor of two, in contrast to a scalar model where the “RWA error” even diverges in the near field.

Discussion:

Collective emission in layered systems is affected by guided modes, with the strongest effect for dipoles oriented along the planes, which is the most common orientation for emitters in 2D materials. Making the RWA in the light-matter interaction has subtle effects: it affects neither single-emitter spontaneous-emission rates, nor the collective emission rates of two identical emitters. But for three or more identical emitters or for two slightly detuned emitters, the RWA does influence the calculated collective rates.