



## Application of Lorenz-Mie Theory in Graphics

**Frisvad, Jeppe Revall**

*Published in:*  
Mie Theory 1908-2008

*Publication date:*  
2008

*Document Version*  
Early version, also known as pre-print

[Link back to DTU Orbit](#)

*Citation (APA):*  
Frisvad, J. R. (2008). Application of Lorenz-Mie Theory in Graphics. In W. Hergert, & T. Wriedt (Eds.), *Mie Theory 1908-2008: Present developments and interdisciplinary aspects of light scattering* (pp. 26-28). Universität Bremen.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Application of Lorenz-Mie Theory in Graphics

**Frisvad, J. R.**

**DTU Informatics, Technical University of Denmark, Richard Petersens Plads, Building 321, DK-2800 Kgs. Lyngby, Denmark**

This talk has two purposes: firstly to show what the Lorenz-Mie theory has been used for in computer graphics, secondly to discuss the generalisations of the theory that are available and why they are important in a graphics context. Some details are provided in the following.

In order to produce photo-realistic images that capture a digitally modelled scene, we need scattering properties for all the materials that are present in the scene. To acquire such scattering properties, one option is to measure them. Another option is to compute them from the particle contents of the materials. The second option is made possible by the Lorenz-Mie theory, and it provides us with a very flexible way of modelling how the appearance of a material changes when we change its contents. Nevertheless, the Lorenz-Mie theory has only reluctantly been adopted in graphics. In the two papers [1,2] where the theory was first considered for graphics applications, it was found to be either too complicated [1] or too restricted [2] to be useful. The problematic restrictions are that the material should consist of nearly spherical particles embedded in a non-absorbing host medium. This significantly limits the number of materials that can be modelled by the original version of the theory. Even so, the original theory has proven useful for modelling the appearance of some special materials: Callet [3] used the theory to model pigmented materials (such as paints, plastics, inks, and cosmetics which consist of pigmented particles in a transparent solvent). Jackèl and Walter [4] used it to model the atmosphere and rainbows. Nishita and Dobashi [5] used the theory to model various other materials consisting of particles in air (clouds, smoke/gas, fog/haze, snow, and sand). Most recently, we have seen a development towards generalisation of the Lorenz-Mie theory such that it becomes useful for a wider range of materials. These new developments were adapted for graphics by Frisvad et al. [6]. I will discuss this and other generalisations of the theory. Figure 1 exemplifies how we can use the Lorenz-Mie theory to compute the visual significance of each individual component in a material. Of course, we can also use it to compute the appearance of materials with different mixed particle concentrations.



Figure 1: From the paper by Frisvad et al. [6]: Rendered images of the components in milk as well as mixed concentrations. The scattering properties of the components and the milk have been computed using a new scheme for robust evaluation of the Lorenz-Mie coefficients when the host medium is absorbing. From left to right the glasses contain: Pure water, the host medium (water and dissolved vitamin B2), water and protein, water and fat, skimmed milk, regular milk, and whole milk.

## References

- [1] T. Nishita, Y. Miyawaki, and E. Nakamae, "A shading model for atmospheric scattering considering luminous intensity distribution of light sources," *Computer Graphics (Proceedings of ACM SIGGRAPH 87)* 21(4), 303–310 (July 1987).
- [2] H. Rushmeier. "Input for participating media," in *Realistic Input for Realistic Images*, ACM SIGGRAPH 95 Course Notes (1995).
- [3] P. Callet, "Pertinent data for modelling pigmented materials in realistic rendering," *Computer Graphics Forum* 15(2), 119-127 (1996).
- [4] D. Jackèl and B. Walter, "Modeling and rendering of the atmosphere using Mie-scattering," *Computer Graphics Forum* 16(4), 201-210 (1997).
- [5] T. Nishita and Y. Dobashi, "Modeling and rendering of various natural phenomena consisting of particles," In *Proceedings of Computer Graphics International 2001*, IEEE Computer Society, 149-156 (2001).
- [6] J. R. Frisvad, N. J. Christensen, and H. W. Jensen, "Computing the scattering properties of participating media using Lorenz-Mie theory," *ACM Transactions on Graphics* 26(3), Article 60 (July 2007).